Central Eyre Iron Project Environmental Impact Statement



APPENDIX Q MARINE AND COASTAL ENVIRONMENT IMPACT ASSESSMENT



COPYRIGHT

Copyright © Iron Road Limited, 2015

All rights reserved

This document and any related documentation is protected by copyright owned by Iron Road Limited. The content of this document and any related documentation may only be copied and distributed for the purposes of section 46B of the *Development Act, 1993 (SA)* and otherwise with the prior written consent of Iron Road Limited.

DISCLAIMER

Iron Road Limited has taken all reasonable steps to review the information contained in this document and to ensure its accuracy as at the date of submission. Note that:

- (a) in writing this document, Iron Road Limited has relied on information provided by specialist consultants, government agencies, and other third parties. Iron Road Limited has reviewed all information to the best of its ability but does not take responsibility for the accuracy or completeness; and
- (b) this document has been prepared for information purposes only and, to the full extent permitted by law, Iron Road Limited, in respect of all persons other than the relevant government departments, makes no representation and gives no warranty or undertaking, express or implied, in respect to the information contained herein, and does not accept responsibility and is not liable for any loss or liability whatsoever arising as a result of any person acting or refraining from acting on any information contained within it.



REPORT

CENTRAL EYRE IRON PROJECT: PORT SITE MARINE ASSESSMENT

E-F-34-RPT-0039

| Revision | Issue Date | Revision Description | Document Writer | Authorised By |
|----------|------------|-------------------------------|------------------|---------------|
| Α | 21/01/2015 | Draft for client review | Greg Barbara | N Bull |
| | | | Kingsley Griffin | |
| | | | Chris Gosling | |
| 0 | 7/04/2014 | Final | Greg Barbara | N Bull |
| 1 | 16/06/2015 | Updates based on final review | Greg Barbara | C Gosling |



Contents

| Lis | t of Ta | bles. | | 6 |
|-----|---------|-------|---|----|
| Exe | ecutive | e sum | ımary | 7 |
| 1 | Intro | oduct | tion | 9 |
| | 1.1 | Reg | ulatory context | 10 |
| | 1.1. | 1 | Commonwealth legislation | 10 |
| | 1.1. | 2 | South Australian legislation and policy | 10 |
| 2 | Met | hodo | ology | 14 |
| | 2.1 | Req | uirements of Guidelines | 14 |
| : | 2.2 | Inve | stigations Undertaken | 16 |
| | 2.2. | 1 | Bathymetry | 17 |
| | 2.2. | 2 | Hydrodynamic Environment | 17 |
| | 2.2. | 3 | Seabed Conditions | 19 |
| | 2.2. | 4 | Water Quality | 22 |
| | 2.2. | 5 | Habitats, Flora and Fauna | 24 |
| | 2.2. | 6 | Noise and Vibration | 27 |
| | 2.2. | 7 | Invasive Marine Species | 27 |
| | 2.2. | 8 | Fisheries and Aquaculture | 27 |
| : | 2.3 | App | roach to Impact and Risk Assessment | 28 |
| | 2.3. | 1 | Impact Assessment | 28 |
| | 2.3. | 2 | Risk Assessment | 29 |
| 3 | Prop | osec | Development | 31 |
| : | 3.1 | CEIF | Infrastructure | 31 |
| | 3.1. | 1 | Port Facility | 31 |
| ; | 3.2 | Port | Site Selection Process | 36 |
| | 3.2. | 1 | Existing Export Facilities | 37 |
| | 3.2. | 2 | Greenfield Export Facility Options Considered | 38 |
| | 3.3 | Mar | ine Study Area | 39 |
| 4 | Desi | ign M | leasures and Mitigation Strategies | 41 |
| | 4.1 | Site | Selection and Alignment of Infrastructure | 41 |
| | 4.2 | Con | struction Techniques | 42 |
| • | 4.3 | Stor | mwater and Runoff | 42 |
| • | 4.4 | Envi | ronmental Management Procedures | 43 |
| • | 4.5 | Nois | se | 44 |

| | 4.6 | Dus | it | 45 |
|---|-----|--------|--|-----|
| | 4.7 | Мо | nitoring | 45 |
| 5 | Phy | /sical | Environment | 47 |
| | 5.1 | Bat | hymetry | 47 |
| | 5.1 | .1 | Existing Environment and Key Findings | 47 |
| | 5.1 | .2 | Impacts and Risks: Bathymetry | 48 |
| | 5.1 | .3 | Conclusions | 49 |
| | 5.2 | Нус | drodynamic Environment | 50 |
| | 5.2 | .1 | Existing Environment and Key Findings | 50 |
| | 5.2 | .2 | Impact and Risk: Hydrodynamics | 52 |
| | 5.2 | .3 | Conclusions | 53 |
| | 5.3 | Sea | bed Conditions | 53 |
| | 5.3 | .1 | Existing Environment and Key Findings | 54 |
| | 5.3 | .2 | Impacts and Risks: Seabed conditions | 61 |
| | 5.3 | .3 | Conclusions | 63 |
| | 5.4 | Wa | ter quality | 63 |
| | 5.4 | .1 | Existing Environment and Key Findings | 63 |
| | 5.4 | .2 | Impacts and Risks: Water quality | 67 |
| | 5.4 | .3 | Conclusions | 69 |
| 6 | Bio | logica | al Environment | 70 |
| | 6.1 | Coa | astal Habitats | 70 |
| | 6.1 | .1 | Existing Environment and Key Findings | 70 |
| | 6.1 | .2 | Impacts and Risks: Coastal Habitats | 71 |
| | 6.1 | .3 | Conclusions | 93 |
| | 6.2 | Inte | ertidal Habitats | 94 |
| | 6.2 | .1 | Existing Environment and Key Findings | 94 |
| | 6.2 | .2 | Impacts and Risks: Intertidal Habitats | 95 |
| | 6.2 | .3 | Conclusions | 97 |
| | 6.3 | Sub | otidal Habitats and Flora | 98 |
| | 6.3 | .1 | Existing Environment and Key Findings | 98 |
| | 6.3 | .2 | Impacts and Risks: Subtidal habitats | 105 |
| | 6.3 | .3 | Conclusions | 109 |
| | 6.4 | Ber | nthic Fauna | 109 |
| | 6.4 | .1 | Existing Environment and Key Findings | 109 |
| | 6.4 | .2 | Risks and Impact: Benthic fauna | 113 |
| | 6.4 | .3 | Conclusions | 119 |

| 6.5 | Mar | ine Megafauna | 120 |
|------------|----------|--|-----|
| 6 | 5.5.1 | Existing Environment and Key Findings | 120 |
| 6 | 5.5.2 | Impacts and Risks: Marine Megafauna | 127 |
| 6 | 5.5.3 | Conclusions | 137 |
| 6.6 | Prot | ected Fish Species | 138 |
| ϵ | 5.6.1 | Existing Environment and Key Findings | 138 |
| ϵ | 5.6.2 | Impacts and Risks: Protected Fish species | 138 |
| ϵ | 5.6.3 | Conclusions | 160 |
| 6.7 | Nois | se and Vibration | 161 |
| 6 | 5.7.1 | Existing Environment and Key Findings | 162 |
| 6 | 5.7.2 | Impacts and Risks: Noise and Vibration | 164 |
| 6 | 5.7.3 | Conclusions | 165 |
| 6.8 | Inva | sive Marine Species | 165 |
| 6 | 5.8.1 | Existing Environment and Key Findings | 166 |
| 6 | 5.8.2 | Impacts and Risks: Invasive Marine Species | 168 |
| 6 | 5.8.3 | Conclusions | 169 |
| 6.9 | Fish | eries and Aquaculture | 170 |
| 6 | 5.9.1 | Existing Environment and Key Findings | 170 |
| 6 | 5.9.2 | Impacts and Risks: Fisheries and Aquaculture | 174 |
| 6 | 5.9.3 | Conclusions | 175 |
| 7 F | Referenc | es | 177 |
| Appe | ndix A | Bathymetry Survey Reports | 193 |
| Appe | ndix B | Sediment Analysis Results | 194 |
| B1. | Particl | e Size Distribution Lab Results | 195 |
| B2. | Chemi | cal Analysis of Sediments Laboratory Results | 196 |
| Appe | ndix C | Geochemistry of Metal Scavenging: Technical Note | 197 |
| Appe | ndix D | Water Quality Laboratory Results | 202 |
| Appe | ndix E | Infauna Sampling Results | 203 |
| Anne | ndiv E | FDRC Search Result | 204 |



List of Figures

| Figure 2-1 Diagram of oceanographic sensor array | 18 |
|--|--------|
| Figure 2-2 Video transects | 20 |
| Figure 2-3 Sediment sample locations | 21 |
| Figure 2-4 Water sampling locations within study area | 23 |
| Figure 2-5 Conceptual cross section of marine habitat zones discussed in this report | 26 |
| Figure 3-1 Port site overview | 32 |
| Figure 3-2 Offshore infrastructure | 33 |
| Figure 3-3 Proposed Port layout depicting preferred vessel approach vectors, anchorage areas and turnir | ng |
| basins | 35 |
| Figure 3-4 Marine study area | 40 |
| Figure 4-1 Example of observation and shutdown zones – wharf (Sonus 2014) | 46 |
| Figure 5-1 Baseline bed-shear stress for easterly wave conditions (Jacobs, 2014d) | 51 |
| Figure 5-2 Infrastructure scenario bed-shear stress for easterly wave conditions (Jacobs, 2014d) | 51 |
| Figure 5-3 Seabed substrate type, determined from video imagery and sediment grabs | 55 |
| Figure 5-4 Particle size distribution measured north to south across the Cape Hardy study area | 56 |
| Figure 5-5 Particle size distribution out from shore at the Cape Hardy study area (transects parallel to coa | ast at |
| increasing depth) | 57 |
| Figure 5-6 Comparison of metal concentrations in Spencer Gulf waters | 64 |
| Figure 6-1 State / National benthic habitat maps across the area | 100 |
| Figure 6-2 Study area benthic habitat types | |
| Figure 6-3 Benthic habitat map of study area developed from towed video | 103 |
| Figure 6-4 Common dolphin photographed trailing video tow survey vessel in study area | 121 |
| Figure 6-5 Known distribution, migration and aggregation areas of Humpback Whales (DEH, 2005a) | |
| Figure 6-6 Recognised aggregation areas of Blue whales in south-east Australia (DEH, 2005b) | 123 |
| Figure 6-7 Known aggregation locations for southern right whale in South Australia (DSEWPaC, 2012) | |
| Figure 6-8 Geographic distribution of foraging Australian Sea lions across South Australia (high: red, med | ium: |
| orange, low: blue) (DSEWPaC, 2013) | |
| Figure 6-9 Map of shipping routes in and adjacent to Spencer Gulf, based on received Automatic Identifi | cation |
| System data. Areas of high shipping traffic are shown in red. Map generated from | |
| http://www.marinetraffic.com/en/ | |
| Figure 6-10 Port Neill aquaculture exclusion zones and Port Neill aquaculture zone | |
| Figure 6-11 Relative catch percentage of Abalone for Western Zone Region A, taken in 2010/11 | 173 |



List of Tables

| Table 2-1 Relevant requirements of DAC guidelines | 14 |
|---|-----------|
| Table 2-2 Data sources to assess the port site environment | 16 |
| Table 2-3 Criteria for categorising residual project impacts and benefits | 28 |
| Table 2-4 CEIP Infrastructure risk matrix | 29 |
| Table 2-5 Criteria for categorising likelihood | 29 |
| Table 2-6 Criteria for categorising consequence | 30 |
| Table 3-1 Summary of port option site selection and site evaluation criteria | 37 |
| Table 5-1 Study area sediment composition: total metals summary compared with NAGD/ANZECC Guide | elines 58 |
| Table 5-2 Study area sediment composition: other components compared with NAGD/ANZECC Guidelin | es60 |
| Table 5-3 Surface water composition: summary in comparison with EPA Water Quality Guidelines | 65 |
| Table 5-4 Bottom water composition: summary in comparison with EPA Water Quality Guidelines | 65 |
| Table 5-5 Surface and bottom water nutrient and chlorophyll a summary in comparison with EPA Water | · Quality |
| Guidelines | 66 |
| Table 6-1 Bird and reptile species: likelihood of occurrence and potential impacts | 73 |
| Table 6-2 Habitat-forming marine flora likely to occur at Cape Hardy | 104 |
| Table 6-3 Esimated area of subtidal substrate affected by marine infrastructure in the study area | 106 |
| Table 6-4 Richness and density of macroinvertebrate infauna collected from the marine sediment samp | ling |
| locations within the Cape Hardy study area | 111 |
| Table 6-5 Benthic fauna of interest in the study area | 112 |
| Table 6-6 Invertebrate species: likelihood of occurrence and potential impacts | 114 |
| Table 6-7 Protected marine megafauna potentially within study area | 120 |
| Table 6-8 Marine megafauna: likelihood of occurrence and potential impacts | 128 |
| Table 6-9 Fish species: likelihood of occurrence and potential impacts | 140 |
| Table 6-10 Summary of relevant noise limits | 163 |
| Table 6-11 Separation distance required to meet requirements of noise criteria | 163 |
| | |



Executive summary

Jacobs was engaged by Iron Road Limited (Iron Road) to undertake an assessment of the physical and biological values within the marine environment at a proposed port development and ship loader facility. The proposed port is located at Cape Hardy, 7 km south west of Port Neill on the Eyre Peninsula of South Australia. The investigation of the marine environment establishes the baseline condition of physical and biological values, determines the anticipated level of impact to the existing environment, and identifies which values require management during construction and / or operation of the proposed port development to minimise impacts wherever practicable. It also identifies legislative obligations that will need to be fulfilled in order to seek approval for proposed construction and operation of the port facility.

Of the existing and proposed export facilities on the Eyre Peninsula, none have the capacity to support the requirements of Iron Roads operations. As such, it was determined that a greenfield port facility would be required to service the export of iron concentrate from Iron Roads proposed Central Eyre Iron Project. The site of the proposed port was selected based upon a multi-criteria analysis, which assessed environmental, physical and social constraints of a number of potential port sites throughout the Eyre Peninsula, and determined that Cape Hardy was the most appropriate location.

A desktop analysis was undertaken in 2011 (updated in 2013) to determine the potential presence of matters of conservation significance within the study area, as listed under the Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* along with South Australian *National Parks and Wildlife Act 1972*, *Marine Parks Act 2007*, *Native Vegetation Act 1991*, *Fisheries Management Act 2007* and *Environment Protection Act 1993*. Marine surveys were undertaken between November 2011 and August 2012, to record baseline hydrodynamic, bathymetry, water quality and seabed conditions at the study area. The presence of flora and fauna was also recorded.

The location of the proposed port, Cape Hardy, is located within the Spencer Gulf. The marine habitats found at Cape Hardy are typical of the southern Spencer Gulf, with the following key characteristics identified:

- The marine habitats at Cape Hardy were observed to be in generally healthy condition. Shallow sandy areas were generally dominated by mature seagrass meadows. Temperate rocky reef habitat was limited to shallow areas adjacent the rocky headlands and a few sub-tidal rocky ridges.
- Comparison with data from other ports in the region shows that Cape Hardy has similar or lower levels
 of metals to Spencer Gulf waters, which have high levels of metals when compared with oceanic
 waters
- There is relatively deepwater (20m) near to shore (within 800m) across the site with the deepwater areas generally displaying only sparse assemblages of invertebrates. In the deepest areas with the finest sediments, there was some evidence of the formation of microbial matting.
- Shallow sandy areas are inhabited by mature seagrass meadows of predominantly *Posidonia* spp. In deeper waters, "clumps" of invertebrates including large ascidians, sponges and bivalves grow on the silty bottom. Small areas of rocky reef exist only in intertidal areas and adjacent to the headlands within the study area. These habitats generally follow depth and sediment contours, and appear to be stable in their distribution (based on the lack of obvious habitat loss and complexity of the habitats).
- Intertidal communities at Cape Hardy are not considered unique and are comparable with habitats commonly observed in the Spencer Gulf. No rare or protected species known to exclusively occur in the intertidal zones at Cape Hardy were identified.

The iron concentrate intended for shipment by Iron Road is magnetite-based and known to be insoluble in seawater, and are therefore highly unlikely to cause measurable elevation in dissolved iron concentrations in either the water column or surrounding sediment. There are expected to be no impacts to water quality from



iron induced algal blooms as the marine waters surrounding Cape Hardy are neither high in nutrients nor low in iron as required to stimulate iron fertilisation.

The hydrodynamic model developed for the proposed port determined that the embayments of Cape Hardy are already relatively isolated from the longshore drift currents of the Spencer Gulf and that changes to the coastline from the proposed marine infrastructure would not significantly alter coastal processes (less than 1% change in sediment transport per annum). Similarly, the location of the port does not require dredging for access to water of sufficient depth, avoiding large scale impacts to the seabed.

The proposed port is located within the Port Neill Aquaculture Exclusion Zone and approximately 2 km from the Port Neill Aquaculture Zone. There are no active aquaculture leases in either zone, with historically approved leases having been withdrawn or surrendered. The port site is located within a number of areas designated for specific shellfish and finfish fisheries although no conflict with fishery operations is anticipated due to depth requirements for commercial fishing methods.

The design of the marine infrastructure and jetty at the proposed port avoids areas of dense benthic flora which dominate the northern extent of the study area. No breeding colonies or nursery grounds for marine fauna were identified within 5 km of the study area and although eastern Hooded Plover (rated as Vulnerable in South Australia and Vulnerable and Marine under the EPBC Act) were identified using the beach adjacent the proposed jetty, the sandy beach areas of the site will be preserved. The study area is not considered to be a significant shorebird habitat.

A wide range of marine megafauna may be present in the area at different times or for limited periods. Species most likely to occur in the waters around Cape Hardy include the Australian sea lions, New Zealand fur seals, bottlenose dolphins and common dolphins. One whale species, the Southern Right Whale may occur in low numbers on a seasonal basis. A range of fish species are also likely to occur at Cape Hardy including the Leafy seadragon, ornate cowfish, Port Jackson shark, magpie perch, leatherjackets and toadfish; each of which were sighted during the video survey

Piling, drilling and underwater noise represent a potential impact to marine megafauna and fish species, however this impact is for a limited duration and will be mitigated through the establishment of observation and shutdown zones (shutdown of piling/drilling equipment in the event of observing marine megafauna) and soft start procedures to allow fauna to vacate the area.

The introduction of invasive marine species was identified as the greatest risk to the marine environment during construction and operation of the proposed port. Invasive marine species could be introduced to the port site via a range of mediums, including as biofouling on vessel hulls, jack-up barge legs, anchors, anchor chains, mooring lines, internal boat compartments, sediment transported in or on vessels, or in any seawater onboard vessels / barges including ballast water, in bilge, and inside pipes or pumps. Invasive marine species can exist in low numbers or persist as cysts in an area and can rapidly increase in numbers after a disturbance to the environment or removal of competitive indigenous species. Although all vessels utilising the port site will be required to comply with the national guidelines relevant to biofouling and ballast water, the introduction of IMS remains a high risk to the marine and coastal environment.



1 Introduction

Iron Road Limited (Iron Road) is proposing to develop an ironmining and minerals processing operation near Warramboo, approximately 25 km south east of Wudinna on the Eyre Peninsula of South Australia. Significant infrastructure is required to provide the logistics chain to enable export of the iron concentrate from the proposed mine to market. The required ancillary infrastructure includes a deep-water port facility on the east coast of the Eyre Peninsula, a standard gauge railway line from the port to the mine, a water pipeline for process water supplied from a borefield, a 275 kV transmission line, and a long term employee village at Wudinna to provide accommodation for the mine site workforce. The overall project is referred to as the Central Eyre Iron Project (CEIP). The infrastructure components (excluding the proposed mine) are herein referred to as the CEIP Infrastructure.

The proposed port development is at a greenfield site, approximately 7 km south west of Port Neill in an area known as Cape Hardy. The site provides a natural deep-water location with no dredging required. The port will have capacity to export 70 Mtpa of product (including 21.5 Mtpa of iron concentrate by Iron Road). The port is designed to support Panamax and Capesize vessels, with a 1.2 km jetty structure that incorporates a tug harbor, marine offloading facility and cargo wharf. Onshore, the port facility will incorporate materials handling facilities, car parking and internal access roads, stormwater management and ancillary facilities such as an administration building, emergency services building, control room(s), warehouse, ablutions facility and crib room, laboratory and fuel storage. Temporary workforce accommodation will also be located at the port site during construction to service the port and infrastructure corridor works. The proposed port development is to support Iron Road's operations, exporting 21.5 Mtpa of iron concentrate. Any additional infrastructure or activities proposed by third party users of the port facility would be subject to a separate approvals process.

Jacobs was engaged by Iron Road to undertake an assessment of the marine and coastal environment at the proposed port development site. The objectives of this study were to characterise the existing marine and coastal environment so that marine environmental impacts and risks associated with the development of the proposed port could be assessed and appropriate mitigation measures identified.

The Minister for Planning declared the CEIP Infrastructure to constitute a 'Major Development' pursuant to Section 46 of the *Development Act 1993* on 15 August 2013 (republished 22 August 2013 and varied 29 May 2014), following which Iron Road submitted a development application in June 2014. In November 2014, the Development Assessment Commission (DAC) issued the Guidelines for the Cape Hardy deep sea port, infrastructure corridor and long term employee village (the Guidelines), establishing the requirements for the preparation of an Environmental Impact Statement (EIS). The Guidelines outlined the parameters to determine the type, extent and condition of the marine and coastal environment at the proposed port development site and to assess the impacts resulting from the development of the CEIP Infrastructure.

Jacobs has undertaken a range of desktop, field and quantitative studies of the marine and coastal environment at the proposed port development site since 2011. The findings of these investigations have been used to determine the expected and possible impacts from the construction and operation of the CEIP Infrastructure which are presented in this report.



1.1 Regulatory context

The following section provides an overview of the legislative framework relevant to the marine and coastal environment during the construction and operational phases of the project.

1.1.1 Commonwealth legislation

The Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) prescribes the Commonwealth's role in environmental assessment, biodiversity conservation, and the management of protected areas. Under the provisions of the EPBC Act, actions that are likely to have a significant impact on a matter of National Environmental Significance are identified as "controlled actions" and cannot be undertaken without referral to the Department of the Environment (DoE) for consideration and approval.

With regard to the marine and coastal environment of the CEIP Infrastructure, the EPBC Act provides specific protection for the following Protected Matters:

- Threatened species and ecological communities the EPBC Act lists threatened fauna species under
 the following categories: extinct in the wild, critically endangered, endangered, and vulnerable. Listed
 species are afforded protection due to their threatened conservation status. Threatened species
 which are known to occur or may occur within proximity to the proposed site are highlighted in
 Section 6 of this report.
- Migratory species including those listed under International Agreements such as the Japan-Australia
 Migratory Bird Agreement (JAMBA), the China-Australia Migratory Bird Agreement (CAMBA), the
 Republic of Korea-Australia Migratory Bird Agreement (ROKAMBA), or the Convention on the
 Conservation of Migratory Species of Wild Animals (the Bonn Convention). Many of the marine
 mammals that may potentially occur in the area are listed as migratory and are protected as such
 within Commonwealth waters; these species are also discussed in Section 6 of this report.
- Some species are listed as Marine under the EPBC Act and protected within Commonwealth marine areas. The Commonwealth marine area is defined as any part of the sea, including the waters, seabed and airspace that are between 3 and 200 nautical miles from the coast (that are not otherwise classified as State waters). The Spencer Gulf sits entirely within State waters and as such, Commonwealth Marine listed species are not subject to Commonwealth marine protection. Despite this, the following report assesses potential impacts to all EPBC listed species, including those not protected within State waters. Marine listed species which could occur in the study area include the following groups:
 - Sea-snakes (Families Hydrophiidae and Laticaudidae);
 - Seals, both eared and true seals (Families Otariidae and Phocidae);
 - Marine turtles (Families Cheloniidae and Dermochelyidae);
 - Seahorses, sea-dragons, pipefish and the ghost pipefish (Families Syngnathidae and Solenostomidae);
 - All bird species that occur naturally in the area; and
 - All cetaceans (whales, dolphins and porpoises).

1.1.2 South Australian legislation and policy

A number of legislative instruments that are specific to South Australia and State waters are also relevant to the marine and coastal environment of the CEIP Infrastructure during the approvals process, construction and operational phases. In general, State legislation is only relevant to State waters. The definition of state waters encompasses all waters within within 3 nautical miles from the coast (including the whole Spencer Gulf).



1.1.2.1 Development Act 1993

The *Development Act 1993* (Development Act) controls development in South Australia, establishing procedures to assess different forms of development. Part 4, Division 2 of the Development Act relates to Major development or projects and is applicable to the CEIP Infrastructure development.

The Minister for Planning declared the CEIP Infrastructure a 'major development' pursuant to Section 46 of the *Development Act 1993* on 15 August 2013 (republished 22 August 2013 and varied 29 May 2014). The DAC issued the Guidelines for the preparation of an EIS in November 2014, requiring an assessment of the relevant environmental impacts associated with the development of the CEIP Infrastructure. The EIS will be made available for public and State agency comment during the assessment period, with the final decision whether or not to approve the CEIP Infrastructure to be made by the Governor of South Australia on the advice of the Department for Planning, Transport and Infrastructure (DPTI) and the Minister for Planning.

1.1.2.2 Harbours and Navigation Act 1993

The Harbours and Navigation Act 1993 governs the safe, efficient and reliable movement of cargo vessels within South Australia. The Act also controls the efficient and effective administration and management of South Australian harbors and harbor facilities, and the safe movement of shipping within harbors and in South Australian waters. Of specific relevance to the marine environment, the Act also applies to any marine parks, and seeks to further the objectives of the Marine Parks Act 2007. To operate a port within South Australia the port area must be declared by the Minister and a Port Operators Agreement approved by DPTI who require a biosecurity plan and Oil Spill Contingency Plan to be provided by the port operator.

1.1.2.3 National Parks and Wildlife Act 1972

The National Parks and Wildlife Act 1972 (NPW Act) allows for the protection of habitat and wildlife through the establishment of parks and reserves (both on land and in State waters). It also provides for the use of wildlife through a system of permits allowing certain actions, i.e. keeping, selling, trading, harvesting, farming, hunting, and the destruction of native species.

The NPW Act assigns species to state conservation categories; Endangered (Schedule 7), Vulnerable (Schedule 8), and Rare (Schedule 9). These conservation categories are based on the categories and definitions of the International Union for Conservation of Nature (IUCN) red list categories and criteria. A number of marine species identified as potentially occurring at the port site have been defined under the NPW Act and are discussed in Section 6 of this report. Where species protected under the NPW Act are present at the port site, a permit maybe required prior to undertaking any works which may impact the species. The National Parks and Wildlife (Protected Animals – Marine Mammals) Regulations 2010 also provides guidance on how vessels should operate within proximity to marine mammals and will be relevant to vessels during construction and operational phases.

1.1.2.4 Native Vegetation Act 1991

The objectives of the *Native Vegetation Act 1991* (Native Vegetation Act) include the conservation, protection and enhancement of the native vegetation in South Australia and, in particular, remnant native vegetation, in order to prevent further loss of quantity and quality of native vegetation (critical habitat) in the State. The Native Vegetation Act defines native vegetation as "a plant or plants of a species indigenous to South Australia including a plant or plants growing in or under waters of the sea." Under the Native Vegetation Act, marine vegetation such as seagrasses, which are present in the marine study area are protected. Where removal of native vegetation is proposed, consent is required from the relevant authority.



1.1.2.5 Fisheries Management Act 2007

In addition to the management and regulation of commercial fisheries, the *Fisheries Management Act 2007* aims, to provide for the conservation and management of the aquatic resources of the State and the control of exotic aquatic organisms and disease in aquatic resources. The Act includes the prohibition of taking certain marine species such as crabs during spawning and blue groper or seadragons at any time, as well as the protection of marine fishes in Aquatic Reserves.

1.1.2.6 Marine Parks Act 2007

The Marine Parks Act 2007 covers 19 marine parks around the State with each park consisting of up to four zones - general managed use, habitat protection, sanctuary and restricted access - that afford different levels of protection to the marine environment and organisms. Within general managed use zones there is no change to the existing use of an area. Habitat protection zones are designed to protect the seafloor habitat and biodiversity within a marine park by preventing the removal or damage to, habitat or the functioning of ecosystems. Sanctuary zones, which include approximately 6% of the State coast, are areas of high conservation value and prohibit the removal or harm of plants, animals or marine products as well as prohibiting mining, trawling or recreational fishing activity, however low impact recreation such as boating, swimming or diving are permitted. The highest level of protection is in restricted access zones, which are off limits to the public. When assessing a marine area it is important to understand the existing environment and rules applicable to each park and its zones. Apart from restricted access zones it is not a legislative requirement for shipping to avoid traversing marine parks. The proposed port site is not located within a marine park.

1.1.2.7 Aquaculture Act 2001

The Aquaculture Act 2001 provides regulation of marine and inland aquaculture. The Act prescribes the classes of aquaculture permitted in the aquaculture zones as well as setting criteria for the determination of applications for licences or in the making of other decisions in relation to the zone. Aquaculture leases are discussed in more detail in Section 6.9.

1.1.2.8 Protection of Marine Waters (Prevention of Pollution from Ships) Act 1987

The Protection of Marine Waters (Prevention of Pollution from Ships) Act 1987 enacts Australia's commitment to the International Convention for the Prevention of Pollution from Ships (MARPOL) in South Australia. MARPOL governs the following areas: Annex I (Oil), Annex II (Noxious Liquid Substances), Annex III (Harmful Substances in Packaged Forms), Annex IV (Sewage), Annex V (Garbage), and Annex VI (Air Pollution). The Act is applicable to all Annex areas excluding MARPOL Annex IV (sewage), as Commonwealth legislation is applicable. All ships operating in and out of the proposed port will be bound by these commitments.

1.1.2.9 Coast Protection Act 1972

The *Coast Protection Act 1972* provides a legislative framework to make provision for the conservation and protection of the beaches and coast of South Australia. The jurisdiction of the Act includes all land:

- within the mean high water mark and the mean low water mark on the seashore at spring tides, or
- above and within 100 m of the mean high water mark, or
- below mean low water mark and within three nautical miles

The Act is implemented via the Coast Protection Board who is the primary authority and prescribed body in South Australia for the management of the coast including coastal protection and advice on coastal development. The Board has a number of functions including:



- protect the coast from erosion, damage, deterioration, pollution and misuse
- restore any part of the coast that has been subjected to erosion, damage, deterioration, pollution or misuse
- develop any part of the coast aesthetically, or to improve it for those who use and enjoy it
- manage, maintain and develop those coast facilities that the Board is responsible for
- report to the Minister where required
- carry out, or be involved in, research into the protection, restoration or development of the coast

South Australia has been divided into six Coast Protection districts, each with its own specific management plans and policies which are used to guide coastal development decisions. The port site falls within the remit of the Eyre Coast Protection District.

1.1.2.10 Natural Resources Management Act 2004

The Natural Resources Management (NRM) Act 2004 aims to achieve ecologically sustainable development in the State by establishing an integrated scheme to promote the use and management of natural resources, including coastal resources that recognises and protects the intrinsic natural values. The Act is implemented through the State NRM Council and eight regional NRM Boards who develop and review regional plans for the management of natural resources. Coastal development is reviewed against regional plans, in consultation with the Coast Protection Board.

1.1.2.11 Environment Protection Act 1993

The Environment Protection Act 1993 provides for the protection of the environment and defines the Environment Protection Authority's (EPA) functions and powers. The Act promotes ecologically sustainable development, the use of precautionary principles to minimise environmental harm, and outlines environmental obligations including responsibilities under the Environment Protection (Water Quality) Policy 2003. The Act requires polluters to bear an appropriate share of the costs and responsibilities for protecting the environment from their activities.



2 Methodology

The following section provides an overview of the methodology employed to meet the requirements of the Guidelines relevant to the marine and coastal environment. The environmental impact and risk assessment process that was undertaken is summarised based on the relevant marine and coastal environment considerations.

2.1 Requirements of Guidelines

The Guidelines establish the requirements to prepare an EIS for all components of the CEIP Infrastructure. Not all aspects of the CEIP Infrastructure and the Guidelines are relevant to the marine and coastal environment. Table 2-1 indicates the requirements outlined in the guidelines relevant to the marine and coastal environment, and where the requirement is addressed in this report.

Table 2-1 Relevant requirements of DAC guidelines

| | Item | Requirements | Discussion |
|--------------------|-------|--|-------------|
| | 4.3.1 | Describe the impacts of jetty construction and tug wharf on the foreshore, intertidal, seabed and benthic communities (especially any nursery/spawning areas), and any mitigation measures that may be used. | Section 6 |
| | 4.3.2 | Describe the coastal engineering requirements for the location, orientation and type of jetty structure. | Section 3 |
| | 4.3.3 | Describe the impacts of any blasting activities, pile driving or screw piling activities on marine communities, especially turbidity/disturbance, vibration and underwater noise on vulnerable or sensitive receptors (including marine mammals) and any mitigation methods that may be used. | Section 6.7 |
| | 4.3.4 | Describe the design and operational measures to protect water quality and prevent stormwater and other run-off from the site affecting the coastal and marine environment, during both construction and operation. | Section 4 |
| Marine | 4.3.5 | Describe the impact of any incidental concentrate spillage and dust emissions (point source and fugitive) during ship loading operations on the marine environment, especially water quality. | Section 5.4 |
| Coastal and Marine | 4.3.6 | Describe how ship loading operations will minimise incidental concentrate spillage and dust emissions (point source and fugitive) during loading operations to avoid causing harm to marine or coastal flora or fauna species, and any mitigation measures that may be used. | Section 4 |
| ö | 4.3.7 | Describe the potential impacts of increased shipping traffic and activities in the Spencer Gulf from offshore anchoring, transhipment or pilotage (especially on marine fauna, water quality, recreational activities and amenity), including effects on commercial and recreational fishing and aquaculture. | Section 6 |
| | 4.3.8 | Describe how marine pests on the jetty will be monitored and managed. Detail the response procedure that will be followed in the event of a new pest record. | Section 6.8 |
| | 4.3.9 | Investigate the sedimentary profiles in the area of construction and associated ship docking/manoeuvring areas, to determine if there are risks from the exposure of fine sediments or clays that would impact adversely on water quality (turbidity) and contribute to the production of sediment plumes in the region. | Section 5.4 |



| | Item | Requirements | Discussion |
|----------------------------|--------|---|-------------|
| | 4.3.10 | Detail measures to protect nearby beach and/or rocky foreshore areas during and after construction, including potential marine and terrestrial protection areas or associated buffers. | Section 4 |
| | 4.3.11 | Describe existing sand movement and water flow characteristics through and around the jetty structure area, to identify any possible changes to beach profiles or sedimentation on sensitive flora and fauna, and to determine sand management requirements. | Section 5.2 |
| | 4.3.12 | Quantify and detail the extent, condition and significance of native vegetation (individual species and communities) that currently exist at the Cape Hardy site | Section 6 |
| ne) | 4.3.13 | Quantify and detail the extent, condition and significance of native vegetation (individual species and communities) that may need to be cleared or disturbed during construction and the ability of communities or individual species to recover, regenerate or be rehabilitated. | Section 6 |
| Native Vegetation (Marine) | 4.3.14 | Describe measures to deliver any significant environmental benefit that is required by the Native Vegetation Act 1991. Identify measures to minimise and mitigate vegetation clearance, including incorporating any remnant stands in the layout design, and to compensate for any loss of native vegetation and habitat. | Section 6 |
| Native Ve | 4.3.15 | Describe strategies to manage and monitor invasive weed species to protect terrestrial (particularly the Hambidge Wilderness Protection Area), coastal and marine species. | Section 6.8 |
| | 4.3.16 | Identify impact avoidance, minimisation and mitigation measures and their effectiveness. | Section 4 |
| | 4.3.17 | Describe how the proposal is not inconsistent with any relevant EPBC Act guidelines, conservation advice and/or recovery plans. For instance, the conservation advice for the Eyre Peninsula Blue Gum (Eucalyptus petiolaris) Woodland Threatened Ecological Communities | Section 6 |
| Native Fauna (Marine) | 4.3.18 | Quantify and detail the extent, condition and significance of native fauna (individual species and communities) that currently exist at the Cape Hardy site, the infrastructure corridors and the long term employee village. Identify sensitive receptors (i.e. species or lifehistory stages with particular sensitivity to construction or operational processes). | Section 6 |
| | 4.3.19 | Quantify and detail the extent, condition and significance of potential native fauna habitat loss or disturbance during construction and operation and the ability of communities and individual species to recover, especially for resident or migratory shore birds, and Threatened, Endangered and Protected Species (TEPS) under the EPBC Act and the South Australian National Parks & Wildlife Act 1972 (NP&W). | Section 6 |
| īţi | 4.3.20 | Describe the measures taken to address displaced native fauna (if any). | Section 4 |
| N | 4.3.21 | Detail appropriate buffer distances that will be required between the proposed development and TEPS, including feeding areas, nesting sites and roosting sites. | Section 4 |
| | 4.3.22 | Outline the effect of light pollution, noise emissions and vibrations on TEPS (including those listed under the EPBC and NPW Act's) and how these will be managed. | Section 6 |



| Item | Requirements | Discussion |
|--------|---|-------------|
| 4.3.23 | Describe the impacts of introduced species, especially vermin and nuisance species that can be attracted to port facilities. | Section 6.8 |
| 4.3.24 | Consider the potential cumulative impacts on marine fauna as a result of the proposed development and other planned or existing port facilities in the Spencer Gulf region. | Section 6 |
| 4.3.25 | Identify impact avoidance, minimisation, mitigation and offset (where appropriate) measures and their effectiveness. | Section 4 |
| 4.3.26 | Describe how the proposal is not inconsistent with any relevant EPBC Act guidelines, conservation advice and/or recovery plans. For instance, the recovery plan for the endangered Southern Right Whale | Section 6 |

2.2 Investigations Undertaken

A comprehensive suite of nine environmental characteristics were assessed to provide an overview of the existing environment at the proposed port site to provide a basis for the subsequent environmental impact assessment. The methods used to identify environmental values and conduct the impact assessment ranged from desktop literature review, to in-field survey or sampling and detailed modelling based upon existing or collected data. The environmental parameters and methods of investigation applied to each are summarised in Table 2-2 below. A detailed overview of the methods employed in assessing existing values and impacts to the marine and coastal environment is outlined in the following Section.

Table 2-2 Data sources to assess the port site environment

| Environmental Characteristic | | Data Sources and study method |
|-------------------------------------|-----------------------------|--|
| | Bathymetry | Marine navigation charts from the Australian Hydrographic Office (AHO) Personal communications with Captain Walter Ferrao (DPTI) High-resolution site specific bathymetric survey (Hydro Survey 2012) |
| Physical Environment | Hydrodynamic Environment | Available data and literature from relevant local authorities, including the CSIRO, SARDI and BOM Port Spencer marine baseline quantitative surveys (Golder Associates, 2012) Site-specific surveys including Acoustic Doppler Current Profiling (ADCP) and temperature, salinity, tide and wave climate logging Hydrodynamic and wave modelling (Jacobs 2014d) |
| | Seabed Conditions | Towed video survey of seabed and mapping of seabed features Seabed sediment sampling to ascertain particle size distribution, and chemical analysis and detection of contaminants Cape Hardy geophysical surveys including side scan sonar, subbottom profiling, continuous marine seismic refraction and geotechnical borehole calibrations (MES, 2012) |
| | Water Quality | Publically available published literature including data from the EPA Port Spencer marine baseline quantitative surveys (Golder Associates, 2012) Water sampling and analysis of surface and bottom waters |



| Envi | ronmental Characteristic | Data Sources and study method |
|------------------------|--|---|
| Biological Environment | Marine and Benthic Habitats and Flora | Publically available published literature National and State benthic habitat map (Nature Maps) EPBC Protected Matters database Biological Database of South Australia Atlas of Living Australia online database Towed video benthic habitat survey |
| | Benthic Fauna, Fish Species and Marine Megafauna | Publically available published literature EPBC Protected Matters database National and State benthic habitat map Atlas of Living Australia online database Towed video benthic habitat survey Incidental sightings during benthic habitat surveys |
| | Noise and vibration | Underwater noise modelling report (Sonus, 2014) Marine fauna species noise threshold based on existing published literature |
| | Fisheries and Aquaculture | Aquaculture Zones – Port Neill Policy 2008 PIRSA Aquaculture Public Register Publically available published literature – SARDI fisheries reports Incidental sightings during benthic habitat surveys |
| | Invasive Marine Species | Global Invasive Species Database Australian Government Department of Agriculture resources (including interactive map of known IMS) Publically available published literature. EPBC Protected Matters Database Biological Database of South Australia Atlas of Living Australia online database Opportunistically during review of video tow survey (none identified) |

2.2.1 Bathymetry

In addition to a desktop review of the bathymetry of the site, a detailed site survey was undertaken. The bathymetric survey was completed by Hydro Survey Australia (Flinders Ports), a registered hydrographical surveyor. Swath coastal survey vessels covered all areas between the shallowest safely navigable depth along the coast and the outer extents of the study area (Figure 3-4). Data was analysed by Hydro Survey and utilised in the production of nautical charts referring to the Australian Height Datum (AHD) and offset for LAT using tide data collected over a complete lunar cycle at Tumby Bay and calibrated by the National Tide Centre (NTC).

2.2.2 Hydrodynamic Environment

In order to understand the local characteristics of tidal fluctuations, currents and stratification at Cape Hardy, site-specific oceanographic surveys were undertaken. Jacobs commissioned oceanographic contractors ASR Ltd and HOV Environment Ltd in conjunction with Dive Connect Ltd to deploy an array of oceanographic sensors at the site. The sensor array included an Acoustic Doppler Current Profiler (ADCP), temperature and salinity probes, the setup of which is indicated in Figure 2-1 (below). The ADCP logged current flow, current direction and wave height, whilst the temperature and salinity probes characterised water column mixing and stratification.

The oceanographic monitoring arrays were deployed for a total of six months of data collection within the study area, over a period beginning on 31 January 2012 with final retrieval of the equipment on 22 September 2012. The data collection covered summer, autumn and winter into early spring of 2012. There were periods of no data collection when the array was retrieved to download data, replace batteries and recalibrate the instruments.

Hydro Survey Australia deployed a tide gauge at a site approximately 2 km south of Cape Hardy. The tide gauge recorded 34 days of tide level data. The NTC undertook an analysis of this data to produce tidal planes at Cape Hardy, which are summarised in the Hydrodynamic Modelling report (Jacobs, 2014d).

On completion of the ADCP deployment, Jacobs (2014d) developed hydrodynamic modelling that included a wave model, bed shear stress analysis and a sediment transport model, to describe changes in sediment movement and coastal waves that may occur with the construction and operation of the proposed port. This hydrodynamic modelling aimed to predict the effects of the proposed wharf facility and vessel moorings within the study area, and the geographical extent of any impact on coastal processes. When coupled to water quality and sediment sampling data, the hydrodynamic modelling aimed to determine sediment movement during construction as well as potential changes to coastal sand movement due to the introduction of physical structures.




Figure 2-1 Diagram of oceanographic sensor array



2.2.3 Seabed Conditions

Visual observations of seabed geomorphology were made from a towed video survey on a grid of transects that extended 10 km along the coast out to a depth of over 20 m (Figure 2-2). Altogether, transects totalling 47.5 km in length were surveyed in the study area. A small, lightweight, high definition video camera was deployed from the survey vessel at 90 degrees from the seafloor to achieve the optimum view of the seabed and towed at a constant speed and height above the seabed. The video footage was recorded to two hard drives (one as a back-up) for post-field processing and analysis of substrate morphology. Images from the video were geo-referenced with the latitude and longitude coordinates of the vessel's position using a differential global positioning system (DGPS) mounted on the survey vessel, allowing for the production of geo-referenced seabed maps.

Substrate morphology was classified and described for all transects from the video footage. All transect footage was reviewed and stills of bare or visible substrate extracted for detailed review and classification. Substrate was visually classified as silt, sand or reef. Layers were created using the Kriging function in the interpolation tool in ArcMap (v10). This function creates a raster surface from point data. Outputs were transformed into vector data layers and draft maps reviewed by the video analysts. Where necessary the maps were manually adjusted. Additional polygon layers were created to ensure features that were rarely recorded along each transect were included on the maps as the interpolation tool smooths data to produce the layers and consequently can dropout data.

The sampling design included 16 sediment sample locations in a strategic grid pattern across the study area (Figure 2-3). A boat-deployed Petite Ponar grab sampler was utilised for sampling the sediment, however due to site A6 having very dense seagrass coverage and matting remote sediment sampling was not possible. Particle size distribution (PSD) samples were taken from a total of 15 sites (Figure 2-3). Samples for PSD analysis were stored in sealed plastic jars and chilled before transport to the University of South Australia for analysis using a Mastersizer 2000 laser diffractometer.

Sediment chemical composition was also sampled at the aforementioned 15 sites (Figure 2-3). Samples were stored in glass containers, chilled, and sent to the laboratory for analyses of arsenic (As), barium (Ba), beryllium (Be), cadmium (Cd), chromium (Cr), copper (Cu), cobalt (Co), iron (Fe), lead (Pb), manganese (Mn), mercury (Hg), nickel (Ni), vanadium (V) and zinc (Zn), organochlorin pesticides, total recoverable hydrocarbons, phosphorus organophosphorus pesticides, total petroleum hydrocarbons, benzene, toluene, ethylbenzene, xylenes, naphthalene (BTEXN) and tributylin. All analysis of sediment chemical composition was completed by ALS Environmental Laboratories and results presented as either the measured values for each analyte or as the level of detection (LOD) for analytes below the LOD of the instrumentation.



Figure 2-2 Video transects




Figure 2-3 Sediment sample locations



Sediment sampling was intended to provide a snapshot of sediment composition for description of the study area. As not all analytes tested are covered by a single relevant standard for marine sediments, the South Australian EPA standard for the production of waste derived fill (WDF) (EPA, 2013) was used in conjunction with the National Assessment Guidelines for Dredging (NAGD) (NAGD, 2009). It should be noted that the NAGD are specific for dredging projects within Commonwealth waters and the WDF are intended for the reuse of excavated sediments; and therefore not required for this project; however, these values have been used along with the Australian and New Zealand Environment Conservation Council (ANZECC) guidelines as these guidelines provide the highest level of scrutiny for marine sediments.

Although the NAGD screening levels and ANZECC guidelines relate to dredge related spoil disposal, the guidelines were developed on the basis of the effects of these substances on marine organisms and provide a conservative approach to the potential impacts of construction, which may disrupt, suspend or otherwise cause the movement of sediments at the site. The proposed development would occur within State waters and dredging or dredge disposal is not required for this project, however construction works will generate localised sediment plumes.

The detailed methodology and findings of the offshore geophysical investigation are provided in the MES (2012) report.

2.2.4 Water Quality

In addition to a desk-based review of local water quality information for the region, a detailed site survey of levels of metals and nutrients in the water column was also undertaken. Thirty water samples were collected from 1 m below the surface water at the sampling points indicated on Figure 2-4. Thirty samples were also taken at the same locations from approximately 2 m off the seabed. Samples were taken using a 2.2 L van Dorn water sampler. All water sample collection was undertaken by Jacobs' marine scientists on-board a vessel supplied and operated by Dive Connect. At all sampling locations, plastic sample containers for total metal and nutrient concentrations (total nitrogen (N), nitrite (NO_2), nitrate (NO_3), ammonia (NO_3), total Kjeldahl nitrogen (TKN) and total phosphorus (P)) were filled directly from the van Dorn, chilled in coolers and sent to ALS Environmental Laboratories within 24 hours of collection for analysis. The phytoplankton pigments chlorophyll- α and pheophytin were also analysed at the sites described above for surface water samples only.




Figure 2-4 Water sampling locations within study area



2.2.5 Habitats, Flora and Fauna

A desktop review of all available habitat data sources using a 10 km search area from Cape Hardy was undertaken including the State Habitat Map (Nature Maps) and the EPBC Protected Matters Search Tool (PMST). The EPBC Protected Matters database is maintained by the Australian Department of the Environment (DoE). This database includes likely distribution of EPBC-listed species, ecological communities and protected areas. This was supported by a search of published literature relevant to the Spencer Gulf, including the Eyre Peninsula Coastal Action Plan, state and national herbarium, and museum records.

Key steps in assessing whether the proposed development would pose significant risks to Protected Matters are assessed in accordance with DoE (2013). In determining significance of impacts, consideration was given to species and habitat abundance within and outside the impact area, sensitivity of the Protected Matter to the proposed action, and the condition and importance of habitat impacted. In addition to total species or habitat abundance, the presence of critical or core habitat areas were also considered such as:

- key breeding or foraging grounds
- population strongholds
- interruptions to migratory pathways
- exclusions to adjoining foraging grounds

Coastal habitat can provide key breeding or foraging areas for shorebirds and these areas may be protected under State or Commonwealth legislation. The widely accepted and applied approach to identifying internationally important shorebird sites throughout the world has been through the use of criteria adopted under the Ramsar Convention on Wetlands. According to this approach, a wetland is considered internationally important if it regularly supports:

- >1% of the population of one of the species migrating to Australia each year, or
- >20,000 shorebirds

A shorebird area is nationally significant if it supports:

- Regularly at least 0.1% of the population of one species of shorebird migrating to Australia each year,
 or
- Regularly >2,000 shorebirds (any mix of species), or
- At least 15 species of shorebirds

The BDBSA and Atlas of Living Australia (ALA) database were reviewed with a 50 km buffer from the study area. The larger search buffer was used for the BDBSA and ALA search in order to capture as many recorded sightings as possible and assist with determining key areas for species distribution.

Aerial imagery (visible bands only) was reviewed for the nearshore areas of the study area in order to determine the extents of intertidal reef areas. The following assumptions were made when designating reef areas:

- Areas of potential intertidal reef were identified visually (without spectral analysis)
- Where possible, bathymetry data (Hydro Survey, 2012) was used to assist in determining reef areas, basing reef extent on sudden changes in the seabed slope
- The marine geophysical report (MES, 2012) was also used to cross-check the likely extents of reef
- Reef extent was also cross-checked with towed video footage
- The shore line was based on the current watermark in the imagery



Examples of intertidal 'habitats of importance' include seagrass beds or temperate reef systems (including EPBC or NPW listed species), while 'species of interest' could include the seagrass species garweed (*Zostera muelleri ssp. mucronata*), protected under the NPW Act or leafy seadragons (*Phycodurus eques*), protected under the EPBC Act.

High resolution benthic habitat mapping was undertaken to improve understanding of the habitats present within the study area and enable a comprehensive assessment of the potential impacts. Towed video transects were conducted following a grid pattern (Figure 2-2). Tows were undertaken by Dive Connect in November 2011. A small, lightweight, high definition video camera was deployed to run parallel with the seafloor and towed at a constant speed and height above the seabed. The video footage was recorded to two hard drives (one as a back-up) for post-field processing and more detailed analysis including taxa identification. Images from the video were geo-referenced with the latitude and longitude coordinates of the vessel's position using a DGPS mounted on the survey vessel, allowing for the production of geo-referenced benthic habitat maps. Video footage from each transect was analysed to quantify percentage coverage of benthic flora and describe the dominant habitat types. As footage was time stamped and geo-referenced, transects were divided up into 10 m blocks to classify dominant habitat types for each metre (point data) of transect. Image classification first assessed key features in the field of view (i.e. bare substrate, seagrass, rocky reef and sediment type) to determine dominant habitat types. The footage was then reviewed by a second analyst to verify classifications and document secondary features such as percent coverage and density of flora or fauna assemblages and identification of taxa. The image classifications were used to map the benthic habitat within the study area using GIS techniques.

When species or habitat types of conservation significance were observed along the video transects (including EPBC or NPW listed species identified in the desktop assessment), their position and extent were logged. Examples of 'habitats of importance' include seagrass beds and temperate reef systems, while 'species of interest' include the seagrass species garweed (*Z. muelleri* ssp. *mucronata*) protected under the NPW Act and leafy seadragons, protected under the EPBC Act.

Jacobs undertook a baseline terrestrial flora and fauna survey at the proposed port to improve understanding of the coastal habitats in the study area and enable a comprehensive assessment of the potential impacts. A detailed report of port site terrestrial flora and fauna survey is provided by Jacobs (2014a). As the port site terrestrial flora and fauna survey focused on terrestrial animals (e.g. mammals, birds and reptiles) the marine assessment also reviewed the likelihood of protected marine bird and marine reptile species occurring in the study area. In addition to the terrestrial survey at the port site, opportunistic sightings of marine birds along the Cape Hardy coast made during the subtidal marine surveys were also recorded.

Information on infauna assemblages was obtained from sediment samples collected from a total of 15 sample sites across the study area in a strategic grid arrangement (Figure 2-3). Sample volumes of at least 400 ml per sample were obtained at each site and samples were sent off to separate labs for assays of sediment analysis, PSD, as well as infauna counts. All collection and sample preparation was completed by Jacobs' staff on a vessel supplied and operated by Dive Connect in November 2011. A boat-deployed Petite Ponar grab-sampler was utilised for sampling the sediment. For the infauna analysis, samples were sieved through a 500 micrometre sieve to remove fine sediment, and organisms retained on the sieve transferred to plastic jars, chilled, and fixed in 10% buffered formalin for 24 hrs before being transferred to 95% ethanol for shipment to a laboratory for identification. Benthic Australia undertook taxonomic identification and counting of the samples, to provide the total number of infauna species and abundance of infauna per cm³ per sampled



transect (for raw results see Appendix E). The towed video footage collected from the study area was also reviewed for the presence of epifauna (those animals living on the seabed) species and communities.

The distribution of many marine organisms is governed by the tide, where an organism's physiological tolerance to desiccation and exposure or submergence and salinity dictate where an organism can survive. As such, findings from investigations were separated into marine habitats based on their geographic location within the natural tidal range (refer to Figure 2-5):

- Areas above the high tide mark were deemed to be coastal habitats and classed as the 'coastal zone'
- Areas below the high tide mark but above the low tide mark are classed as the 'intertidal zone'
- Areas below the low tide mark are classed as the 'subtidal zone'




Figure 2-5 Conceptual cross section of marine habitat zones discussed in this report

The likelihood of occurrence and impact to each of the species identified by desktop and field investigations as potentially occurring within the marine or coastal environment are discussed in:

- Section 6.1 Coastal habitats, fauna and marine birds
- Section 6.2 Intertidal habitats
- Section 6.3 Subtidal habitats
- Section 6.4 Benthic fauna and invertebrate species
- Section 6.5 Marine megafauna
- Section 6.6 Fish species



2.2.6 Noise and Vibration

In addition to a desktop-based review of the current guidelines and literature for the effects of noise on marine fauna, a site-specific underwater noise modelling study was commissioned to better understand sound propagation and sensitive receivers at Cape Hardy. The site-specific underwater noise modelling and assessment was undertaken by Sonus (2014). The assessment considered the impacts of noise on marine mammals, turtles, penguins, fish and cephalopods (e.g. cuttlefish).

Noise predictions were conducted using the RAMGeo acoustic model in the AcTUP acoustic "toolbox". The RAMGeo acoustic model considers the bathymetry, profile of speed of sound in water and interaction with the different materials in the seabed. Underwater acoustic noise models calculate the transmission loss as a function of distance and frequency for a single direction. The overall noise at a distance is then calculated by subtracting the transmission loss from each of the noise sources. A detailed description of the underwater noise modelling methodology is presented in Sonus (2014).

2.2.7 Invasive Marine Species

A search of the online databases, registers and relevant published peer reviewed papers was undertaken for known IMS within Spencer Gulf and South Australia, along with a desktop assessment of threats posed by IMS to commercial fishery interests, aquaculture and the marine environment of the study area. Species considered were limited to those that have either previously been detected within South Australia or are highlighted by the National or State biosecurity authorities as a high priority species for their potential impacts and risk of establishing themselves in South Australia. No IMS were recorded during review of the video tow benthic habitat data.

2.2.8 Fisheries and Aquaculture

A search of the online PIRSA Aquaculture Public register (on 1 December 2013) for aquaculture interests around Cape Hardy was undertaken, along with a desktop assessment of registered commercial fishery interests in the area and SARDI stock assessment reports for all commercial species dating back to the 1990s. The data review included data collected during the site-specific towed video surveys as well as review of published peer reviewed papers relevant to area.



2.3 Approach to Impact and Risk Assessment

The approach to the assessment of impacts and risks to the marine environment as a result of the proposed Cape Hardy development aligns with the broader approach to impact and risk assessment employed for the whole of the CEIP Infrastructure. The environmental impact assessment for the CEIP Infrastructure focuses on the major issues associated with the project, being those impacts identified as either medium or high. The impacts identified as low or negligible have been addressed only to the extent necessary to demonstrate that they have been considered. Where identified, benefits associated with construction and operation of the project have also been identified and described, as required by the Guidelines. The identified impacts and benefits were categorised as being negligible, low, medium or high. Criteria were developed to standardise the assessment and categorisation of impacts and benefits for the project. An overview of this process is discussed in further detail below.

2.3.1 Impact Assessment

The impact assessment process recognises that, even with controls in place, normal or planned construction and operation of the project will result in changes to environmental, community and economic values. These changes may be positive (benefits) or negative (impacts). The identified impacts and benefits were categorised as being negligible, low, medium or high based on extent and duration of the predicted impacts. Criteria were developed to standardise the assessment and categorisation of impacts and benefits for the project (Table 2-3).

Table 2-3 Criteria for categorising residual project impacts and benefits

| Category | Residual Impacts | Residual Impacts | Residual Benefits |
|------------|--|---|---|
| | Legislative criteria exist | Legislative criteria do not exist | |
| Negligible | A negative change below detectable limits. | A negative change below detectable limits. OR No change to protected environmental value(s) ³ . | A positive change below detectable limits. |
| Low | Detectable negative change that is within regulatory limits/standards. | A short term (< 3 y) negative change affecting receivers located within the project area ² boundary (local receiver) only. | A short term (<3 y) positive change experienced within the project area only. |
| Medium | A periodic and temporary non- compliance of a regulatory limit/standard ¹ | A long term (>3 y) negative change affecting receivers located within the project area ² boundary (local receiver) only. OR A short term (<3 y) negative change affecting receivers outside of ⁴ the project area ² boundary, but not regionally | A long-term (>3 y) positive change experienced within the project area ² only. OR A short term (<3 y) positive change experienced outside of ⁴ the project area ² boundary (local receiver), but not regionally. |
| High | A regular or consistent non-compliance. | A negative change affecting regional receivers (Eyre Peninsula), statewide receivers or protected environmental value(s) ³ | A positive change experienced by the region (Eyre Peninsula), the state or by protected environmental value(s) ³ |

¹ Periodic and temporary impact is defined as a daily exceedance of a specified limit occurring no more than once every two weeks.

² Project area is defined as the proposed port site boundary, and the marine study area

³ Protected environmental value is an element of the environment that is afforded protection under legislation, including through licensing and permitting (e.g. listed species, native vegetation, groundwater abstraction, level of service for roads).

⁴Outside of the project area but not regionally is considered to be receiving environments within 5 km of the project area.



The impact assessment was an iterative process. Residual impacts categorised as 'negligible' or 'low' were considered as low as reasonably practicable and not warrant specific control measures, other than standard environmental management measures. 'Medium' and 'High' residual impacts required review, modification, redesign and/or control measures in order to reduce the impact to as low as reasonably practicable.

2.3.2 Risk Assessment

Faults, failures and unplanned events may occur with the potential to cause environmental impact despite best efforts to avoid or mitigate impacts. The impact assessment process has accounted for the possibility of such events occurring via an environmental risk assessment. A key distinction from the impact assessment process is that the consequences of the identified risks may or may not eventuate.

Risk criteria were developed for the project to standardise the assessment and categorisation of risks (see Table 2-4 to Table 2-6). The risk assessment process integrates approaches from the following sources:

- AS 31000: 2009 Risk Management Principles and Guidelines
- HB 203: 2012 Managing Environment Related Risk

Table 2-4 CEIP Infrastructure risk matrix

| | | | Consequences | | | | | |
|-----------|---|----------------|---------------|--------|----------|---------|--------------|--|
| | | | 1 | 2 | 3 | 4 | 5 | |
| | | | Insignificant | Minor | Moderate | Major | Catastrophic | |
| Frequency | Α | Almost certain | Low | Medium | High | Extreme | Extreme | |
| | В | Likely | Low | Medium | High | Extreme | Extreme | |
| | С | Possible | Low | Low | Medium | High | Extreme | |
| | D | Unlikely | Low | Low | Medium | High | High | |
| | E | Rare | Low | Low | Low | Medium | High | |

Table 2-5 Criteria for categorising likelihood

| Descriptor | Level | General Description | Chance p.a. | Frequency |
|----------------|-------|--|-------------|--------------|
| Almost certain | А | This event is expected to occur in most circumstances. Expected to occur at least once each year | >90% | 1/year |
| Likely | В | This event may occur in some given circumstances May occur during any given year | 20% | 1/5 years |
| Possible | С | This event might occur at some time during the project life Not likely to occur in any given year, but is possible | 5% | 1/25 years |
| Unlikely | D | This event could occur at some time Very unlikely to occur in any given year | 0.5% | 1/200 years |
| Rare | E | This event may only occur in very exceptional circumstances Examples of this have occurred historically, but is not anticipated | <0.5% | <1/200 years |

Notes: The intention is to describe the probability or frequency of an event on an annualised basis such that the impacts or exposure (risks) faced by society and the environment are recorded as those present during any given year of the life of the project, including the construction phase.

Table 2-6 Criteria for categorising consequence

| Category | Effect on behaviour of listed fauna | Effect on viability of listed species | Effect on behaviour of fauna (non-listed) | Effect on benthic flora or fauna communities | Marine Water Quality | Landform / Bathymetry |
|--------------|---|--|--|--|---|--|
| Minimal | Insignificant effect | Insignificant effect | Local short term behavioural effect | Local short term decrease in abundance of some species without reduction in local community viability | Minimal change with no significant loss of quality | Insignificant effect |
| Minor | Local short term behavioural effect | Local short term decrease in abundance with no lasting effects on local population. | Local long-term behavioural effect that does not unduly affect the ecology of the population | Local long-term decrease in abundance of some species resulting in little or no change to community structure | Local minor short term reduction or change in quality | Minor change in bathymetry within localised portions of landform |
| Moderate | Local long-term behavioural effect with no significant effects on the ecology of the species | Local long-term decrease in abundance without reduction in local population viability | Local long-term behavioural impact that significantly affects the ecology of the population | Regional long-term decrease in abundance of some species and / or local loss of some species diversity resulting in some change to the community structure | Local minor long term or widespread short term reduction or change in water quality | Widespread minor changes in bathymetry Localised major changes in bathymetry |
| Major | Local long-term behavioural effect that significantly effects the ecology of the species | Regional long-term decrease in abundance and/ or local loss resulting in reduction in regional viability | Local long term behavioural impact that significantly affects the ecology of the species | Regional long-term decrease in abundance of numerous species and / or some loss of species diversity resulting in significant changes to community structure | Widespread (regional) major short term reduction or change in water quality | Major changes in bathymetry result in effects beyond footprint |
| Catastrophic | Regional extinction of the species | Regional extinction of the species | | Regional long-term loss of numerous species resulting in the dominance of only a few species | Regional long term reduction or change in water quality | Widespread and ongoing major changes in bathymetry resulting in effects beyond footprint |



3 Proposed Development

The following section details the proposed CEIP Infrastructure (specifically the proposed port development components) and its relationship with the existing marine and coastal environment. An overview of the site selection process for the proposed port is provided, detailing the environmental, social and economic factors considered.

3.1 CEIP Infrastructure

The port, infrastructure corridor and long term employee village are ancillary infrastructure required to support the operation of the proposed CEIP mine which plans to export 21.5 Mtpa of magnetite concentrate for 25 years. The proposed port and associated shipping movements are the components of the CEIP Infrastructure most relevant to the marine and coastal environment and are discussed in detail below.

3.1.1 Port Facility

The proposed port site at Cape Hardy is located approximately 5 km south of Port Neill. The proposed port will operate 24 hours a day, seven days a week and export 21.5 Mtpa of iron concentrate. An overview of the port site is shown on Figure 3-1. The port will include:

- Rail unloading facility
- Stockpile, conveyors, and bulk materials handling facilities
- Jetty structure which commences from an abutment located on the outer end of the tug harbour breakwater and connects to the ship loading wharf
- Wharf structure for supporting the rail mounted ship loader, conveyors, roadway and services deck areas
- Module offloading facility (MOF) to facilitate the offloading of heavy modules and bulk cargo during project construction and will remain for the duration of the project
- Tug harbor to accommodate tugs and other small service vessels
- Plant and equipment workshop and facilities
- Ancillary port administration, customs and stevedoring facilities car parking and internal access roads
- Stormwater management
- Water supply and wastewater treatment facilities
- Road upgrades and realignments, including secure access gates
- Emergency services facility
- Berthing and mooring structures
- Navigational markers

A ship loader will be located at the end of the jetty. The ship loader is designed to load both Panamax and Cape size vessels on either side of the wharf. The ship loader is able to manoeuvre to reach loading hatches on any of the vessels serviced by the port facility. The ship loader's travel limits will be approximately 240 m, with a radial reach of approximately 50 m. During ship loading, the ship loader boom will be positioned over the loading hatch of the vessel. The ship loader design includes a flared telescopic chute that will extend into the loading hatch to be a short distance above the surface of the hold as the concentrate begins to load. As the hold fills, the chute will rise to maintain this short separation. This will minimise dust emitted during ship loading.



Figure 3-1 Port site overview




Figure 3-2 Offshore infrastructure



3.1.1.1 Tug Harbour

A sheltered tug facility is proposed to accommodate tugs and the other small vessels needed to operate the proposed port. The tug harbour is required for the tugs to moor safely when not in use during extreme wave conditions. Within the tug harbour, a sheltered tug jetty approximately 36 m long and 8 m wide will be constructed perpendicular to the main jetty. The tug jetty will have a steel-piled concrete deck and provide sheltered berthing for the safe mooring of harbour tugs and pilot boats. The tug jetty will be connected to the main jetty via an 18 m long, 4 m wide access way.

3.1.1.2 Module Offloading Facility

The module offloading facility will be located to the east of the main jetty and to the south of the tug harbour. Similar to the tug harbour, it will also incorporate the jetty causeway and the causeway hook in order to provide a secure berth for heavy lift ships.

The roll on, roll off berth will enable heavy modules and other project cargo to be unloaded. The cargo will be driven off the stern ramp of a heavy lift vessel using a self-propelled module transporter. The lift on / lift off wharf (approximately 20 m wide and 80 m long) will enable smaller cargo units to be lifted ashore directly onto road trucks using the heavy lift ship's onboard cranes. The single berth at the module offloading facility will provide for a range of vessel types and sizes and will be able to accommodate a heavy lift ship up to 217 m in length.

3.1.1.3 Jetty and Wharf

A co-linear jetty and wharf alignment is proposed. The jetty and wharf have been orientated to minimise the maximum vertical motion of the vessels as it aligns with the average direction of the long period, large wave height.

The required berth depth for the port facility is approximately -20.1 m Chart Datum, which allows for the draft of a fully laden Capesize vessel, vessel motion and a safety clearance of 10% under keel between the vessel hull and the sea floor. To reach the required water depth, the outer end of the wharf is approximately 1300 m from the shore line.

The jetty will provide access to the shiploading wharf for the iron concentrate, operational personnel and maintenance equipment. The proposed jetty deck is approximately 13 m wide. This provides sufficient width for a 3.6 m wide single lane roadway, jetty conveyor and walkway. The roadway will generally be used by small vehicles but will occasionally be transited by small commercial vehicles and mobile cranes. Lighting, power and telecommunications outlets will be provided at intermittent locations along the jetty.

The jetty links to the wharf deck which will be approximately 24 m wide. The roadway will generally be 4.2 m in width, except for at the end of the wharf where it will be wider to provide a vehicle turn around. The wharf deck will have a height of approximately 11.2 m AHD. This will give the shiploader enough height to be clear of the hatch of an empty Capesize vessel.

Berthing and mooring dolphins will be provided on each of the two berths. The wharf design includes a total of nine dolphins on each of the two berths. The mooring and berthing dolphins maintain berthed and moored vessels at a suitable distance from the wharf structure.



3.1.1.4 Port Approaches

The port site will receive approximately 145 vessels per year, which on average will be three vessels per week (or six vessel movements in / out). With an arrival and departure movement for each ship that visits the port site, this equates to a large vessel movement at the port facility on most days.

Ships will enter Spencer Gulf and then proceed directly to the port site at Cape Hardy. Ships would approach and depart from the port facilities via a preferred approach route from the main Spencer Gulf shipping channel. The fairway (approach and departure vector) is of sufficient natural depth so as not to require dredging. A designated anchorage area located approximately 3 km from shore is proposed for ships waiting to berth and load cargo at the port site (Figure 3-3). The fairway will be marked on charts but is not expected to be delineated in the field by navigation beacons or buoys, unless specifically required to mark navigational obstructions.

Subject to agreement with DPTI, the proposed port operating limit would be designated as the limit of jurisdiction of the port operator. It is anticipated that this will encompass the waters in the immediate area of the port site, but may not encompass the anchorage area as well. The port operator will ensure that vessels bound for Port Neill or other destinations can freely pass the port site.

Navigation lights and day marks will be provided to indicate the preferred approach route to the facility. It is proposed to include a sector light on the outer end of the wharf indicating the alignment of the loading wharf for vessels approaching from the south-east. Another set of sector lights will be positioned on the shoreward end of the wharf to indicate the limits of navigable depth and preferred alignments for vessels approaching from the north-east or at the Inner Berth.



Figure 3-3 Proposed Port layout depicting preferred vessel approach vectors, anchorage areas and turning basins



3.1.1.5 Ship Arrival

Pilotage will be provided for ships arriving at the port site. If the berth is occupied, it is expected that ships will make their way to the anchorage area without needing a pilot. The pilot will board near the anchorage area to bring the ship in along the final part of the approach route, turn the ship with tug assistance, and berth it alongside the wharf. Moored vessels would be facing offshore when at berth.

3.1.1.6 Ship Departure

At completion of loading, a marine surveyor will undertake a draft survey to calculate the total tonnage of product loaded which will then be certified in consultation with the ships master. The associated maximum loaded draft will be agreed given the state of the tide at departure, sea and swell and minimum under keel clearance conditions (allowing for a minimum 10% clearance). Large laden capsize vessels will be restricted from departing during periods of peak ebbing tidal currents.

Tugs will assist vessels to depart from the port facility by pulling the loaded ships off the berth after line release and providing escort until the ship gains sufficient speed.

3.2 Port Site Selection Process

The selection of a preferred port site to export iron concentrate from the CEIP was a three stage process (SKM, 2010). The first stage considered existing export facilities and other proposed export facilities on the Eyre Peninsula. The second stage sought to identify potential greenfield port locations which satisfied environmental, social and engineering design site selection criteria (refer to Table 3-1 for a summary of the site selection criteria). Three broad port zones which could be suitable for a deep-water port were identified from the analysis. These zones were then refined into six port site options, specifically:

- Upper West Eyre Peninsula
 - greenfield location near Elliston
- Lower West Eyre Peninsula
 - greenfield location near Drummond Point
 - greenfield location near Coles Point
- Spencer Gulf
 - greenfield location at Cape Hardy
 - greenfield location near Arno Bay
 - greenfield location near Gibbon Point

The third stage of the study compared the six port options to each other for development suitability based on a set of evaluation criteria. Like the site selection criteria used in Stage 2, the evaluation criteria addressed environmental, social, design and economic aspects of the development (refer to Table 3-1). The evaluation identified Cape Hardy as the most preferred location for a port facility for Iron Road. A summary of the evaluation of the existing export facilities and for each of the greenfield options is provided below.



Table 3-1 Summary of port option site selection and site evaluation criteria

| Category | Site selection criteria (Stage 1 – Existing facilities) | Site selection criteria (Stage 2 – greenfield port locations | Site evaluation criteria (Stage 3 – greenfield port options) | | | |
|-------------|--|--|--|--|--|--|
| Environment | Proximity to Conservation Parks Proximity to Marine Parks | Proximity to Conservation Parks Proximity to Marine Parks | Impact to known habitat of EPBC listed species Impact to native vegetation and areas of high biodiversity value | | | |
| Social | Proximity to built up urban areas | Proximity to sites of recognised heritage value Proximity to aquaculture zones | Impact to aquaculture Proximity to homesteads, dwellings, schools, churches, cemeteries and townships. Impact to cultural heritage sites or areas Impact to tourist areas Impact to visual amenity | | | |
| Design | Capacity to export an additional 21.5 Mtpa Sufficient depth of water for Capesize vessels | Proximity to deep water (a natural water depth of 20 m within reasonable proximity to the shore is preferred to accommodate Capesize ships without the need to undertake significant dredging) Exposure risk (sites with low exposure risk to the prevailing wave/swell are preferred) | Need for dredging to provide vessel access to the berth area Topographical features (avoid areas with cliffs that require extensive engineering) Suitable for exporting a minimum of 21.5Mtpa | | | |
| Economic | Certainty that the facilities would be available within Iron Road's project timeframes | Proximity to Warramboo (sites with a shorter distance to the proposed mine site are preferred) | Length of trestles to the berth area Need for breakwaters to provide adequate shelter for berth areas Ease of access for workforce and port services | | | |

3.2.1 Existing Export Facilities

There are four existing ports in the Eyre Peninsula region; Port Thevenard, Port Bonython, Port Lincoln and Whyalla. The Port of Thevenard, located 3 km from Ceduna, is managed by Flinders Ports and exported approximately 3 million tonnes of produce in 2011. Additional capacity for export is constrained by the depth of water and the site has limited room for expansion as it abuts residential and recreation areas. Due to the water depth, Thevenard is not capable of accommodating the required Capesize vessels for the CEIP.



Port Bonython, located at the head of the Spencer Gulf, is owned by the State Government and operated by Santos for the export of approximately 250,000 tonnes per annum including crude oil, naphtha, propane and butane. A proposal for an expanded operation to support the export of bulk commodities through construction of a new jetty is under consideration by the State Government. The expansion would initially provide for the export of up to 25 Mtpa, and could be expanded to be capable of export of up to 50 Mtpa. If the project goes ahead, the initial 25 Mtpa export capacity may be operational by 2018. The Port Bonython expansion is subject to Government approval and finance. Therefore this option does not provide certainty to Iron Road around when the facility could be available.

Port Lincoln is managed by Flinders Ports and exports between 1-3 Mtpa, dependent on the amount of agricultural product harvested in a given year. Expansion of Port Lincoln to meet the required tonnages is constrained by the need to move product through the township, impacts to the amenity of the community, congestion at the port as a result of existing grain export and potential land use conflicts with nearby fisheries and aquaculture industries.

The Port of Whyalla is South Australia's largest export facility, exporting approximately 6 Mtpa. The Port is owned by the State Government and operated by Arrium who recently expanded the facility to double its capacity to 12 Mtpa. However the increase capacity is already allocated to Arrium and is significantly less than the 20 Mtpa required by Iron Road for export of its iron concentreate. Further expansion of Whyalla to accommodate Capesize vessels would require large scale dredging and/or extension of the wharf facilities and would not represent an economic or environmentally viable solution.

A greenfield port facility is proposed at Port Spencer by Centrex Metals. The facility is proposed to export up to 20 Mtpa, over a four stage development. To date, only the first stage of the development (capable of exporting up to 2 Mtpa of iron ore and grain) has been designed and received Development Approval by the State Government. The anticipated timing for subsequent stages of development is currently unknown, and Iron Road are unable to rely upon the successful approval of third party projects for their own development.

The existing and proposed export facilities on the Eyre Peninsula do not provide sufficient capacity for the export of 21.5 Mtpa from the CEIP or have uncertain timing. They also do not represent an economic solution to be upgraded to meet Iron Road's requirements. Therefore a new greenfield port site is required for the CEIP.

3.2.2 Greenfield Export Facility Options Considered

As no existing port facilities were identified as suitable for Iron Road's purposes, a multi-criteria analysis was undertaken to identify a preferred greenfield site. Each of the key sites assessed as part of the multi-criteria analysis are discussed below.

3.2.2.1 Near Elliston, Upper West Eyre Peninsula

A location in the southern corner of Anxious Bay, to the north of Elliston was considered as a port option. This option was not preferred as it is subject to moderate to high wave exposure and would therefore involve the construction of an offshore breakwater. It would also require substantial piling work over potentially hard reefs. The site is in relatively close proximity (<2 km) to the Lake Newland Conservation Reserve and the Waldegrave Islands Conservation Park. The wharf and breakwater would be approximately 1 km from the Anxious Bay Aquaculture Zone. For these reasons the site ranked poorly against the environmental, social and engineering criteria.



3.2.2.2 Drummond Point and Coles Point, Lower West Eyre Peninsula

Two potential port locations were identified on the lower west Eyre Peninsula, Drummond Point and Coles Point. Both sites did not offer safe shipping navigation without substantial engineering modifications such as breakwaters or dredging and were therefore not preferred locations for a deep-water exporting port.

The Drummond Point port option site offers no natural protection from dominant waves. Therefore a breakwater to provide protection would be essential. Even with the breakwater, arrivals and departures during bad weather would be difficult and could lead to delays and increase the risk of vessel collisions or environmental damage due to navigational hazards.

The Coles Point port option would also require a significant breakwater, even though it has some indirect protection from south-west waves provided by Point Sir Isaac. The adjacent land side topography is also not ideal, with cliff faces in some locations.

3.2.2.3 Near Arno Bay, Spencer Gulf

A site approximately 4.5 km north of Arno Bay was considered as a port option. The Arno Bay option did not score well against the environmental, social and engineering criteria compared to the other site options in Spencer Gulf. The Inner Arno Bay Aquaculture zone is located approximately 3 km off the coast in this location. Although the jetty and wharf could be located outside the aquaculture zone, vessels arriving and departing from the port would have to traverse the zone. Heritage agreement areas, which protect vegetation, also adjoin the site to the north, east and west. The distance to deep water at the site is approximately 2.5 km, which is relatively long. The construction of a jetty at least 2.5 km posed too many environmental risks, along with engineering challenges. For these reasons this port option was discounted.

3.2.2.4 Gibbon Point, Spencer Gulf

The Gibbon Point port option was discounted as it is located on the southern boundary of the Franklin Harbour Marine Park. The jetty for the port would be located within the Marine Park; however the wharf and berth for vessels would be located outside of the Park. In this location there is also only a small area of land that is likely to be suitable for port infrastructure.

3.2.2.5 Cape Hardy, Spencer Gulf

The option evaluation identified the Cape Hardy port option as the most appropriate location for a port. The option scored well against the environment criteria as it is not in close proximity to a marine park or a conservation reserve and the port land area has been predominantly cleared of native vegetation for farming. The option also scored well against the engineering criteria as deep water is located relatively close to shore (approximately 1.5 km based on historical charts) and the site has some protection from ocean swells and waves. This option also scored well against the social criteria, as it is not located in close proximity to towns or registered heritage sites.

3.3 Marine Study Area

For the purpose of this baseline characterisation of the marine environment and impact assessment, a marine "study area" was identified as the area being potentially impacted by the construction and operation of the proposed port development. The marine study area surrounds the proposed port site at Cape Hardy covering approximately 2870 Ha (refer Figure 3-4). While the marine study area is assessed in this report, flora and fauna records have been sourced from a wider area (up to 50 km from the site) due to the paucity of records in the region.

JACOBS




Figure 3-4 Marine study area



4 Design Measures and Mitigation Strategies

The following section identifies measures that have been incorporated into the design of the proposed port development to minimise impacts to the marine and coastal environment. Management strategies that have been committed to by Iron Road to further minimise the risk of additional impact are also outlined.

4.1 Site Selection and Alignment of Infrastructure

A key consideration in finalising the port site selection process (Section 3.2) was the minimisation of physical and biological alterations to the existing marine environment. The selected site at Cape Hardy is readily accessible to deep water and does not require dredging (or subsequent disposal of dredge spoil) to make the site suitable for larger vessels. Similarly, no blasting within the marine environment is proposed which may alter the benthic substrate. As such, no major changes to the existing bathymetry will be required that may alter local currents. Avoiding the need for dredging and blasting will also:

- Minimise increased sedimentation and turbidity during construction which in turn minimizes potential impacts to water quality
- Minimise increased sedimentation that could smother seagrasses and other habitat areas
- Minimise noise and vibration emissions (particularly associated with blasting) within the marine environment that could impact marine fauna

The design and alignment of the jetty was refined in conjunction with hydrodynamic modelling (Jacobs, 2014d) to minimise changes to sediment deposition across the study area. The use of a pylon structure for the jetty significantly reduces potential effects to hydrodynamics as opposed to a rock armoured or earthen structure. As a result, the predicted rate of increased sediment deposition due to changes in the hydrodynamics is less than 1% per annum. This is considered to be within the range of natural fluctuation and readily incorporated by the seagrass meadows that naturally accrete sediments (refer to Section 5.2 for more information).

The vessel turning basins are located in areas of greater than 10% under-keel clearance over areas of sparse or no sea grass coverage. As such, vessel movements will minimise damage to seagrass meadows as a result of anchoring or vessel scour.

The proposed port site selection avoids known critical habitat, breeding colonies, foraging grounds and haul out areas for coastal fauna, fish species, benthic fauna and marine megafauna. Similarly, the port site is not within any known migration paths for whale. The site selection also provides an adequate separation distance to sensitive industries such as aquaculture areas and commercial fishing grounds. Conflict with existing aquaculture or fisheries operations is not anticipated.

The marine infrastructure has been aligned to minimise footprint within coastal, intertidal, subtidal and seagrass areas that may represent habitat for marine fauna. Less than 0.1 Ha of rocky intertidal shore will be affected by the proposed development. As a result, 99% of the study area's intertidal habitat and 100% of sandy intertidal areas will remain intact. Similarly, the alignment avoids areas of dense flora coverage (providing >50% seagrass coverage), with areas of dense seagrass in the north of study area outside of the construction footprint. As such, only 2.65 Ha of predominantly sparse seagrass is proposed to be cleared. Areas of sandy beach which may be utilised by protected species such as the Hooded Plover, Beach Slider or Bight Coast skink have also been avoided. This was achieved through the minimisation of the projects footprint within sandy beaches and the marine environment. The jetty, tug harbour and MOF infrastructure occupy a footprint less than 17 Ha.



Avoiding the northern parts of the study area that support dense sea grass also resulted in the avoidance of identified areas with high metal concentration, thereby minimising the likelihood of releasing metals from sediments.

As the port site is relatively isolated from a hydrodynamic perspective (refer Section 5.1), any IMS that may colonise the area would not be readily transported by currents beyond the study area and into new regions.

4.2 Construction Techniques

The rock armour for the MOF and tug harbour will be built out from a small headland (Figure 3-2) using roughly 1 m³ boulders and reclaimed clean fill from the landside construction area. By not relying on large sea based equipment for construction, there is no requirement for dredging and therefore impacts from mobilisation of sediments associated with these large vessels are removed. Similarly, avoiding the use large of sea based construction equipment will reduce the number of vessels needing to access the near shore area, minimising vessel scour, propwash and anchoring impacts. Vessel scour, propwash and anchoring can resuspend sediments and cause turbidity which would result in diminished water quality, coastal erosion and damage to habitats.

The jetty itself will be constructed in stages using a jack-up barge for impact piling to minimise disturbance to the seabed. The staged construction will also minimise long term alterations to site hydrodynamics and bathymetry to localised areas, with construction equipment in place for short durations only.

Heavy vehicle traffic, machinery movement, excavation, and construction of the MOF and tug harbour have the potential to crush habitat or disturb the flightless juvenile hooded plover, species that forage amongst the seagrass wrack, or nesting areas. To minimise adverse effects to coastal habitat, vehicle access to the beach areas will be restricted, with no storage or laydown areas for equipment on beach areas to protect the habitat of the hooded plover, beach slider and Bight Coast skink.

Construction of the marine infrastructure will be staged over an approximate 18 month period which will provide mobile animals an opportunity to vacate the immediately impacted area and re-enter the study area once construction activities have ceased.

Impact piling using a jack-up barge is the proposed method of construction and will avoid disturbance of the seabed as a result of dredging or blasting. Hollow pilings of 700mm diameter will be driven until resistance into the seabed and then filled with concrete to provide structural strength. As the hollow pilings will be easier to drive into the seabed, minimal disturbance to sediments is expected. As such, there will not be widespread disturbance of sediments that could be colonised by IMS. Disturbed areas during construction will be limited to the anchoring and securing of the jack-up barge and support vessels. As the jack-up barge legs will have only localised disturbance to the seabed this will minimise areas of disturbed substrate that could be colonised by IMS.

4.3 Stormwater and Runoff

It is expected that the majority of rainfall at the port site will infiltrate the sandy soils, rather than running off into the marine environment. Where possible, existing permeable areas covered by vegetation will be maintained and supplemented as described in Jacobs (2014a). However, run-off at the site will be altered by the construction of non-permeable infrastructure including roadways, hardstands and buildings at the site (Figure 3-1). The run-off from this infrastructure will be captured and either directed into evaporation / infiltration ponds with capacity to capture up to a 90th percentile of run-off events with any over flow from



greater events discharged into vegetated swales away from the marine environment. No wastewater discharge into the marine environment is proposed with full containment of all wastewater and sewage on site.

The capture and treatment of stormwater at the site is to eliminate the negative effects to water quality from an increase in fresh water run-off, and / or the release of potential contaminants or sediments within stormwater. This will also prevent high sediment loads, pollutants or high velocity flows from entering the marine environment or creating erosion of the coast and subsequent impacts to benthic coastal, intertidal and subtidal areas, fauna, and fish species.

4.4 Environmental Management Procedures

A CEMP and OEMP will be developed for the CEIP Infrastructure, including the proposed port. All staff and vessels will be required to comply with the CEMP / OEMP, and will be obliged to implement the following procedures:

- Vessels and staff will be prohibited from bringing their pets onto site to prevent disturbance or predation of native animals
- Procedures for managing fuel and chemical storage and distribution areas at both land and marine based sites. An OSPC will also be developed as required under the Port Operators Agreement to prevent either direct spills into the marine environment or seepage of spilled fuel into soil or groundwater subsequently entering the marine environment
- Establishment of a safety zone around piling activities to monitor the movement and behavior of marine mammals
- Establishment of shut down procedures should a marine mammal be identified within 500 m of piling activities
- Notification and demarcation of navigational hazards at the site in accordance with the Harbours and Navigation Act
- IMS control procedures incorporating:
 - enforcement of Commonwealth guidelines for ballast water management keeping vessels and equipment clean and well maintained
 - ensuring antifouling is up to date
 - reporting any suspected IMS to the Project Manager who will inform the relevant state authority
 - All vessels and staff will comply with all Commonwealth or State regulations relating to marine pests
 - The importation of all materials and goods to South Australia for the construction of this project is to comply with relevant Australian legislation

IMS control procedures will form part of their CEMP and OEMP. All vessels and equipment entering the construction site will adhere to the CEMP for procedures to mitigate importation or transfer of IMS to the site.

Spills of contaminants from vessels are often considered a major source of water pollution at port sites (OECD, 1997). In practice, such spills are often small and difficult to track. The release of potential pollutants by marine vessels is governed by Commonwealth (*Protection of the Sea Act 1981* and subsequent amendments) and International Marine Organisation (IMO) guidelines and legislation. As with any operational port all vessels accessing the wharf are required to comply with IMO Ballast water management plans (BWM) and MARPOL controls for marine pollution.



The proposed port site is not located within critical habitat, breeding colonies, foraging grounds or migration parts for marine megafauna. Despite this, a megafauna management plan will be developed that will cover both the construction and operational phases of the project to minimise impact. The megafauna management plan will include:

- A description of all threats to the megafauna species expected in the area
- A plan to monitor megafauna habitat use and behaviour, using appropriate survey techniques for mapping of potential threats to the expected whale species arising from the port construction and operations
- Identification and indication of noise sources and strategies to manage/mitigate noise impacts
- Mitigation measures to manage the impact of port operation and shipping, arising from port construction and operation activities, including underwater noise and the risk of vessel strike
- Set up of megafauna safety zone around piling activities to include an observation and exclusion zone, with the following activities undertaken prior to commencement of piling:
 - Designated megafauna observers will monitor for animals within the observation zone 30 minutes prior to commencing piling
 - Provided no animals were sighted within the 30 minutes soft start piling procedures would commence
 - Soft start piling would involve slowly ramping up of impact piling this slow build up in impact
 piling will allow animals within the wider area the opportunity to move away from the noise
 source and avoid potential noise impacts
 - The designated megafauna observers will continue to observe for megafauna within the observation zone and call for a stop works if an animal is sighted within the designated exclusion zone

A waste management plan will be developed to identify, separate and provide adequate waste disposal for all waste streams including kitchen wastes, soil (from foundations and clearance), hazardous items (e.g. sewage) and hydrocarbons. All waste will be sorted and stored within controlled contained areas until it can be removed from site by a suitable waste disposal company. Bunding will be used to prevent leaching of soluble waste or stormwater run-off carrying pollutants into drains or groundwater, and ultimately the marine environment.

Vessel speed will be restricted within the port by using tugs for manoeuvring large vessels to minimise vessel scour and changes to sediment PSD or damage to identified seagrass habitat that helps to stabilise sediment. Vessel movements will also be restricted to designated areas with sufficient depth / under keel clearance to avoid vessel scour and reduce the risk of vessels running aground. Anchorage areas for large cargo vessels are located in deep water (>20 m) and away from reefs and dense seagrass.

4.5 Noise

The following measure will be implemented during underwater piling to minimise adverse affects to marine fauna:

- Pre-start procedure The presence of marine mammals, turtles and penguins will be visually
 monitored by a suitably trained crew member for at least 30 minutes before the commencement of
 the soft start procedure.
- Soft start procedure If marine mammals, turtles or penguins have not been sighted within, or are
 unlikely to enter, the shut down zone during the pre-start procedure, the soft start procedure may
 commence in which the piling impact energy is gradually increased over a 0 minute time period.
 Visual monitoring should continue during the soft start procedure. Where visibility is poor or when it



is dark, the soft start procedure should be postponed until visual inspections of the safety zones can be made.

- Normal operation procedure If marine mammals, turtles or penguins have not been sighted within
 or are unlikely to enter the shut down or observation zone during the soft start procedure, piling may
 be increased to full impact energy. Visual monitoring should continue during normal operation. The
 soft start procedure should be repeated where piling is stopped for more than 30 minutes.
- Stand-by operations procedure If a marine mammal, turtle or a penguin is sighted within the observation zone during the soft start or normal operation procedures, the operator of the piling rig should be placed on stand-by to shut-down the piling rig. Visual monitoring should continue during stand-by operation.
- Shut-down procedure If a marine mammal, turtle or a penguin is sighted within or about to enter the shut-down zone, the piling activity should be stopped. Visual monitoring should continue and where these marine fauna are observed to move out of the shut-down zone, or it has not been seen for 30 minutes, the piling activities should recommence using the soft start procedure.

An observation zone and shutdown zone will be implemented for each stage of construction within the marine environment in accordance with Sonus (2014). An example of the observation and shut down zones for the wharf are depicted in Figure 4-1.

4.6 **Dust**

The risk of iron ore entering the marine environment from dust emissions or an accidental spill is expected to be largely mitigated via the use of covered conveyors, telescopic shiploader and veneering of stockpiles. It is reasonable to assume that there will always be some level of dust emission, which has been modelled (Jacobs 2014b) and discussed further below in Section 5.4.

4.7 Monitoring

Iron Road have committed to minimising impacts to the marine environment, marine fauna and habitats (i.e. water quality) by monitoring for impacts, including regular monitoring and establishing reference sites. Monitoring programs will be based on reference sites north and south of the port and will include:

- Ongoing monitoring of the intertidal habitat, subtidal habitat, water quality and for the presence of IMS in conjunction with State and Commonwealth biosecurity authorities
- Monitoring for changes in benthic habitat health and navigational safety
- Monitoring habitat health at the site that will be able to identify any divergences from the anticipated negligible impacts during construction and operation

JACOBS




Figure 4-1 Example of observation and shutdown zones – wharf (Sonus 2014)



5 Physical Environment

Registered hydrographic surveys were conducted to obtain accurate data of the bathymetry and to produce nautical charts of the Cape Hardy study area for navigation and engineering design purposes. Offshore geophysical investigations were undertaken to characterise the seafloor geomorphology and depth of soft sediment. Water and sediment chemistry and composition were also characterised and were used to develop a hydrodynamic model for the area (Jacobs, 2014d).

The hydrodynamic model predicts that the embayments of Cape Hardy are already relatively isolated from the longshore drift currents of the gulf, and that changes to the coastline from the proposed infrastructure would not significantly alter coastal processes (Jacobs, 2014d). In deeper water the soft bottom substrate layer becomes consistently thicker, with finer sediments, moving from a sandy environment overlying gneiss or gneiss gabbro, outcropping nearshore to a silty environment at the deepest extent of the surveyed area. The silt environments typically supported sparse mixed small algae and aggregations of invertebrates (sponges, ascidians and both motile and sessile crustaceans). The water quality and seabed conditions within the study area indicate the site is similar to other uncontaminated coastal areas within the Spencer Gulf, with generally low nutrient availability and naturally elevated levels of metals.

5.1 Bathymetry

The bathymetry of the proposed port site is of key importance as the wharf must be able to accommodate Capesize vessels at all states of the tide. Based on previous wharf development proposals and discussions with the Department of Transport Planning and Infrastructure (DPTI) Principal Advisor for Navigation Captain Walter Ferrao, the minimum Lowest Astronomical Tide (LAT) depth required for safe anchorage and passage of a Capesize vessel would be 20 m. In order to select a jetty alignment that would provide suitable depth and access for vessels, a review of the bathymetry of the area was required. Although marine charts are available for the area, the resolution of these charts is insufficient for adequate navigational and engineering design purposes so additional bathymetric surveys were undertaken.

5.1.1 Existing Environment and Key Findings

The Spencer Gulf has a length of 300 km and maximum width of 130 km, and is semi-enclosed by the Eyre Peninsula to the west and Yorke Peninsula to the east. The gulf has a typical depth of 40 to 60 m near the mouth, in the south. In the northern reaches of the gulf, and north of Cape Hardy, maximum depths reduce to 7-20 m (Bullock, 1975). The existing charts indicate that Cape Hardy would provide depths of around 20 m at a distance of 1.5 km from shore. However the resolution of the historic charts could not be used for a detailed assessment as they were last surveyed in the 1960s with a resolution of +/- 2 m.

The key findings of the detailed hydrographic survey at Cape Hardy (Appendix A) include:

- Sub-tidal bathymetry at Cape Hardy generally has low relief and is free of significant navigational hazards.
- Rocky projections within the study area are restricted to the in-shore areas close to headlands and intertidal zones.
- The 20 m LAT depth contour is generally at 750 to 1200 m from shore.

A detailed hydrographic map of the study area with local benchmarks and sounding charts is available in the full report from Hydro Survey Ltd attached as Appendix A.



5.1.2 Impacts and Risks: Bathymetry

Impacts and risks to the bathymetry at the site (and surrounds) could potentially occur as a result of:

- Construction of rock armour walls of the MOF and tug harbour
- Construction of infrastructure such as pylons, buoys and navigational markers resulting in erosion / build up of sediments
- Vessel scouring and anchoring
- Dredging
- Invasive marine species (IMS) build-up, or loss of benthic habitat resulting in erosion of sediments
- Alteration of hydrodynamic environment

An assessment of these potential impacts and risks is provided below.

5.1.2.1 Impacts

The construction of the MOF and tug harbour will permanently alter the bathymetry within the project area as they will be physical structures in the environment. This is considered to represent a medium impact on the bathymetry as the effects will be long term and confined to the footprint of the infrastructure.

The hydrodynamic modelling undertaken at the site to assess the potential impacts from development of the marine infrastructure at the Cape Hardy port predicts minimal changes to existing natural sediment dynamics (approximately 1%), and therefore on the bathymetry within the study area (Jacobs, 2014d). As no erosion of the coastline or silting of the tug harbour is expected based on the hydrodynamic model, impacts to bathymetry are considered negligible.

Operational ports have the potential to introduce invasive marine species (IMS). Some IMS such as mussels have the potential to build up biological reefs and change the bathymetry of an area, or smother benthic habitat (in the case of some algae) causing seagrass loss that could alter the bathymetry of an area by destabilising sediment and causing sediment loss. The OEMP for the port will outline monitoring and management measures to assess and control the introduction and build-up of IMS, and as such the impacts to bathymetry at the site as a result of IMS are not expected and considered to be negligible. The possibility of management measures failing is not a planned event and is therefore covered as a risk below.

As site selection and alignment of the jetty will avoid areas of dense seagrass (Section 6.3) and provide adequate depth for vessels maintenance, dredging is not required for vessel access. As such, no impacts to bathymetry are expected as a result of dredging.

5.1.2.2 Risks

Changes due to scour around jetty pilings, vessel scour or sedimentation inside the tug harbour, or other structure effects on sedimentation represent a risk to the bathymetry of the site. Vessel scour and hydrodynamic changes can not only change turbidity and sedimentation rates, but also PSD and total organic carbon (TOC). This could subsequently result in flow-on changes in benthic community composition which may influence bathymetry. The likelihood of changes to the bathymetry as a result of scour and sedimentation is considered possible and the consequence of any change is considered minor, resulting in a low risk rating.



Spills of product, loss of cargo or vessel sinking could also alter the bathymetry of the site, however these risks are considered to be low as the likelihood of occurrence of these events is unlikely (rare for vessel sinking) and the consequence on bathymetry is minor (moderate for vessel sinking).

As mentioned above operational port facilities have the potential to introduce IMS and some of these (e.g. mussels) have the potential to build up and to form biological reefs, therefore changing the bathymetry of an area. In addition, if left unmanaged, some IMS (e.g. algae) can smother benthic habitat causing seagrass loss that could alter the bathymetry of an area by destabilising sediment and causing sediment loss. Active monitoring and management of IMS will be included in the OEMP for the site. As with any operating port, the introduction of IMS is considered possible, and there is a risk that management measures fail to keep IMS from building up at the site and altering the bathymetry. The likelihood of management measures failing is considered to be possible, and the consequence to bathymetry of failure is considered to be moderate. The risk to bathymetry from IMS is therefore considered to be medium.

5.1.3 Conclusions

The hydrographic survey of Cape Hardy identified no significant navigational hazards within the study area. The hydrographic survey was used in conjunction with the ecological surveys to select an alignment for the jetty which avoids the need for dredging to achieve adequate depth. No blasting for placement of the pilings is required. As dredging or blasting will not be required to construct the marine infrastructure, impacts to the bathymetry as a result of construction are considered negligible. Scour around the site will be minimised during construction and operation by restricting vessel movements to areas of adequate depth. Hydrodynamic modelling of the site indicated negligible impacts to natural sediment movements at the site as a result of the development, largely because of the predominantly open piling structure. The development avoids seagrass areas and as a result erosion impacts to the bathymetry of the area are not anticipated.

Impacts to the bathymetry of the site are anticipated as a result of the construction of the MOF and tug harbour as these permanent structures will affect a physical change to the seabed floor. The OEMP is expected to mitigate impacts to bathymetry from IMS, either from the formation of biological reefs, or the destruction of seagrass.

The risks to bathymetry as a result of the project include unanticipated sedimentation or erosion due to scour around jetty pilings, vessel scour or sedimentation inside the tug harbour. Product spillage, loss of cargo and vessel sinkage is considered a low risk to bathymetry. Inappropriate management of IMS is considered a medium risk to bathymetry at the site



5.2 Hydrodynamic Environment

Investigation of the hydrodynamic conditions in the Spencer Gulf and the waters around Cape Hardy was undertaken to gain a better understanding of the wind, waves, tides and currents in the Cape Hardy area. Baseline investigations and field data collection were undertaken to develop and calibrate a numerical hydrodynamic model, allowing an assessment of the likely effects of the proposed port facility and moored vessels on the seabed and adjacent coastline and beaches (Jacobs, 2014d).

5.2.1 Existing Environment and Key Findings

The Spencer Gulf is a semi-enclosed body of water extending approximately 300 km inland from the Southern Ocean. It is an inverse estuary, meaning that evaporation exceeds the minimal freshwater input. This results in a trend of increasing temperature and salinity towards the upper reaches, particularly in summer months. Changes in upper gulf waters transport highly saline water out of the gulf along the eastern edge (Middleton *et al.*, 2013). Spencer Gulf also has an eccentric tidal regime, with large diurnal tidal ranges and null or neap tides roughly every fortnight (Harvey *et al.*, 1995). The tidal ranges are from 0.0 to almost 4 m (BOM, 2014).

Spring tides have a large tidal range and occur fortnightly on a full or new moon. The maximum spring tide range is over 3 m, compared to the mean tide range of approximately 1.5 m (DEH, 2003). During null or neap tides, known locally as "dodge tides", the tidal range is 0.0 m and all tidal movements cease for a period of two to three days (Lennon *et al.*, 1987). Tidal, thermohaline, wind, and wave-driven currents play a major role in the gulf's circulation with diffusion in the upper gulf driving water mixing rather than turbulence (Middleton *et al.*, 2013). In the gulf, funnelling tidal movement also affect tidal amplitude; thus the spring tide range at Port Lincoln is 2.0 m, while at Port Augusta (at the head of the tapering Spencer Gulf) it is 3.9 m. As Cape Hardy lays roughly half way up the gulf it is subject to increased tidal amplitude, especially during major storms, which can raise tides 1.0 to 1.5 m above predicted heights (Harvey and Caton, 2010; Jacobs, 2014d).

The results of the oceanographic ADCP monitoring indicated that the site is exposed to both ocean swell and wind-generated waves. The oceanic swell energy appears to approach Cape Hardy from the south-south-east while the sea waves can approach from a wider range of directions (Jacobs 2014d) as they are influenced by prevailing winds. The largest wind-generated wave heights at the site were generally associated with south easterly storms (Jacobs 2014d). Hydrodynamic modelling of the study area (Jacobs 2014d) predicts that the highest current flows and bed shear are generally experienced at the Cape Hardy point proper with a second, slightly weaker area of high current and bed shear located at a southern headland (Figure 5-1). The proposed jetty and MOF alignment sits between these two areas and modelling has predicted that the structures will have minimal impact on the current flows or bed shear across the site (Figure 5-2). The hydrodynamic modelling predicted the proposed jetty and MOF would cause no significant changes to the hydrodynamics at the site, with only minor reductions in current flow, wave energy, bed shear and sediment transport from south to north (Jacobs, 2014d). The hydrodynamic model indicated that the study area is relatively isolated from the greater gulf longshore drift processes.

JACOBS




Figure 5-1 Baseline bed-shear stress for easterly wave conditions (Jacobs, 2014d)




Figure 5-2 Infrastructure scenario bed-shear stress for easterly wave conditions (Jacobs, 2014d)



5.2.2 Impact and Risk: Hydrodynamics

As dredging will not be required during construction or operation, impacts and risks to the hydrodynamics at the site (and surrounds) could potentially occur as a result of:

- Construction of the rock armour walls of the MOF and tug harbour
- Construction of infrastructure such as pylons, buoys and navigational markers
- Potential alteration of the bathymetry in the project area via:
 - Vessel scouring and anchoring
 - Sediment build up and coastal erosion
 - IMS build-up, or loss of benthic habitat i.e. seagrass loss leading to runaway coastal erosion
- Large cargo spills or vessel run aground

An assessment of these potential impacts and risks is provided below.

5.2.2.1 Impacts

Since no dredging or associated spoil disposal is proposed as part of the construction process, no impacts to hydrodynamics are expected as a result of these activities. Piling construction is only a temporary disturbance resulting in localised, but short term changes to the hydrodynamics of the study area. As such, piling activities are expected to result in a low impact to the hydrodynamics of the site.

During construction and once constructed, the MOF and tug harbour will result in some minor but permanent alterations to the hydrodynamic processes at the site. The presence of marine structures at the Cape Hardy site, particularly the tug harbour and MOF, is expected to result in only localised and minor changes to the hydrodynamics, as detailed by the hydrodynamics modelling undertaken by Jacobs (2014d). Changes would include low-level reduction in the rates of sediment transport and bed shear which will generally follow the existing natural dynamics within the Cape Hardy area, resulting in increased sedimentation around the MOF of less than 1%. Therefore significant changes in broader coastal processes such as erosion or sediment dynamics are not expected. As such, alterations to the hydrodynamic environment as a result of the proposed development a considered to represent a medium impact; long term but localised to the marine study area.

5.2.2.2 Risks

The presence of marine structures at the Cape Hardy site, particularly the tug harbour and MOF, is expected to result in only localised changes to the hydrodynamics as discussed above and in detail by Jacobs (2014d). Changes would include low-level reduction in the rates of sediment transport and bed shear which will generally follow the existing natural dynamics within the Cape Hardy area. Therefore significant changes in coastal processes such as erosion or sediment dynamics are not expected. There is a risk that unanticipated changes to hydrodynamics could occur, for example the build-up of seagrass wrack along the MOF. OEMP monitoring would identify unpredicted build-up and mechanical removal may be undertaken if required. The risk of unanticipated changes to the hydrodynamic environment at the proposed port site are considered low, unlikely, and minor in consequence due to the ability for mechanical management if required.

Some IMS such as mussels have the potential to build up biological reefs consequently changing the bathymetry and therefore ultimately the hydrodynamics of an area. Seagrass has also been shown to influence the hydrodynamics of coastal areas by physically slowing currents. Therefore if a reef-building IMS, seagrass pest or smothering IMS such as *Caulerpa taxifolia* were introduced to the site it could cause seagrass loss, which could in turn lead to alterations in hydrodynamics and sediment dynamics. As with any operating



port the introduction of IMS is considered possible, and the consequences of such an introduction to the hydrodynamics of the site are considered moderate. Therefore, the risk of IMS to hydrodynamics is considered medium. IMS impacts and risks are further discussed in Section 6.8.

Iron ore spills, loss of cargo or vessel sinking pose risks of localised impacts to the hydrodynamics at the site. However these risks are considered low as both iron ore spills and loss of cargo are considered unlikely and expected to have only minor impacts on the hydrodynamics of the site. A vessel sinking is considered a rare event which would have a moderate consequence on the hydrodynamics of the Cape Hardy site Therefore the overall risk associated with a vessel sinking is considered to be low.

5.2.3 Conclusions

By considering the hydrodynamic environment during the design of the port infrastructure and selecting a predominantly pylon jetty, large scale changes to the hydrodynamic environment at Cape Hardy have been avoided. The alignment and location of the MOF and tug harbour mean there is no requirement for dredging, and impacts as a result of these physical structures are predicted to be insignificant to the regional hydrodynamics within Spencer Gulf (Jacobs, 2014d). Only minor, localised reductions in bed shear and current speeds in the nearshore areas of Cape Hardy are expected and these minor changes to the site hydrodynamics will not pose a significant risk to marine ecology or the marine environment.

- Localised sediment movements or seabed scour could occur from shipping activities in the nearshore
 environment. Effective implementation of a considered CEMP and OEMP would minimise potential
 effects to the marine environment.
- The increased rates of sedimentation and/or decreases in light penetration of the water column due to scouring and sediment suspension could have an effect on the ecology at Cape Hardy, depending on the sensitivity of the habitat to changes in both light penetration and in sedimentation rates. Impacts and risks to ecological habitats are discussed in more detail in Section 6

The site selection, design and targeted environmental management plans are expected to result in low to medium level impacts to the hydrodynamics at Cape Hardy. Similarly, risks to the hydrodynamic environment at Cape Hardy are expected to be low to medium

5.3 Seabed Conditions

Seabed conditions are determined by local geology, oceanographic processes, climatic conditions and ecological interactions. The coast at Cape Hardy is generally considered 'moderate energy', being within the Spencer Gulf, protected from oceanic swell and with a moderate tide range. The site is located in a long-term geologically stable area and there are no significant waterways discharging nearby to supply terrigenous sediments or nutrients.

An understanding of the seabed conditions at the site is important to inform or identify:

- preferential locations for siting and routing of jetty and berths
- the sediment type and suitability as habitat
- the potential contaminated or naturally toxic sediments with the potential to be re-mobilised during construction
- the effects of construction activities on sediment movement and transport along the coast



5.3.1 Existing Environment and Key Findings

The nature of the seabed is illustrated in Figure 5-3 and the following points are highlighted:

- The surface sediments generally changes from sandy to silty substrate at between 19-21 m of depth.
- Rocky reef was observed in a discrete, shallow-water patch close to the point of Cape Hardy. It is
 worth noting that transects avoided shallow water and rocky projections for vessel and equipment
 safety. Based on aerial imagery and the MES (2012) report it is likely that additional rocky reef
 substrate is present along the near-shore areas associated with the capes and intertidal zones.
- Toward the north of the sampled areas, sand substrate was observed to extend the full distance of the areas, with no silt observed within the northern extent of the study area.
- The sub-bottom profiling survey and seismic refraction profiles from the study area are consistent with the observed surface morphology indicating that gneiss and gneiss gabbro outcropping occurs near shore adjacent to and in line with the headlands (MES, 2012).
- The thickness of hard substrate gneiss and gneiss gabbro outcropping increases with distance from shore, as the overlying soft-bottom layers increase in thickness to around 20 m at roughly 1000 m from shore (MES, 2012). This indicates the rock outcropping occurs around the headlands and near shore with deep-water sediments dominated by soft silt layers up to 20 m thick overlying a weathered rock profile..

Detailed PSD results are presented in Appendix B and include:

- Sites to the south of the study area (Transect 1) had a greater proportion of fine sediments than those from the north (Transect 6; Figure 5-4).
- Dominant particle sizes increased closer to shore (Transect A; Figure 5-5) compared to sites sampled further offshore (Transects B, C and D), which generally had a greater portion of fine sediments.

These patterns in PSD are consistent with the visible substrate characteristics noted during the video survey, as represented in Figure 5-3.

The sediment composition is likely to be related to the bathymetry of the study area, and level of seagrass cover. Areas of no seagrass generally consist of fine sands or silt and areas of seagrass cover composed of unsorted shell grit and sands. This is consistent with many seagrass beds acting as both a trap for sediments and wave energy dissipater preventing sorting of sediments. The distribution of seagrass can also be dependent on depth as a function of light requirements in deeper waters and exposure/high wave energy in the intertidal zone. In areas of no or little seagrass cover (i.e. the intertidal zone) wave energy is able to remove fines in shallow areas and deposit the fine sediment offshore in deeper waters where less energy reaches the bottom.






Figure 5-3 Seabed substrate type, determined from video imagery and sediment grabs






Figure 5-4 Particle size distribution measured north to south across the Cape Hardy study area






Figure 5-5 Particle size distribution out from shore at the Cape Hardy study area (transects parallel to coast at increasing depth)



Results of the metal and organic contaminant analysis of the sediments are presented in Table 5-1 and Table 5-2. Results for metals were as follows:

- Across the study area, a large range of values was detected for each metal
- With the exception of As, for which two samples were above NAGD/ANZECC guidelines found in the far north eastern extent of the study area all other samples were below NAGD/ANZECC guideline levels of high and low effects on organisms (Table 5-1)
- It should be noted that Australian sediments are commonly naturally high in As and Ni (NAGD, 2009)
- Across the study area the concentrations of As, Cr and V increased with depth into the north-east
- In the most northerly and offshore area, two sites (C6 and D6) showed significantly elevated concentrations of the metals (As, Cr, Fe and V). These areas had quite coarse sediments compared to other areas at the same distance offshore, which is likely to reflect that these sites had more bioturbation compared to southern sites. It was noted during the sample collection that sediments at these sites had a strong sulfurous odour along with visible organic matter in the form of algal matting, tunicates and polychaetes. This combination of benthic fauna and anoxic sediments can be responsible for the higher concentration of metal ions via 'metal scavenging', a common product of sediments with decreased redox potential. This phenomenon is discussed further in Appendix C

Table 5-1 Study area sediment composition: total metals summary compared with NAGD/ANZECC Guidelines

| | | | ٦. | *_ | ANZ | ZECC* | | | Results | |
|----------------|-------|-------|------------|-------------------|------|-------|------|-------|----------------|--------|
| Analyte Metals | units | LOD** | SA EPA WDF | NAGD guideline | Low | High | Min | Max | Mean ± SE | Median |
| Arsenic (As) | mg/kg | 1 | 20 | 20 | 20 | 70 | 2.08 | 51.1 | 10.4 ± 3.6 | 5.9 |
| Barium (Ba) | mg/kg | 10 | 300 | n/a | n/a | n/a | 10 | 10 | 10 ± 0.0 | 10 |
| Beryllium (Be) | mg/kg | 1 | 20 | n/a | n/a | n/a | 1 | 1 | 1 ± 0.0 | 1 |
| Cadmium (Cd) | mg/kg | 0.1 | 3 | 1.5 | 1.5 | 10 | 0.1 | 0.1 | 0.1 ± 0.0 | 0.1 |
| Chromium (Cr) | mg/kg | 1 | 400 | 80 | 80 | 370 | 5.5 | 24 | 12.1 ± 1.5 | 10.5 |
| Copper (Cu) | mg/kg | 1 | 60 | 65 | 65 | 270 | 1 | 2.4 | 1.5 ± 0.1 | 1.3 |
| Cobalt (Co) | mg/kg | 0.5 | 170 | n/a | n/a | n/a | 0.5 | 0.8 | 0.6 ± 0.0 | 0.5 |
| Iron (Fe) | mg/kg | 50 | n/a | n/a | n/a | n/a | 2460 | 14000 | 5645 ± 968.7 | 4545 |
| Lead (Pb) | mg/kg | 1 | 300 | 50 | 50 | 220 | 1 | 2.9 | 1.8 ± 0.2 | 1.9 |
| Manganese (Mn) | mg/kg | 10 | 500 | n/a | n/a | n/a | 10 | 14 | 11.4 ± 0.4 | 11 |
| Mercury (Hg) | mg/kg | 0.01 | 1 | 0.15 | 0.15 | 1 | 0.01 | 0.01 | 0.01 ± 0.0 | 0.01 |
| Nickel (Ni) | mg/kg | 1 | 60 | 21 | 21 | 52 | 1 | 3.2 | 1.5 ± 0.2 | 1.2 |
| Vanadium (V) | mg/kg | 2 | n/a | n/a | n/a | n/a | 2 | 57.4 | 15.3 ± 4.6 | 10.5 |
| Zinc (Zn) | mg/kg | 1 | 200 | 200 | 200 | 410 | 1.2 | 3.6 | 2.2 ± 0.2 | 2.1 |

NOTE: *As not all analytes have SA EPA standards the NAGD/ANZECC guidelines for sediments have also been referenced as a guide.

^{**}Level of Detection (LOD) is the level of detection in the case of Cd and Hg the sampled values were below the LOD. S.E. = standard error of the mean.



The levels of organic contaminants in marine sediments within the study area were below the detection level of the laboratory analysis (Table 5-2; Appendix B). As such, results were below relevant SA EPA standards and NAGD/ANZECC guidelines.

The results of analysis for which NAGD/ANZECC guidelines exist have been presented in Table 5-2, with a complete listing of all results presented in Appendix B.

- The mean levels of organochlorin pesticides detected at Cape Hardy were consistently below the maximum concentrations for the SA EPA WDF standards and ANZECC guidelines of high levels of contamination. It should be noted that the limit of detection (LOD) of the analysis was often above the NAGD/ANZECC guidelines for low levels of contamination and therefore apparent exceedances of these guidelines may occur as a result of the limitations of the testing method. It is recognised that the ANZECC guidelines are of uncertain ecological relevance
- The mean concentration of total petroleum hydrocarbons (sum C6-C36) was 34.4 mg/kg, which is substantially lower than the NAGD screening level and below the SA EPA WDF acceptable levels in sediments.
- Similarly, levels of the BTEXNs Benzene, Toluene, Ethyl benzene and Total Xylenes were below SA EPA WDF standards
- Tributyltin, an organotin compound, was below NAGD/ANZECC guidelines



Table 5-2 Study area sediment composition: other components compared with NAGD/ANZECC Guidelines

| | | | | S | ANZE | ECC* | Sample values | | | | |
|---------------------|------------|-----------|---------------|-----------------|------|------|---------------|----------|------|--------|--------|
| | | * | ⋖ | guidelines * | | | | | | | |
| | | **007 | SA EPA WDF | ide * | | | | | | | |
| Analyte | units | _ | S > | ng | Low | High | Min | Max | Mea | n ± SE | Median |
| Organochlorin p | | OCs) | | _ | | 3 | | | | _ | |
| cis-Chlordane | mg/kg | 0.05 | 2 | 0.5 | 0.5 | 6 | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Dieldrin | mg/kg | 0.05 | 2 | 280 | 0.02 | 8 | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| 4.4`-DDE | mg/kg | 0.05 | | 2.2 | 2.2 | 27 | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Endrin | mg/kg | 0.05 | | 10 | 0.02 | 8 | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| 4.4`-DDD | mg/kg | 0.05 | | 2 | 2 | 20 | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| 4.4`-DDT | mg/kg | 0.2 | 2 | 1.6 | 1.6 | 46 | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| alpha-BHC | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Hexachlorobenzene | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| beta-BHC | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| gamma-BHC | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| delta-BHC | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Heptachlor | mg/kg | 0.05 | 2 | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Aldrin | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Heptachlor epoxide | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| trans-Chlordane | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| alpha-Endosulfan | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| beta-Endosulfan | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Endrin aldehyde | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Endosulfan sulfate | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Endrin ketone | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Methoxychlor | mg/kg | 0.2 | | | | | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| OC surrogate | % | | | | | | | | 89.3 | ± 6.8 | 91 |
| Dibromo-DDE | | | | | | | | | | | |
| Organophospho | rous pesti | cides (OF | Ps) | | | | | | | | |
| Dichlorvos | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Demeton-S-methyl | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Monocrotophos | mg/kg | 0.2 | | | | | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| Dimethoate | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Diazinon | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Chlorpyrifos-methyl | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Parathion-methyl | mg/kg | 0.2 | | | | | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| Malathion | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Fenthion | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Chlorpyrifos | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Parathion | mg/kg | 0.2 | | | | | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| Pirimphos-ethyl | mg/kg | 0.05 | | | | | < 0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Chlorfenvinphos | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Bromophos-ethyl | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Fenamiphos | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Prothiofos | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Ethion | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Carbophenothion | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| Azinphos Methyl | mg/kg | 0.05 | | | | | <0.05 | detected | 0.05 | ± 0.0 | 0.05 |
| OP surrogate DEF | % | 0.1 | | | | | | | 85.1 | ± 9.0 | 87 |



| - | | | S | ANZECC* | | | Sar | | | |
|-------------------------------------|-------------|-----------------|-----------------|---------|--------|--------|----------|-------|----------|---------|
| Analyte | units | LOD** SA EPA | guidelines * | Low | High | Min | Max | Mea | ın ± SE | Median |
| Hydrocarbons | anics | | | 2011 | 111811 | 141111 | IVIUX | 14166 | <u> </u> | Wicalan |
| Total petroleum hydrocarbons | mg/kg | 65 | 550 | | | | | 34.4 | ±10.8 | 32 |
| Total recoverable hydrocarbons | mg/kg | | | | | | | 45.8 | ±12.1 | 43 |
| BTEXN | | | | | | | | | | |
| Benzene | mg/kg | 0.2 1 | | | | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| Toluene | mg/kg | 0.2 | 1 | | | < 0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| Ethyl benzene | mg/kg | 0.2 3.3 | l | | | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| meta- & para-Xylene | mg/kg | 0.2 | | | | < 0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| ortho-Xylene | mg/kg | 0.2 | | | | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| Total Xylenes | mg/kg | 0.5 | ļ | | | < 0.5 | detected | 0.5 | ± 0.0 | 0.5 |
| Sum of BTEX | mg/kg | 0.2 | | | | <0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| Naphthalene | mg/kg | 0.2 | | | | < 0.2 | detected | 0.2 | ± 0.0 | 0.2 |
| Organotin comp | ound | | | | | | | | | |
| Tributyltin | μgSn/k g | 0.5 | 9 | 5 | 70 | 0.5 | 0.5 | 0.5 | ± 0.0 | 0.5 |
| organotin surrogate Tripropyltin | % | 0.1 | | | | | | 102 | ± 23.7 | 107 |

Note: * As not all analytes have SA EPA standards the NAGD/ANZECC guidelines for sediments have also been referenced as a guide.

In contrast to other areas within Spencer Gulf (Ward, 1987; EPA, 2004; Sinclair Knight Merz, 2010) the marine sediments within the study area are considered to be low in metals and uncontaminated.

5.3.2 Impacts and Risks: Seabed conditions

Changes to sediment structure or levels of contaminants in sediments could have an effect on the distribution and abundance of organisms, along with potentially influencing seagrass survival at the site. Sediment redistribution could potentially occur due to alterations in hydrodynamic conditions at the jetty tug harbour and MOF (Jacobs, 2014d).

Impacts to the seabed conditions could occur as a result of the following construction and operational activities, which are individually assessed in further detail below:

- Pile driving techniques to place the jetty pylons will cause suspension of fine particles or redistribution of sediments
- The jack-up barge and other construction equipment could transport contaminated sediment from other areas into the site
- Suspension or exposure of contaminated sediments could potentially occur during piling activities.
- The construction of the tug harbour and MOF could introduce contaminants and increase turbidity during construction
- Vessel scour and anchoring impacts from vessels involved in the marine infrastructure construction have the potential to alter the PSD distribution of sediments

^{**}LOD is the level of detection in all cases the sampled values were below the LOD, as per the NAGD guidelines reporting results for substances below LOD is either 'detected' or 'not detected'. S.E. = standard error of the mean.



- During construction and operation, equipment and fuel loading have the potential to spill hydrocarbons and other chemicals into the marine environment that could contaminate sediments
- Run-off from land based construction or port operation has the potential to introduce contaminants into the marine sediments at the site
- Iron ore handling and stockpiling pose the risk of generating dust which could enter the marine environment and impact on seabed sediments

5.3.2.1 Impacts

There will be a permanent change to the existing seabed as a result of introducing the tug harbour and MOF into the existing environment. Although the footprint of these structures is relatively small, the impact from these structures on the seabed conditions is considered to be medium (due to the permanent nature of the change). Hydrodynamic modelling at the site (Jacobs, 2014d) has predicted that there will be no significant departures from the natural sediment transport processes across the site.

Shipping operations are expected to modify the seabed to a small degree due to the installation of navaids, anchor drops, propwash (in the intertidal) and vessel scour. Standard vessel management plans are part of CEMP and OEMP would control vessel movements and restrict impacts to the seabed conditions from these activities to low.

The sediments are not expected to significantly change as a result of the port development at the site. Dust emissions are proposed to be controlled via engineered solutions on the landward side and the use of chemical veneering of ore concentrate stockpiles. Air quality modelling has shown dust emission into the marine environment will be negligible (Jacobs, 2014b) and it is expected that dust will be dispersed by consistent breezes and strong currents in the marine environment or incorporated into the seagrass meadows via natural sediment trapping processes. The effect of the release of iron into the marine environment is discussed in more detail in Section 5.4, and is considered highly unlikely to cause a measurable change in iron concentration in marine sediments or have toxic effects as iron ore is biologically inert.

5.3.2.2 Risks

If there was a large scale loss of seagrass at the site, sediment transport processes would be at risk of substantial change, which could result in the redistribution of sediment, or changes to seabed conditions. However, the likelihood of widespread seagrass loss at the site is considered to be unlikely with only minor changes to seagrass coverage confined to the infrastructure footprint anticipated. The consequences of widespread seagrass loss to the seabed conditions would be minor (a local long term decrease in abundance of some species) meaning the overall risk to seabed condition is considered low. Impacts and risks to seagrasses are discussed further in Section 6.3.

The creation of hardstand areas for the land based facilities creates the risk of increased flows of run-off channelling into the marine environment that could change the sediment loading and distribution at the site. Run-off from the land based operations also has the risk of introducing contaminants into marine sediments. Run-off risks will be mitigated via the landside stormwater capture design which is proposed for the development. As such, the likelihood of stormwater run-off into the marine environment is considered to be unlikely with minor consequences. The risk to seabed condition is therefore considered to be low.

There is a risk that seabed conditions could be impacted as a result of spills and leakages during fuel loading or refuelling, oil change or lubricant escape from the shiploader, anti-fouling paints or escape of iron ore into



the marine environment. An effectively implemented OEMP would reduce the likelihood and consequences of a spill. A minor spill event is considered to be possible (i.e. a 5% chance per annum), the consequences to seabed conditions are considered to be minor (local effects only), the risk of spills and leaks to seabed conditions would be considered to be low. A major spill event would be considered unlikely rather than possible, with elevated consequences of a moderate rating due to the more widespread implications. In this instance, a major spill event would be considered a medium risk to seabed conditions.

5.3.3 Conclusions

The substrate characteristics and geochemistry at Cape Hardy are typical of similar Spencer Gulf green-field sites.

- The design and location of the port will not require dredging which will avoid large scale impacts to the seabed.
- The substrate in shallow water areas is generally sandy, becoming coarser closer to shore in the northern areas. The sandy surface substrate is not expected to be a source of long-term resuspension due to the type of construction activities in the shallow water. In the areas of deeper water the sediment becomes increasingly fine, and is expected to be more easily suspended in the water column. However the impact piling method planned for construction (hollow steel tubes to be later filled with concrete) will result in only localised, negligible impacts.
- While impacts from vessel scour in deep water are considered low, propwash and vessel scour from tug movements during operation are expected to have medium impacts to the nearshore areas of the seabed.
- There will be a permanent impact on the seabed from the placement of the MOF and tug harbour. Although the area of the impact will be small scale due to the footprint of the structures, the permanent nature of the change escalates this impact to medium.
- Effective CEMP and OEMPs would minimise the risk of accidental releases into the marine environment but there remains a low risk from minor spills and a medium risk from major spills to the seabed condition as a result of the introduction of shipping activities into the environment.
- As the areas of elevated metals are restricted to the north of the study area outside the development footprint and all Cape Hardy samples were below guideline levels, sediments are considered to be uncontaminated, and therefore the redistribution of contaminants due to sediment movement during construction and operation is unlikely to be of concern. There is only a low possibility that sediments with elevated metals or contaminants will be exposed during construction and transported across the site. A suitable CEMP would reduce risks from sediment resuspension should they arise.

5.4 Water quality

Water quality is driven by the natural biogeochemistry of the environment as well as anthropogenic activities. Ultimately, water quality conditions drive the ecology of the broader marine environment by determining the availability of light, nutrients, oxygen and other parameters, as well as the exposure of organisms to pollutants. This makes water quality a valuable indicator of the condition of the marine system. Small disruptions to the dynamics of variables such as temperature, salinity, pH or light penetration can trigger a biological response. The following section describes the results of a survey which aimed to characterise the water quality at the site, and to assess potential impacts upon water quality as a result of the development.

5.4.1 Existing Environment and Key Findings

South Australian marine waters are characterised by low levels of dissolved nutrients, which is supported by the results of the water quality surveys at Cape Hardy. This is due to its physical isolation from strong oceanographic currents and minimal terrestrial runoff from rivers or creeks into the gulf that could supply



nutrients. The existing water quality conditions for Cape Hardy described in this section are based on publicly available data as well as spot samples for metals and nutrients collected from Cape Hardy in November 2011. Concentrations of total metals at Cape Hardy were compared against EPA data from other locations in Spencer Gulf i.e. Port Augusta, Port Germein, Port Pirie, Whyalla, and Port Hughes (EPA, 2004; EPA, 2009). A plot of this data comparison can be seen in Figure 5-6. From this comparison, it is clear that the concentrations of the measured metals at Cape Hardy are below or at similar levels to existing port locations within Spencer Gulf.




Figure 5-6 Comparison of metal concentrations in Spencer Gulf waters

EPA trigger guidelines for metal contaminants were used for comparison as a guide to ascertain if levels were above those considered to be of concern however guidelines are not available for all analytes. For some of the analytes, the sampled values were below the LOD of the laboratory, and in those cases to be conservative, values were presented as the LOD rather than zero. Results from metal analysis are presented in Table 5-3 and Table 5-4. A summary of the results is bulleted below and the full ALS Environmental Laboratory results are included in Appendix D.

• With the exception of Cu, the levels of metals in surface and bottom water samples taken within the study area were all below EPA trigger values (Table 5-3, Table 5-4)

JACOBS

- Cu values were consistently at or just above the EPA trigger value of 0.01 mg/L
- Concentrations of Fe in surface waters were all at the LOD, hence there is no variation in the data and no median was presented (Table 5-3)
- Concentrations of Fe in bottom waters were higher than surface water concentrations

Table 5-3 Surface water composition: summary in comparison with EPA Water Quality Guidelines

| Analyte | Unit | LOD | SA EPA | Surface Water - Sampled Values | | | | | | | |
|----------------|------|--------|---------|--------------------------------|---------|---------|----------|--------|--|--|--|
| Total Metals | | | Trigger | Lowest | Highest | Average | ± SE | Median | | | |
| Arsenic (As) | mg/L | 0.001 | 0.05 | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | | |
| Barium (Ba) | mg/L | 0.001 | n/a | 0.006 | 0.007 | 0.006 | ± 0.000 | 0.006 | | | |
| Beryllium (Be) | mg/L | 0.001 | n/a | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | | |
| Cadmium(Cd) | mg/L | 0.0001 | 0.002 | 0.0001 | 0.0006 | 0.0001 | ± 0.0000 | 0.0001 | | | |
| Cobalt (Co) | mg/L | 0.001 | n/a | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | | |
| Chromium (Cr) | mg/L | 0.001 | n/a | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | | |
| Copper (Cu) | mg/L | 0.001 | 0.01 | 0.01 | 0.02 | 0.010 | ± 0.002 | 0.01 | | | |
| Iron(Fe) | mg/L | 0.05 | n/a | 0.05 | 0.05 | 0.05 | ± 0.000 | n/a | | | |
| Manganese (Mn) | mg/L | 0.001 | n/a | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | | |
| Mercury (Hg) | mg/L | 0.0001 | 0.0001 | 0.0001 | 0.0001 | 0.0001 | ± 0.0000 | 0.0001 | | | |
| Nickel (Ni) | mg/L | 0.001 | 0.015 | 0.002 | 0.004 | 0.002 | ± 0.001 | 0.002 | | | |
| Lead (Pb) | mg/L | 0.001 | 0.005 | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | | |
| Vanadium (V) | mg/L | 0.01 | n/a | 0.01 | 0.01 | 0.01 | ± 0.000 | 0.01 | | | |
| Zinc (Zn) | mg/L | 0.005 | 0.05 | 0.005 | 0.028 | 0.007 | ± 0.001 | 0.005 | | | |

NOTE: SE = standard error of the mean. EPA guidelines for total metals trigger where used. n/a = no trigger. Values in **bold** are at or below the LOD i.e. As, Be, Cr, Fe, Hg, Mn and Pb were all below the LOD for all samples.

Table 5-4 Bottom water composition: summary in comparison with EPA Water Quality Guidelines

| Analyte | Unit | LOD | SA EPA | Bottom Water - Sampled Values | | | | | | |
|---------------------|-------|--------|---------------|-------------------------------|---------|---------|----------|--------|--|--|
| Total Metals | Offic | LOD | Trigger | Lowest | Highest | Average | ± SE | Median | | |
| Arsenic (As) | mg/L | 0.001 | 0.05 | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | |
| Barium (Ba) | mg/L | 0.001 | n/a | 0.005 | 0.007 | 0.006 | ± 0.000 | 0.006 | | |
| Beryllium (Be) | mg/L | 0.001 | n/a | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | |
| Cadmium (Cd) | mg/L | 0.0001 | 0.002 | 0.0001 | 0.0001 | 0.0001 | ± 0.0000 | 0.0001 | | |
| Cobalt (Co) | mg/L | 0.001 | n/a | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | |
| Chromium (Cr) | mg/L | 0.001 | n/a | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | |
| Copper (Cu) | mg/L | 0.001 | 0.01 | 0.01 | 0.012 | 0.01 | ± 0.00 | 0.01 | | |
| Iron (Fe) | mg/L | 0.05 | n/a | 0.05 | 0.17 | 0.081 | ± 0.043 | 0.055 | | |
| Manganese (Mn) | mg/L | 0.001 | n/a | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | |
| Mercury (Hg) | mg/L | 0.0001 | 0.0001 | 0.0001 | 0.0001 | 0.0001 | ± 0.0000 | 0.0001 | | |
| Nickel (Ni) | mg/L | 0.001 | 0.015 | 0.003 | 0.006 | 0.004 | ± 0.001 | 0.005 | | |
| Lead (Pb) | mg/L | 0.001 | 0.005 | 0.001 | 0.001 | 0.001 | ± 0.000 | 0.001 | | |
| Vanadium (V) | mg/L | 0.01 | n/a | 0.01 | 0.01 | 0.01 | ± 0.0000 | 0.01 | | |
| Zinc (Zn) | mg/L | 0.005 | 0.05 | 0.005 | 0.014 | 0.008 | ± 0.004 | 0.006 | | |

NOTE: SE = standard error of the mean * there are no EPA guidelines for these dissolved metals in marine waters, total metals trigger where used instead. n/a = no trigger Values in **bold** are at or below the LOD i.e. As, Be, Cr, Fe, Hg, Mn and Pb were all below the LOD for all samples.

JACOBS

The nutrient compositions of surface and bottom water samples at Cape Hardy are presented in Table 5-5. The full laboratory results are presented in Appendix D with a summary of key findings below.

Concentrations of nutrients in surface and bottom waters at Cape Hardy were quite low, and often at or below the LOD (Table 5-5). The low nutrient levels measured around Cape Hardy is typical of Spencer Gulf which has been shown to be nutrient poor, including phosphorous (Middleton et al., 2013)

There was little difference in concentrations between surface and bottom waters, suggesting that the water column was well mixed and homogeneous in nutrient composition (Table 5-5)

All sample averages were below EPA trigger values for ammonia, total oxidised nitrogen (sum of nitrite and nitrate), total nitrogen, and total phosphorus. The concentrations of chlorophyll a (and pheophytin) were low at Cape Hardy, with the average concentration in surface waters 1.4 mg/m³, and ranged from 1 to 3 mg/m³. The low levels of chlorophyll a and nutrients suggest that the Cape Hardy site is oligotrophic, which is characteristic of waters in Southern Australia (Russell et al., 2005) and particularly Spencer Gulf (Middleton *et al.*, 2013)

Table 5-5 Surface and bottom water nutrient and chlorophyll a summary in comparison with EPA Water Quality Guidelines

| Analyto | Unit | LOD SA EPA | | | Sampled Values | | | | | |
|--|---------|------------|---------------|------|----------------|--------|---------|--------|--|--|
| Analyte | Onit | LOD | Trigger* | Min | Max | Averag | ge ± SE | Median | | |
| | | Surface | water sample: | | | | | | | |
| Ammonia as N | mg/L | 0.01 | 0.2 | 0.08 | 0.12 | 0.1 | ± 0.0 | 0.1 | | |
| Nitrite as N | mg/L | 0.01 | n/a | 0.01 | 0.01 | 0.01 | ± 0.00 | 0.01 | | |
| Nitrate as N | mg/L | 0.01 | n/a | 0.01 | 0.02 | 0.01 | ± 0.00 | 0.01 | | |
| Nitrite + nitrate (oxidised nitrogen) as N | mg/L | 0.01 | 0.2 | 0.01 | 0.02 | 0.01 | ± 0.00 | 0.01 | | |
| Total Kjeldahl nitrogen (TKN) as N | mg/L | 0.1 | n/a | 0.1 | 0.6 | 0.2 | ± 0.0 | 0.2 | | |
| Total nitrogen as N | mg/L | 0.1 | 5 | 0.1 | 0.6 | 0.2 | ± 0.0 | 0.2 | | |
| Total phosphorus as P | mg/L | 0.01 | 0.5 | 0.01 | 0.09 | 0.02 | ± 0.00 | 0.02 | | |
| Chlorophyll-a + pheophytin | mg/m³ | 1 | n/a | 1 | 3 | 1.4 | ± 0.1 | 1 | | |
| | | Bottom | water sample: | s | | | | | | |
| Ammonia as N | mg/L | 0.01 | 0.2 | 0.08 | 0.16 | 0.11 | ± 0.00 | 0.105 | | |
| Nitrite as N | mg/L | 0.01 | n/a | 0.01 | 0.01 | 0.01 | ± 0.00 | 0.01 | | |
| Nitrate as N | mg/L | 0.01 | n/a | 0.01 | 0.04 | 0.01 | ± 0.00 | 0.01 | | |
| Nitrite + nitrate (oxidised nitrogen) as N | mg/L | 0.01 | 0.2 | 0.01 | 0.04 | 0.01 | ± 0.00 | 0.01 | | |
| Total Kjeldahl nitrogen (TKN) a N | as mg/L | 0.1 | n/a | 0.1 | 1 | 0.2 | ± 0.0 | 0.1 | | |
| Total nitrogen as N | mg/L | 0.1 | 5 | 0.1 | 1 | 0.2 | ± 0.0 | 0.1 | | |
| Total phosphorus as P | mg/L | 0.01 | 0.5 | 0.01 | 0.14 | 0.03 | ± 0.00 | 0.02 | | |

NOTE: SE = standard error of the mean * values of n/a indicates no EPA guidelines for marine waters



5.4.2 Impacts and Risks: Water quality

Water quality could potentially be affected by alterations to the chemical dynamics of nutrients, contaminants or suspended particles. Even slight alterations to water quality can cause disproportionate biological responses by impacting on phytoplankton or epiphytic algae and, over long periods the loss of habitat-forming macrophytes such as seagrass or macroalgae. Changes in water quality, particularly from pollution can also directly impact on fauna, including habitat forming species, which can lead to habitat loss or changes of species composition within the ecological community of a site.

Impacts could potentially occur to water quality as a result of the following, which are individually assessed in further detail below:

- Dredging / blasting during construction
- Leakage or spills of stored fuels, waste and chemicals used on site
- Chemical or waste spills from the jack-up barge (during construction) or vessels (construction and operation)
- Erosion caused by earthworks and road building during construction
- Alterations to drainage due to construction of hardstand areas increasing run-off of freshwater and input of nutrients, suspended solids or other contaminants
- Wastewater and stormwater from the landside construction site which potentially carry contaminants into the marine environment
- Oil spill during fuel loading of vessels or fuel transfer
- Nutrient or iron stimulation of phytoplankton communities
- Introduction of IMS algae
- Elevated turbidity as a result of piling during construction or vessel scour

5.4.2.1 Impacts

The iron concentrate intended for shipment by Iron Road is magnetite-based and known to be insoluble in seawater, and is therefore highly unlikely to cause a measurable elevation in dissolved iron concentrations in either the water column or surrounding sediment in the event of dust or accidental release to the marine environment. For there to be an iron 'fertilisation' effect on phytoplankton or algae, the receiving waters need to be very low in iron but high in other nutrients. Such environments are referred to as High-Nutrient Low-Chlorophyll (HNLC) environments (Bowie et al., 2001). From previous surveys of the gulf it is clear that the Spencer Gulf environment has relatively high levels of metals including iron (Table 5-3 and Table 5-4) when compared with oceanic waters (Martin et al., 1994; Thompson et al., 2008). The marine environment in the Lipson Island to Port Neill area is considered low in dissolved and total nutrients based on previous sampling (Table 5-5; Golder Associates, 2012; Middleton et al., 2013). Middleton et al. (2013) noted that concentrations of all macronutrients within the Spencer Gulf are limited and primary productivity within the gulf is restricted by these nutrient limitations. As the marine waters surrounding Cape Hardy are neither high in nutrients nor low in iron, they do not fit the definition of a HNLC system (Hutchins et al., 1999). Therefore it is considered highly unlikely that iron fertilisation could occur as a result of air-borne emissions or an iron ore spill. There are expected to be no impacts to water quality from iron induced algal blooms.

Increased current and wave energy during storm conditions is an important part of nutrient dynamics, resuspending nutrient-rich sediments, but also resulting in elevated turbidity close to shore for a series of days after storm events. The solid tug harbour and MOF may influence how sediments are redistributed nearshore by causing settlement adjacent to the leeward side of the harbour and MOF. The impacts to water



quality following storm events as a result of the MOF and tug harbour are considered low (short durations following storm events in a localised area).

Minor (low) impacts to water quality are expected from temporary increases in turbidity as a result of vessel movements and vessel scour (localised and temporary in nature).

As a result of the mitigation measures designed for containment of run-off and wastewater at the site impacts on water quality as a result of surface water run-off during construction and operation are expected to be negligible. There is a risk of surface water run-off during extreme storm events, which is discussed further below.

5.4.2.2 Risks

The containment of stormwater and wastewater within the port site in all but the most extreme storm events means that the risk to water quality as a result of sediments or contaminants within run-off is considered low; an unlikely event with minor consequences.

As the sediments in the study area were found to be uncontaminated, it is considered unlikely that the proposed development will disturb or cause the redistribution of any contaminated sediments. Appendix C discusses the ability of sediments and infauna / epifauna to accumulate metals and the effects of sediment re-suspension on water chemistry. Consequences of any resuspended contaminants on water quality are considered minor – local and short term. It is therefore considered a low risk that existing metals or other contaminants would be released from marine sediments and impact water quality.

An effective CEMP and OEMP would reduce the likelihood of oil spill and leaks during construction and operation which could degrade water quality at the site. A minor spill would be considered possible, but with only minor consequences. As such, the risk to water quality from a minor spill or leak would be considered low. A major spill or leak would have moderate consequences for water quality, but are considered to be an unlikely event. As such, the risk from a major oil spill or leak is considered to be medium.

As with any operating port the introduction of IMS is a possibility, however the likelihood of introducing an algal species that could have an affect on water quality is considered unlikely due to the Commonwealth requirements for vessels to discharge ballast water in open waters and the requirement for operators to adhere to a CEMP and OEMP for the port. Given the consequences of such an introduction on water quality would be considered minor (localised and short term), the risk to water quality from IMS is considered to be low.



5.4.3 Conclusions

Water quality and chemistry at Cape Hardy is typical for the Spencer Gulf as a region. Nutrient concentrations are low, as expected in this system (Russell *et al.* 2005). Comparison with data from other ports in the region shows that Cape Hardy has similar or lower levels of metals to Spencer Gulf waters (Figure 5-6), which have high levels of metals when compared with oceanic waters (Martin *et al.*, 1994; Thompson *et al.*, 2008). The Spencer Gulf system does not fit the requirements for iron fertilisation to take place.

Combining the knowledge gained from investigations into the substrate geomorphology, chemistry and water quality, it is considered that the site at Cape Hardy is uncontaminated, and that contamination is unlikely to occur as a direct result of incidental emissions, sediment movement or redistribution during the construction and operational phases of this development.

The key risks to water quality at the Cape Hardy site include:

- Elevated turbidity and/or redistribution of contaminants due to construction
- Elevated turbidity and/or redistribution of contaminants due to ship movements and/or altered hydrodynamic conditions
- Vehicle and equipment maintenance and operations, vehicle washing, possible dust control
 chemicals, storage of fuel, lubricants, other chemicals, various aerial emissions including dust are all
 risks to the water quality at the site
- Stormwater and wastewater entering the marine water quality during extreme storm events
- Alterations to drainage due to construction of hardstand areas resulting in increased run-off of freshwater and input of nutrients, suspended solids or other contaminants

It is expected that the majority of these risks are mitigated to medium or less, due predominantly to the site selection and the design standards used for capture and minimisation of run-off emissions from the port site. Due to these measures impacts to the water quality from waste water or run-off are expected to be low. The planned construction methods for the marine infrastructure and adherence to standard mitigation strategies outlined within a comprehensive CEMP and OEMP will further mitigate risks.



6 Biological Environment

The assessment of the biological environment commenced with a desktop study of marine flora and fauna, to determine the potential presence of species and habitats of conservation value within the study area. The EPBC Act and NPW Act were utilised as the major reference points in this component of work. Key species of concern identified during the investigation included protected marine mammals, birds, seahorses, pipefish, and habitats such as seagrass meadows. Seagrass areas are of high conservation value and are protected under the Native Vegetation Act.

A number of EPBC listed species were sighted during field investigations including the hooded plover, common dolphin and leafy seadragon. Other sightings of interest include juvenile Port Jackson sharks (*Heterodontus portusjacksoni*) and a long-snouted boarfish (*Pentaceropsis recurvirostris*).

The key benthic marine habitats and species of interest or conservation value were mapped and identified during December 2011 field expeditions within the study area.

The Cape Hardy biological environment is typical of the Spencer Gulf region. The shallow sandy areas are inhabited by mature seagrass meadows, predominantly *Posidonia* spp. In deeper waters, "clumps" of invertebrates including large ascidians, sponges and bivalves grow on the silty bottom. Small areas of rocky reef exist in rocky areas adjacent to the headlands and the intertidal zones of the study area. These habitats generally follow depth and sediment contours, and as such appear to be stable in their distribution (based on the lack of obvious habitat loss and relative age of the habitats). In the deepest areas with the finest sediments, there is evidence of the formation of microbial matting.

The marine habitats throughout the study area were observed to be in generally healthy condition. Shallow sandy areas are generally dominated by mature seagrass meadows. Temperate rocky reef habitat was limited to shallow areas adjacent the rocky headlands and a few sub-tidal rocky ridges.

6.1 Coastal Habitats

The coastal habitat of the Spencer Gulf is varied, supporting a wide range of habitats and species groups. In the upper reach of the gulf, tidal flats and mangroves are key habitats. In the lower reaches of the gulf sandy beaches backed by dunes or rocky cliff coasts tend to dominant. There are a number of species known to be endemic to the Spencer Gulf. Each of these species has been identified in the upper gulf, as many ecological investigations have focused on this area, whereas information on the southern reaches of the gulf (including the study area) is limited. To assess the biological importance of the Cape Hardy site it was necessary to determine which habitats and species exist at, or utilise the study area, as well as identify any species that are protected by legislation and may trigger further regulatory requirements. Rocky shores or sandy beaches with seagrass provide essential habitat to a wide range of bird species as well as some reptiles and mammals.

6.1.1 Existing Environment and Key Findings

The EPBC Act Protected Matters Search Tool (PMST) found no Threatened Ecological Communities that occurred within 10 km of the study area. However, 56 birds of national conservation significance and two skink species of state significance were identified as having the potential to occur, based upon records from the Biological Database of South Australia (BDBSA), as well as results from the EPBC PMST. In addition to the desktop study, species of conservation significance were also identified at the site during the marine surveys and the terrestrial flora and fauna survey (Jacobs, 2014a).



During the marine field surveys, a total of 14 bird species listed under the EPBC Act were observed within the study area, four of which are also listed under the NPW Act schedules. Unlike many areas in the northern regions of Spencer Gulf, the Cape Hardy coastal area contained no mangrove trees and no flora species listed under the EPBC Act. The dune habitat at Cape Hardy was not considered suitable for conservation significant samphire species such as *Tecticornia flabelliformis*.

For the purposes of this report only the 56 conservation listed marine bird species and two lizard species identified as potentially occurring within the coastal habitat are further considered, as the terrestrial flora and fauna report (Jacobs, 2014a) assesses the on-shore native vegetation and terrestrial mammals, reptiles and birds. Of the 56 protected bird species, 30 are considered unlikely to occur within the study area, 12 are considered to potentially occur in the study area and 14 were actually observed in the study area by survey teams (Table 6-1). While the site contains suitable habitat for the two lizard species they are only considered as potentially occurring at the study area as all known populations are found on the west coast of the Eyre Peninsula.

Based on the desktop reviews of the available literature, BDBSA, EPBC, ALA and field surveys at the port site there are no known large breeding or roosting sites for any shorebird species within the study area with only one species of shorebird (hooded plover) known to nest at the site. Therefore the study area is not considered an internationally or nationally significant shorebird area. Furthermore, of the nationally and internationally significant shorebird areas on the Eyre Peninsula only Franklin Harbour and Sleaford Bay are on the east coast of the Eyre Peninsula (Caton *et al.*, 2011a). As both these locations are greater than 50 km from the study area, it is unlikely those colonies of shorebirds would rely on habitat within the study area. All of the other nationally and internationally significant shorebird areas on the Eyre Peninsula: Baird Bay, Sceale Bay (including Seagull Lake), Streaky Bay (including Acraman Creek), Lake Newland Conservation Park including the ocean beach), Tourville Bay, Murat Bay, Eyre Island, St Peter Island, Coffin Bay (Point Longnose and Gunyah Beach) and Venus Bay (including the islands within the bay) are on the western coast of the Eyre Peninsula (Caton *et al.*, 2011a).

6.1.2 Impacts and Risks: Coastal Habitats

Impacts and risks to the landside coastal environment associated with the proposed port facility have been discussed in detail by Jacobs (2014a). The following section discusses the impact on coastal habitats of fauna, marine bird species and reptiles that have the potential to be impacted by the construction and operation of the proposed development.

The impacts and risks involved with the proposed development are short-term impacts during the construction phase, such as noise impacts from piling, as well as long-term impacts throughout the ongoing presence of structures and operations at the site. Impacts to coastal habitats could occur as a result of:

- Habitat loss or exclusion
- Trampling
- Entanglement in debris or infrastructure or ingestion of non-biological waste such as plastics
- Artificial lighting at night has the potential to disturb nesting birds or interfere with their navigation and foraging
- Stormwater runoff (including erosion)
- Coastal erosion from vessel scour or propwash
- Run-off including pesticides, herbicides, spills (e.g. oil) or other pollutants
- Introduction or increased attraction of invasive, feral or pest species.



6.1.2.1 Impacts

As previously outlined, a number of design and mitigation strategies have been implemented to minimise disturbance of coastal habitat including remnant vegetation and sandy beach areas. The coastal habitat at Cape Hardy is not considered to be core habitat for any EPBC or NPW listed species. CEMP and OEMP procedures will restrict access to sandy beach areas and the engineering design has avoided this habitat. As such, no significant change to the protected environmental values associated with coastal fauna and habitats will occur, with the overall impact considered to be negligible.

Noise impacts from construction vessel movements and construction activities (such as piling) have the potential to disturb fauna utilising coastal habitats, as well as marine fauna (detailed in Section 6.5) that may be prey for coastal species. Coastal fauna known or likely to frequent the port site such as the Hooded Plover, are susceptible to disturbance by noise. There are no large breeding colonies of birds that utilise the port site for habitat or foraging purposes, however, Hooded Plovers are known to frequent the port site and were identified during field surveys. Hooded Plovers are known to have territorial ranges of over 30 ha with several nesting beaches several kilometres apart within their territory (SPRAT, 2014). As such, it is expected breeding pairs disturbed by construction could readily relocate to nearby beaches and be able to return once construction has ceased. Shorebird prey species such as fish or cephalopods will be temporarily displaced from the port site as a result of underwater noise emissions (refer Section 6.7.2). Displacement of species will be localised (around noise sources) and on a temporary short term basis. As such, local populations of protected coastal fauna will experience short term disturbance during construction, considered to be a low impact. Feeding or foraging behaviour of seabirds will not be significantly affected, nor will key habitat be unusable as a result of construction noise emissions.

The introduction of artificial light sources during construction and operation of the proposed port will be required for safe night-time operations and way finding. Artificial light sources attract a number of marine species which may be suitable prey for coastal fauna, thus attracting additional marine birds and affecting the abundance of resources for resident species. As light spill will be minimised through the use of directional lighting (i.e. oriented to a specific area) and the port site is not utilised by any large breeding colonies no change to protected environmental values is anticipated and the overall impact is considered to be negligible.

An assessment of the likelihood of occurrence and impacts to each of the 56 protected bird species and 2 protected lizards species identified as potentially present at the port site is provided in Table 6-1 below.



Table 6-1 Bird and reptile species: likelihood of occurrence and potential impacts

| Protected Species p | otentially present at th | ie port | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|--------------------------------|--|---|--|---|--|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Birds | | | <u>'</u> | | | | |
| Acanthiza iredalei iredalei | Slender Billed Thornbill | Vulnerable (EPBC); Vulnerable (NPW) | Unlikely to occur at the port. This species prefers dense samphire, dense chenopod shrubland, or other low shrubland habitat not present within the project area. This species was not recorded during Jacobs' site surveys. No records occur in a BDBSA 10 km buffer search from site (BDBSA, 2013). This species has potential to overfly the development. | Loss of habitat through clearance | As the preferred habitat for this species is not found at this site, it is highly unlikely that the development area would disturb a significant area of the foraging grounds or range of this species. The species range is extensive, including inland and coastal areas of Australia (SPRAT, 2014). Therefore, any potential impacts to this species from the construction and operation of the port at Cape Hardy are not considered significant. | N | |
| Anthus novaeseelandiae | Australasian Pipit, Richard's Pipit | Marine (EPBC) | Present. This species was sighted within the study area during the 2011 Jacobs port site flora and fauna surveys and was sighted at road sidings beside farmland during 2012 Jacobs site visits. BDBSA records exist at Port Neill and near Lipson Cove from 2000-2008 (BDBSA, 2013. The species is found across Australia (Simpson and Day, 1999) as well as New Guinea, New Zealand, as well as being widespread across Africa and Asia. | Loss of habitat through clearance; oil spill impacts to intertidal; entanglement in marine debris; vehicle traffic | As the geographic range of this species covers the entire Australian continent this development would not pose any significant loss to habitat or foraging area for this species. Therefore, any potential impacts to this species from the construction and operation of the port at Cape Hardy are not considered significant. | N | |
| Apus pacificus | Fork-tailed Swift | Migratory, Marine (EPBC) | Unlikely to occur at the port. There are no known nesting sites are within 5 km of the proposed development with the nearest recorded sightings near Port Neill in 1988 and 1998 (BDBSA, 2013). While this species was not observed during Jacobs' site surveys it may overfly the study area but given the species has not be recorded in the area for over decade it is unlikely to be a frequent visitor to the port. | Loss of habitat through clearance | The development represents only a small part of the migratory range of this species, which extends over most of the Australian continent (SPRAT, 2014). As such there would be no significant loss of habitat for this species and therefore impacts to this species from habitat loss are not considered significant. | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|-----------------------------|---|--|---|---|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Ardea alba | Great Egret, White Egret | Migratory, Wetland, Marine (EPBC) | Unlikely to occur at the port. Although this species may have the potential to overfly the area, this species was not sighted during Jacobs' surveys, and the preferred habitat of this species is floodplains, rivers, shallow wetlands and intertidal mudflats (Simpson and Day, 1999). Considering that no preferred habitat exists within the development area (i.e. no tidal creeks or mudflats) there is low potential for the species to occur within the study area. | Loss of habitat through clearance; oil spill impacts to intertidal; entanglement in marine debris | The development represents only a small part of the migratory range of this species, which extends over the entire Australian landmass including offshore islands (SPRAT, 2014). As such there would be no significant loss of habitat for this species and therefore impacts to this species from habitat loss are not considered significant . | N | |
| Ardea ibis | Cattle Egret | Migratory, Wetland, Marine (EPBC); Rare (NPW) | Unlikely to occur at the port as the species preferred habitat of mudflats or tidal creeks does not exist within 10 km of the port (BDBSA, 2012). There are no records for this species in the area and it was not sighted during Jacobs' site surveys. | Loss of habitat through clearance; oil spill impacts to intertidal; entanglement in marine debris | This species is migratory and has a large range covering most of the Australian continent (SPRAT, 2014). As such there would be no significant loss of habitat for this species and therefore impacts to this species from habitat loss are not considered significant . | N | |
| Arenaria interpres | Ruddy Turnstone | Migratory, Marine (EPBC); Rare (NPW) | Possible species will forage within the study area. The PMST did not identify this species for the area however it has been recorded at Port Neill from 1998-2009 (BDBSA, 2013), 7 km northeast of the development. The species is considered moderately common with a preference for rocky shores with seagrass wrack which it forages amongst (Simpson and Day, 1999). The species was not sighted during the site surveys but as there are small areas of rocky shore within the study area it is possible that this species will occur at the port site. | Loss of habitat through clearance; oil spill impacts to intertidal; entanglement in marine debris | This species is migratory and has a large range covering most of the Australian coastline (SPRAT, 2014). Therefore, despite the likelihood of the species being found at the site its critical habitat would not be affected, nor would the development represent a significant area of the species foraging grounds or range. Therefore impacts to this species from habitat loss are not considered significant. | N | |
| Calidris acuminata | Sharp-tailed sandpiper | Migratory, Wetland, Marine (EPBC) | Possible species will forage within the study area, as the port area offers coastal shore areas and the species has been recorded at Port Neill in 2000, 7 km north-east of the development (BDBSA, 2013). | Loss of habitat through clearance; oil spill impacts to intertidal; entanglement | This species is migratory and has a large range covering large areas inland Australian and coastline (SPRAT, 2014) therefore despite the likelihood of the species occurring at the port, its critical habitat would not be affected nor would the development represent a significant area of the species foraging grounds or range. Impacts | N | |



| Protected Species p | ootentially present at t | he port | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|-----------------------------|------------------------------|---|---|---|---|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | in marine debris | to this species from the port are not considered significant . | | |
| Calidris ruficollis | Red-necked stint | Migratory, Marine (EPBC) | Possible will forage within the study area. This species is considered common or abundant across most of Australia and found in very large flocks in coastal and inland shores (Simpson and Day, 1999). There are no known breeding colonies on Eyre Peninsula, however as the port area offers coastal shore areas and the species has been recorded between the years 1998-2009 at Port Neill, 7 km north-east of the development (BDBSA, 2013). | Loss of habitat through clearance; oil spill impacts to intertidal; entanglement in marine debris | This species is migratory and has a large range covering numerous sections of the Australian coastline and some inland sites (SPRAT, 2014). Therefore, despite the potential for this species to occur at the port, its critical habitat would not be affected nor would the development represent a significant area of the species foraging grounds or range. Impacts to this species from the port are not considered significant. | N | |
| Catharacta skua | Great skua | Marine (EPBC); Vulnerable (NPW) | Unlikely to occur at the port site. This species was not observed during port site surveys, and no records of this species occur within 10 km of the study area (BDBSA, 2013). The species uses open-ocean habitats as well as coastal habitats (Simpson and Day 2004). This species has a large feeding range covering the entire southern Australian coastline and offshore in the Southern Ocean (SPRAT, 2014). Therefore although it may have the potential to overfly the development, the area would not represent a significant area of the species foraging grounds or range. | Loss of habitat through clearance or disturbance, oil spill impacts to foraging areas; entanglement in marine debris | It is known to visit southern Australian waters but is highly mobile and ranges widely. It is not expected that the Great skua is reliant upon any habitat resources located within the study area or surrounding habitats. Consequently, it is not expected that the construction and operation of a port facility at Cape Hardy will significantly impact this species. Impacts to this species from the port are not considered significant. | N | |
| Coracina novaehollandiae | Black-faced cuckoo-shrike | Marine (EPBC) | Present. This species was sighted within the development area during the 2011 site flora and fauna survey (Jacobs, 2014a). The species prefers most wooded areas including suburban areas (SPRAT, 2014). This species is common throughout Australia with a range including all of Australia including Tasmania (Simpson and Day, 2013). | Loss of habitat through clearance or disturbance | Despite this species existing in the area of the port site, the site does not represent critical habitat for the species as its habitat and range is extensive and (SPRAT, 2014). The species is considered common throughout its range; the development would not pose any significant threat to the species due to its known ability to adapt to human settlements. Impacts to this species from the port are not considered | N | |



| Protected Species po | otentially present at t | he port | Likelihood of Occurrence | Likelihood of si | Likelihood of significant impacts | | | |
|---|------------------------------|---|--|--|--|-----------------------------|--|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | | |
| | | | | | significant. | | | |
| Corvus mellori | Little raven | Marine (EPBC) | Present. This species was sighted within the development area during the site flora and fauna survey and previously recorded in the surrounding areas at Port Neill, Lipson Island and Lipson Cove from 1998-2008 (BDBSA, 2013). This species is common across farmlands and suburban areas in South Australia, Victoria and NSW (SPRAT, 2014). This species readily exploits human developments for habitat and food sources. | Loss of habitat through clearance or disturbance; Attraction to development and subsequent entanglement in marine debris; pollution | Due to the distribution and adaptability of this species, any potential impacts to this species from the construction and operation of the port are not considered significant. | N | | |
| Coturnix pectoralis | Stubble quail | Marine (EPBC) | Present. This species was sighted within the development area during the 2011 site flora and fauna survey (Jacobs, 2014a). This species can be found in many grassy areas with a widespread distribution across Australia including many inland areas (Simpson and Day, 1999). This species is not considered threatened in any region across Australia (SPRAT, 2014). | Loss of habitat through clearance or disturbance; vehicle traffic; predation from feral pests or dogs; pollution; entanglement with debris | As the development does not represent a significant geographic area of habitat any potential impacts to this species from the construction and operation of the port are not considered significant. | N | | |
| Chalcites basalis = Chrysococcyx basalis | Horsfield's bronze-cuckoo | Marine (EPBC) | Present. This species was sighted within the development area during the 2011 site flora and fauna survey (Jacobs, 2014a). This species is common throughout Australia with its range covering all of Australia and offshore islands including Tasmania. This species is found in wooded areas including farmlands and suburban areas as long as there are some trees (SPRAT, 2014). | Loss of habitat through clearance or disturbance | Despite this species being known to exist in the area of the port site, its range is extensive and the species common (SPRAT, 2014). The development does not represent critical habitat for this species. Given its ability to adapt to human settlements any potential impacts to this species from the construction and operation of the port are not considered significant . | N | | |



| Protected Species po | tentially present at the | he port | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|--------------------------------------|---------------------------------------|---|---|---|--|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Charadrius ruficapillus | Red-capped plover | Marine (EPBC) | Likely that species occurs in the area. The species was recorded 1 km away from the study area in 2009 and on numerous occasions at both Lipson Island to the South and Port Neill to the north (Golder Associates, 2012; BDBSA, 2013). | Loss of habitat through clearance or disturbance; oil spill impacts to intertidal; vehicle traffic; predation from feral pests or dogs; pollution; entanglement with debris | This species is migratory and has a large range covering numerous sections of the Australian coastline and some inland sites (SPRAT, 2014). Therefore, despite the high potential for this species to occur at the port, its critical habitat would not be affected nor would the development represent a significant area of the species foraging grounds or range. Impacts to this species from the construction and operation of the port are not considered significant. | N | |
| Charadrius veredus | Oriental plover, Oriental dotterel | Migratory, Wetland (EPBC) | Unlikely to occur at port site. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). Males of this species are conspicuous with red breasted breeding plumage (Simpson and Day, 1999), and would be expected to be easily identified if they were present in the wider area. The lack of sightings of this bird in the greater area suggests it is unlikely it would occur at the port site. | Loss of habitat through clearance or disturbance; oil spill impacts to intertidal; entanglement with debris | This species is migratory and has a large range covering numerous sections of the Australian coastline and inland sites (SPRAT, 2014). Therefore, given the specie's extensive range and the development would not represent critical habitat or a significant area of the species foraging grounds or range. Impacts to this species from the construction and operation of the port are not considered significant . | N | |
| Diomedea epomophora epomophora | Southern royal albatross | Vulnerable, Migratory, Marine (EPBC); Vulnerable (NPW) | Unlikely to occur at port site. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). This species does not breed in Australia, with all recorded breeding pairs nesting on islands offshore of New Zealand (SPRAT, 2014). This species is capable of global circumnavigation and therefore could in theory overfly the area but as they prefer open oceanic areas any visiting birds would be transients (SPRAT, 2014). Low potential to be present with no nesting sites in the area. | Entanglement in marine debris; disturbance; oil spills; pollution | As no critical habitat is present within the development area there is a low potential for this species to occur near the development area. The port site does not represent a significant area of the species foraging grounds or range. Impacts to this species from the construction and operation of the port are not considered significant. | N | |



| Protected Species p | otentially present at tl | he port | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|------------------------------------|--------------------------|---|--|--|---|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Diomedea epomophora sanfordi | Northern royal albatross | Endangered, Migratory, Marine (EPBC); Endangered (NPW) | Unlikely to occur at port site. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). The species does not breed in Australia, all breeding pairs nest on offshore New Zealand islands (SPRAT, 2014). The species range is wide covering the Southern Ocean from 36° S to at least 49°, including Australia and South America (SPRAT, 2014). This species is known to forage offshore in South Australian and Tasmanian using updrafts from open ocean wave fronts for lift (SPRAT, 2014). With no nesting sites in the area and its preferred foraging in open waters this species is therefore unlikely to frequent Spencer Gulf | Entanglement in marine debris; disturbance; oil Spills; pollution | As no critical habitat is present within the development area there is a low potential for this species to occur near the development area. The port site does not represent a significant area of the species foraging grounds or range. Impacts to this species from the construction and operation of the port are not considered significant. | N | |
| Diomedea exulans amsterdamensis | Amsterdam albatross | Endangered, Migratory, Marine (EPBC) | Unlikely to occur at port site. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). There is a low potential for this species to be present, as no suitable nesting sites exist in the area. According to SPRAT (2014) the main habitat range is in south-west Western Australia. Therefore it is unlikely this species would occur in the development area. | Entanglement in marine debris; disturbance; oil Spills; pollution | This species is not considered to frequent the Spencer Gulf area as its primary habitat is in south-western Western Australia and therefore its presence is unlikely. As the development does not represent critical habitat impacts to this species from the construction and operation of the port are not considered significant. | N | |
| Diomedea exulans antipodensis | Antipodean albatross | Vulnerable, Migratory, Marine (EPBC) | Unlikely to occur at port site. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). The Antipodean albatross is endemic to New Zealand, and forages widely in open water in the south-west Pacific Ocean, Southern Ocean and the Tasman Sea, notably off the coast of NSW (SPRAT, 2014). This species is not known to frequent South Australian waters and it unlikely to be present at the port site, as no suitable nesting sites exist in the area. | Entanglement in marine debris; disturbance; oil spills; pollution | This species is not considered to frequent the Spencer Gulf area as its primary foraging habitat is in south-eastern Australia and New Zealand therefore its presence is unlikely. As the development does not represent critical habitat impacts to this species from the construction and operation of the port are not considered significant. | N | |
| Diomedea exulans exulans | Tristan albatross | Endangered, Migratory, Marine (EPBC) | Unlikely to occur at port site. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). This species has a large feeding range covering the sub Antarctic and Southern Ocean with breeding occurring only on Gough Island and Inaccessible Island in the Atlantic Ocean, with its main foraging area in open water near Cape of Hope, South Africa | Entanglement in marine debris; disturbance; oil spills; pollution | This species has a large range covering the entire southern Australian coastline and offshore in the Southern Ocean. Therefore, while it may have the potential to overfly the development, the area will not represent a significant area of the species' foraging grounds or range. Impacts to this species from the construction and operation | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|------------------------|---|--|--|--|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | (SPRAT, 2014). The species is very large with a wingspan of 3.5 m making it rather conspicuous, given there are no records of the species near the development area and its preferred habitat is the Southern Ocean, it is unlikely to frequent Spencer Gulf. | | of the port are not considered significant. | | |
| Diomedea exulans gibsoni | Gibson's albatross | Vulnerable, Migratory, Marine (EPBC) | Unlikely to occur at port site. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). According to SPRAT (2014) distribution of this species is predominantly the Tasman Sea between NSW and New Zealand with all breeding occurring on offshore islands south of New Zealand. With no breeding sites in Australia and foraging areas around New Zealand it is considered highly unlikely this species would be found near the development area. | Entanglement in marine debris; disturbance; oil Spills; pollution | This species is not considered to frequent the Spencer Gulf area as its primary habitat is in south-eastern Australia and offshore islands near New Zealand. Therefore as its presence is unlikely impacts to this species from the construction and operation of the port are not considered significant. | N | |
| Diomedea exulans (sensulato) | Wandering albatross | Vulnerable, Migratory, Marine (EPBC); Vulnerable (NPW) | Unlikely to occur at port site. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). This species has a large feeding range as a circumpolar species nesting on Macquarie and Herald Island, Antarctic (SPRAT, 2014). This species prefers pelagic areas for foraging (SPRAT, 2014). Therefore, it is unlikely this species would frequent Spencer Gulf or overfly the development area. | Entanglement in marine debris; disturbance; oil Spills; pollution | This species has a large feeding range covering the entire southern Australian coastline and offshore in the Southern Ocean. Therefore, the port area will not represent a significant area of this species' foraging grounds or range. Impacts to this species from the construction and operation of the port are not considered significant. | N | |
| Eudyptula minor | Little penguin | Marine (EPBC) | Likely that species occurs in the area. This species is known to nest on Lipson Island (Edyvane 1999a; b; Madden-Hallett et al., 2011) 5 km to south of the development area. The BDBSA indicates a sighting in 1999 at Port Neill to the north (BDBSA, 2013). Given that the adult birds' foraging range is up to 20 km (Seraux et al., 2011) it is likely that the species will frequent the port site from the Lipson Island colony. This species is distributed along the southern coast in Australia from Perth to Brisbane, including Tasmania and offshore islands. Little Penguins are also found in New Zealand (SPRAT, 2014). Breeding burrows are found on rocky cliffs and vegetated sand dunes, usually on islands but also along remote beaches (Simpson and Day, 1999). | Disturbance of foraging during construction; underwater noise impacts during construction; entanglement with marine debris; feral predators or dogs; oil spills; | The population at Lipson Island is likely to be impacted by the presence of the port. However the species is found along Australia's southern coast and New Zealand (SPRAT, 2014) with breeding colonies in several other areas. The port site does not represent a significant area of habitat or foraging area and therefore impacts from the development on this species as a whole are not expected to be significant. | N | |



| Protected Species p | otentially present at t | he port | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|----------------------------|-------------------------|---|---|--|--|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | pollution | | | |
| Falco cenchroides | Nankeen kestrel | Marine (EPBC) | Present. This species was sighted within the development area in 2011 during the Jacobs port site flora and fauna survey. There are also a number of records of this species both at Port Neill and near Lipson Cove from 1998-2008 (BDBSA, 2013). This species is widespread across all of Australia including Tasmania and other offshore islands (SPRAT, 2014). | Loss of habitat through clearance or disturbance | This species breeds throughout Eyre Peninsula, Kangaroo Island, most of eastern SA, VIC and NSW and sparsely in WA, QLD, NT and TAS with a preference for most terrestrial habitats except dense woodland (Simpson and Day, 2010). As the port site does not represent a large geographic area of critical habitat for the species, impacts from the development on this species are not expected to be significant. | N | |
| Haematopus fuliginosus | Sooty oystercatcher | Rare (NPW) | Likely that species occurs in the area. There is the potential for this species to overfly or forage within the development area. This species was recorded during the Lipson Island flora and fauna study (Madden-Hallett et al., 2011). While it was not sighted during the flora and fauna survey at the site (Jacobs, 2014a) it has been recorded at Port Neill and Lipson Island in the years 1982-2008 (BDBSA, 2013) north and south of the development area. This species prefers rocky coastline; Cape Hardy includes some rocky outcrop areas that this species may use for foraging. Therefore the presence of this species at the port is considered likely. | Entanglement in marine debris; loss of habitat through clearance or disturbance; oil spills; pollution; feral predators or dogs | This species is distributed across all coastal areas of Australia (Simpson and Day, 1999). It nests on offshore islands and forages on adjacent rocky shores (Caton et al., 2011a). Despite the potential for this species to be present at the port site it is highly unlikely the port site would represent a significant area of this species' foraging grounds or range. Therefore, despite its potential presence there will be no significant impact on it or its habitat. This species is not an EPBC listed species and therefore not a matter of National environmental significance. Impacts from the development on this species are not expected to be significant | N | |
| Haematopus Iongirostris | Pied oystercatcher | Rare (NPW) | Likely that species occurs in the area. There is the potential for this species to overfly or forage within the development area as sightings have been recorded at Port Neill between 1998-2001 (BDBSA 2013) along the coast, north-east of the development. This species prefers sandy beach coastline or estuaries; Cape Hardy includes sandy beach areas that this species may use for foraging. | Entanglement in marine debris; loss of habitat through clearance or disturbance; oil spills; pollution; feral | The distribution of this species includes southern New Guinea and all coastal areas in Australia except the rocky cliffs of the Great Australian Bight (Geering et al., 2008). The species is found along most sandy beaches on Eyre Peninsula (Caton et al., 2011a). Despite the potential of this species to be present at the port site it is highly unlikely the port site would make up a significant area of its foraging grounds or range. This species is not an EPBC listed species, and | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|-----------------------------|---|--|--|---|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | predators or dogs | therefore despite its likely presence, disturbance to this species is not considered a matter of National environmental significance. Impacts from the port on this species are not expected to be significant | | |
| Haliaeetus leucogaster | White-bellied sea- eagle | Migratory, Marine (EPBC); Endangered (NPW) | Present. This species was recorded by Jacobs during the 2011 port site survey, overflying the development area. The BDBSA also has a number of records for this species in 1998-2008 at locations north and south of the port site at Port Neill, Lipson Island and Lipson Cove. Therefore, it is highly likely that this species will be seen in the vicinity of the harbour. The nests of this species are conspicuous as it prefers nesting on elevated platforms, poles or trees and there are no known breeding sites for this species within 10km of the port (BDBSA, 2013). | Entanglement in marine debris; loss of habitat through clearance or disturbance; oil spills; pollution | This species is migratory and has a large range covering the entire Australian coastline and many inland areas (SPRAT, 2014; Simpson and Day, 1999). The species is not known to nest in the port area and does not tend to establish new nests near settlements. Therefore the port site does not represent critical habitat or significant areas of its geographical range. Impacts from the development on this species are not expected to be significant | N | |
| Halobaena caerulea | Blue petrel | Vulnerable, Marine (EPBC) | Unlikely that species occurs in the area. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). This species is known to predominantly occupy sub-Antarctic open-ocean habitats, and does not tend to range far from breeding colonies in sub-Antarctic territory. This species is an uncommon visitor to Australian waters (Simpson and Day 2004, DEWHA 2010) with just a few records of sightings in the Great Australian Bight between Kangaroo Island and Esperance (SPRAT, 2014). Therefore, it is unlikely this species would frequent Spencer Gulf or overfly the development area. | Entanglement in marine debris; loss of habitat through disturbance; oil spills; pollution | This species is not considered to be directly reliant upon habitat near the port site as this species is an uncommon visitor to Australia and has a large feeding range in sub-Antarctic openocean. This species is unlikely to overfly the port and therefore the port site would not represent a significant area of its foraging grounds or range. Impacts from the development on this species are not expected to be significant | N | |
| Hydroprogne caspia | Caspian tern | Migratory; Marine (EPBC) | Present. There are no known nesting sites for this species at the port site however this species was sighted in the development area during the 2011 port site flora and fauna survey (Jacobs, 2014a), and a number of BDBSA records show sightings both at Port Neill and Lipson Cove from 1966-2009 (BDBSA, 2013). | Entanglement in marine debris; oil spills; pollution; loss of habitat | This species is found across Australia's coastal areas including inland waterways (Simpson and Day, 1999). Breeding of this species is known to occur in all Australian States and NT with breeding colonies found in SA from the Coorong to Ceduna and inland at Lake Eyre and Lake | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of si | Likelihood of significant impacts | | | |
|---|---------------------------|---|--|--|--|-----------------------------|--|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | | |
| | | | | through clearance or disturbance | Goyder (SPRAT, 2014). Given the extent of the species' distribution and breeding sites in Australia and the lack of identified breeding sites at Cape Hardy, the port site does not represent critical habitat or significant areas of its geographical range. Impacts from the development on this species are not expected to be significant | | | |
| Tringa brevipes listed marine as Heteroscelus brevipes | Grey-tailed tattler | Migratory, Wetland, Marine (EPBC); Rare (NPW) | Unlikely that species occurs in the area. There is one recorded sighting for this species at Port Neill in 2000 (BDBSA, 2013) but species was not sighted during the flora and fauna surveys at the port site in 2011 (Jacobs, 2014a). According to the SPRAT fact sheet for the species "the species is uncommonly recorded along the coasts" of South Australia (SPRAT, 2014). This species' preferred habitat is tidal creeks and mudflats. Therefore, as there is no suitable habitat at Cape Hardy, the species is unlikely to frequent the port site and any individuals that visit the area would be transients merely overflying the site. | Entanglement in marine debris; disturbance; oil spills; pollution | As a migratory species it has a large range covering numerous sections of the Australian coastline. Despite the potential for this species to occur near the development, the species has an extensive range of habitat. The development will not represent a significant area of the species with no suitable foraging grounds. Impacts from the development on this species are not expected to be significant | N | | |
| Hirundapus caudacutus | White-throated needletail | Migratory, Marine (EPBC) | Unlikely that species occurs in the area This species was last sighted in 1984 at Port Neill, north of the site (BDBSA 2013) and was not observed during the port site flora and fauna surveys. This species' range is extensive in the east of Australia but is unlikely to occur on the Eyre Peninsula (SPRAT, 2014). The Australian distribution of this species is generally further east along the eastern seaboard (Simpson and Day 2004). This species uses forested coastal and mountain habitats, as well as farmland and orchards. This species is not considered to be directly reliant upon habitat near the study area and therefore is unlikely to be present. | Disturbance; pollution | This species is not considered to exist in the development area due to its preferred habitat not being present. Considering the extensive habitat range in eastern Australia for this species; Impacts from the development on this species are not expected to be significant | N | | |
| Larus novaehollandiae | Silver gull | Marine (EPBC) | Present. This species was sighted within the development area during the 2011 site survey (Jacobs, 2014a) and is known to breed on Lipson Island (Madden-Hallett <i>et al.</i> , 2011). There are numerous BDBSA records for the species inland and along the coast from Port Neill to Lipson Cove dating from 1947-2009 (BDBSA, 2013). The species is extremely common across Australia, New Zealand and New Caledonia (Simpson and Day, 1999). This | Attraction to infrastructure for roosting or foraging; leading to entanglement in marine | This species is widespread and there are no nesting sites within 5 km of the development area. Even though this species is likely to forage in the area, it is known to exploit wharfs and even rubbish grounds for roosting and foraging, impacts from the development on this species are not expected to be significant | N | | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|---------------------------|---|--|--|--|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | species will utilise virtually any well-watered habitat and is highly adapted to exploiting human settlement, nesting on offshore islands or isolated cliffs (Simpson and Day, 1999). | debris; loss of habitat through clearance or disturbance; oil spills; pollution | | | |
| Larus pacificus | Pacific gull | Marine (EPBC) | Present. This species was sighted within the development area during the 2011 site survey (Jacobs, 2014a). There are also a number of records for this species inland and along the coast from Port Neill to Lipson Cove from 2000-2009 (BDBSA, 2012). This species is endemic to southern Australia, occurring along the coast from Exmouth in WA to Newcastle in NSW and TAS including offshore islands, with an isolated population in southern QLD and is considered moderately common (Simpson and Day, 1999). They generally nest on islands or headlands but their nesting is easily disturbed by human activities. | Attraction to infrastructure for roosting or foraging; leading to entanglement in marine debris; loss of habitat through clearance or disturbance; oil spills; pollution | There are no known breeding sites for this species within 5 km of the port site. This species is known to exploit wharfs and even rubbish grounds for roosting and foraging, so impacts from the development on this species are not expected to be significant. | N | |
| Leipoa ocellata | Malleefowl | Vulnerable, Migratory (EPBC); Vulnerable (NPW) | Unlikely to occur in area. Suitable habitat (large areas of mallee canopy coupled with sandy soils and dense leaf litter) is not found within the port site. Even inactive malleefowl mounds are easily identified and persist in the environment for several years, so their absence during the Jacobs port flora and fauna surveys along with the review of BDBSA records indicate it is unlikely the species utilises the area. The nearest record of species was over 12 km north-east from the site in 1999 (BDBSA, 2013). | Disturbance; vehicle strike, pollution | Suitable habitat for this species does not exist within the development area. Therefore, no loss of habitat will occur through development clearance. Impacts from the development on this species are not expected to be significant. | N | |
| Macronectes giganteus | Southern giant- petrel | Endangered, Migratory, Marine (EPBC); Vulnerable (NPW) | Unlikely to occur in area. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). This species is commonly seen in Australian oceanic waters and following ships (Simpson and Day, 1999). Therefore, given the existing ship traffic in Spencer Gulf the lack of records for the species in the gulf indicates that it is unlikely to be present at the | Entanglement in marine debris; loss of habitat through disturbance; oil spills; | This species is migratory and has a very large range covering the Southern Ocean, all Australian coastline and offshore areas (SPRAT, 2014). There is no known breeding occurring on the Australian mainland, rather all breeding occurs on sub-Antarctic islands (SPRAT, 2014). Therefore, the development will not represent | N | |



| Protected Species p | otentially present at t | he port | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|------------------------|---------------------------|---|--|--|---|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | port site. | pollution | critical habitat or a significant area of the species foraging grounds or range. Impacts from the development on this species are not expected to be significant. | | |
| Macronectes halli | Northern giant- petrel | Vulnerable, Migratory, Marine (EPBC) | Unlikely to occur in area. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). This species has an extensive range covering Australia's temperate coastal and offshore regions (SPRAT, 2014) with similar habitat and behaviour to the Southern species. (Simpson and Day, 1999). This species is considered to have only a low potential to overfly the development area. | Entanglement in marine debris; loss of habitat through disturbance; oil spills; pollution | This species is migratory and has a very large range covering the Australian coastline and offshore areas (SPRAT, 2014). Despite the potential for this species to over fly the development, the development will not represent a significant area of the species foraging grounds or range. Impacts from the development on this species are not expected to be significant. | N | |
| Merops ornatus | Rainbow bee- eater | Migratory, Marine (EPBC) | Unlikely to occur in area. This species has extensive migratory range and passes over much of Australia (excluding Tasmania) (SPRAT, 2014). These birds are vividly coloured making them quite conspicuous (Simpson and Day, 1999). There are no known nesting sites within 10 km of the development area (BDBSA, 2013). However, the species has been recorded in 1988 and 1998 in Port Neill. The species was not observed during Jacobs site surveys. The lack of records for the species in the last decade in the greater area indicates that this species is unlikely to be present in the development area. | Loss of habitat through disturbance; pollution | The proposed activities will not impact on this species or species habitat due to lack of suitable habitat in the development area. This species may have the potential to overfly the port site however it is highly unlikely the development would make up a significant area of its foraging grounds or range as this species range is extensive including most of Australia therefore impacts from the development on this species are not expected to be significant . | N | |
| Neophema petrophila | Rock parrot | Marine (EPBC); Rare (NPW) | Present. This species was sighted within the development area during the 2011 site survey (Jacobs, 2014a) with two records at Port Neill in 2000-2001 (BDBSA, 2013). This species is listed as Rare in SA but is considered reasonably common throughout its range (Simpson and Day, 1999). | Loss of habitat through clearance or disturbance; pollution | This species' population is distributed along Australia's coast from Exmouth in WA to the WA-SA border and from Fowlers Bay to Fleurieu and Kangaroo Island in SA including both gulfs (SPRAT, 2014). The species nests on offshore rocky islands and cliffs, visiting the adjacent coast (Simpson and Day, 1999). Despite the presence of the species at the site, critical habitat for this species exists outside of the development area and as the port will not directly impact on any coastal/offshore islands impacts from the development on this species is | N | |



| Protected Species potentially present at the port | | he port | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|------------------------|---|--|--|--|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | | not expected to be significant. | | |
| Nycticorax caledonicus | Nankeen night heron | Marine (EPBC) | Present. This species was sighted within the development area during the 2011 site survey (Jacobs, 2014a). These birds roost in trees or under wharfs by day and fish at night (Simpson and Day, 1999). The construction of the port may benefit the species by providing additional roosting areas (thus increasing the likelihood of occurrence at the site). | Attraction to infrastructure for roosting; leading to entanglement in marine debris; clearance of habitat; oil spills; pollution | This species is distributed along most of the Australian coast (except the Great Australian Bight) and eastern Australian inland areas (Simpson and Day, 1999). While there may be impacts on individuals of this species due to port operations, the species is common and its range large therefore impacts from the development on this species its habitat is not expected to be significant . | N | |
| Pandion haliaetus | Osprey | Marine (EPBC); Endangered (NPW) | Possible that species occurs in the area. There is potential for this species to be in the area as a juvenile bird was recorded in 2009 within 2 km of the port site and sighting records of the species exist from 2001 at Port Neill (BDBSA). There are no known breeding sites for this species within 10 km of the development area (BDBSA, 2013). These individuals can have a large feeding range of up to 80 km (SPRAT, 2014). It is therefore likely that visiting birds to the site would be vagrants or transient juveniles. | Loss of habitat through clearance or disturbance; entanglement in marine debris; oil spills; pollution (e.g. pesticides) | This species has an extensive habitat range including most of coastal Australia (Simpson and Day, 1999) and extending into New Guinea, Solomon Islands, Indonesia, Philippines, Palau Islands, and New Caledonia (SPRAT, 2014). This species has a large feeding range and although it may have the potential to overfly the development site it is unlikely the site would make up a significant area of its foraging grounds or range. Impacts from the development on this species or its habitat are not expected to be significant. | N | |
| Pelecanus conspicillatus | Australian pelican | Marine (EPBC) | Present. This species was sighted within the development area during the 2011 site survey (Jacobs, 2014a) with multiple records from 1998-2009 at Port Neill (BDBSA, 2013). This species is found throughout Australia, Papua New Guinea and western Indonesia, with occasional reports in New Zealand and various western Pacific islands. The Australian Pelican utilises diverse habitats including inland fresh and saline lakes, dams, rivers, suburban ponds, swamps, estuarine, wetlands, coastal shores and islands | Loss of habitat through clearance or disturbance; entanglement in marine debris; oil | This species is not considered threatened. Apart from habitat destruction the biggest threats to this species are oil spills, pesticide poisoning and damage breeding sites, which are sensitive to destruction by even a single dog (Barbara - personal observation 1993 at Outer Harbour, SA). The species has previously bred on the sand-spit isle at Outer Harbour, Port Adelaide | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|-------------------------------|---|--|--|---|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | (Australian Museum, 2013). Breeding colonies are widespread across Australia (Simpson and Day, 1999). The species breeding is dependent on environmental conditions with one of the largest ephemeral breeding areas being Lake Eyre during years of heavy rainfall. This species also breeds on several offshore islands. | spills; pollution (e.g. pesticides) | opposite the active port (Barbara -personal observation) and is commonly found at all ports throughout Australia. Therefore, the presence of a port does not adversely impact on its behaviour. As the species is capable of coexisting with human developments the construction and operation of the port at Cape Hardy is unlikely to significantly impact on the species. | | |
| Phalacrocorax fuscescens | Black-faced cormorant | Marine (EPBC)) | Likely that species occurs in the area. Several BDBSA records of this species exist south and north of the development area at Lipson Island and Port Neill from 1982-2008. Breeding for this species occurs on Lipson Island (Madden-Hallett et al., 2011) and foraging is likely to occur in the area of the port site. Breeding colonies for this species are widespread across Australia (Simpson and Day, 1999). The species' largest breeding colony is at South Australia's largest operating port, Port Adelaide on the Outer Harbour breakwater (Zoo SA, 2013). Therefore the presence of an active port is unlikely to negatively impact on the species. The species is known to exploit man-made constructions in the marine environment, utilising them for both roosting and fishing locations (Simpson and Day, 1999). | Interactions with vessels and construction machinery due to attraction to infrastructure for roosting; entanglement in marine debris; loss of habitat through clearance; oil spills; pollution | This species is not considered threatened and is found along much of Australia's southern coast (except perhaps the Great Australian Bight) (Simpson and Day, 1999). Considering the active breeding colony at Port Adelaide, this species is able to coexist with large operating ports without adversely impacting on its behaviour. Therefore, construction and operation of the port at Cape Hardy is unlikely to significantly impact on the species. | N | |
| Psophodes nigrogularis leucogaster | Western whipbird (eastern) | Vulnerable (EPBC); Endangered (NPW) | Unlikely to occur in area. The SA Museum has carried out targeted surveys for the Western-Whip bird on Eyre Peninsula in 2006-2007 to determine the species distribution and abundance (ZooSA, 2013) and did not identify the area as habitat for the species. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site (Jacobs, 2014a). The Western Whipbird (eastern) inhabits mallee and thicket vegetation in coastal and inland areas of southern South Australia and. is said to associate with the Red-lored Whistler (<i>Pachycephala rufogularis</i>) which is | Pollution | The population on the Eyre Peninsula is restricted to sites around Coffin Bay National Park and Lincoln National Park (EPBC, 2013; SPRAT, 2014). The Western Whipbird (eastern) is a sedentary bird (Condon 1966) that is capable of making only short-distance flights (Condon 1966). Its inability to traverse long distances in flight probably limits or prevents its dispersal across areas that have been cleared of suitable habitat (SPRAT, 2014). Therefore the species is | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|--|---|---|--|--|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Pterodroma mallic | Soft-plumaged | Vulnerable | listed as Vulnerable under the EPBC Act. It also occurs in areas inhabited by the Malleefowl (<i>Leipoa ocellata</i>) and Regent Parrot (eastern) (<i>Polytelis anthopeplus monarchoides</i>) both listed as Vulnerable under the EPBC Act and the Black-eared Miner (<i>Manorina melanotis</i>) and Mallee Emu-wren (<i>Stipiturus mallee</i>) listed as Endangered under the EPBC Act (SPRAT, 2014). Therefore as there are no records for the species in the greater area, its inability to fly great distances, no suitable floral habitat, nor associated avian species known to occur in the development area, it is considered unlikely the species would occur in the area. | Entanglament | not considered to occur in the area and impacts from the construction and operation of the port at Cape Hardy are unlikely to significantly impact on the species. | N | |
| Pterodroma mollis | Soft-plumaged petrel | Vulnerable, Marine (EPBC) | Unlikely to occur in area. There are no records for this species within 10 km of the study area (BDBSA, 2013) nor was it observed during the flora and fauna survey at the site ((Jacobs, 2014a). This species is not known to breed in Australian territory and inhabits sub-Antarctic oceanic areas (Simpson and Day 2004; Shirihai, 2007; DEWHA, 2010). This species is not considered to be directly reliant upon habitat near the study area. | Entanglement with marine debris; oil spills; pollution (e.g. poisoning from pesticides) | This species has a large feeding range covering temperate and sub Antarctic waters in the South Atlantic, southern Indian and western South Pacific Oceans (SPRAT, 2014) therefore the species has the potential to overfly the development area. However the only known breeding colony of the species is on an offshore island south of Tasmania with all other colonies spread across islands in the Southern Ocean from South Africa to South America (SPRAT, 2014). Given the extent of the species range and its absence of breeding colonies in South Australia, is unlikely that the port site would make up a significant area of its foraging grounds or range. Impacts from the construction and operation of the port are unlikely to significantly impact on the species. | N | |
| Puffinus carneipes = Ardenna carneipes | Flesh-footed Shearwater, Fleshy-footed Shearwater | Migratory, Marine (EPBC) | Unlikely to occur in area. This species was not observed during the flora and fauna survey at the site (Jacobs, 2014a). One log describing a group of 15 birds of unknown <i>Puffinus</i> sp. was recorded in 2009, with behaviour described as "feeding near dolphins", approximately 10 km south of the development area (BDBSA 2013). As the sighting did not identify the birds beyond genus it is possible the <i>Puffinus</i> sp. recorded were Fleshy-footed shearwaters, however it is considered unlikely as the nearest colony is over 90 km away and a similar species the Short-tailed | Entanglement with marine debris; oil spills; pollution (e.g. poisoning from pesticides) | The Puffinus carneipes or Ardenna carneipes is a trans-equatorial migrant widely distributed across the Indian and Pacific Oceans with a global population of up to 380000 pairs nesting throughout the globe (SPRAT, 2014). This species is considered abundant throughout its range in Australia (Simpson and Day, 1999). The species is known to breed on 41 offshore islands within Australia with South Australia's key | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence Like | | Likelihood of significant impacts | | | |
|--|----------------------------|--|---|--|---|-----------------------------|--|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | | |
| | | | Shearwater is thought to roost on Lipson Island (Madden-Hallett et al., 2011) and is therefore a more likely candidate for the sighting | | breeding site located on Smith Island near Port Lincoln (SPRAT, 2014) > 90 km to the south of the development area. As the species has a large feeding range it has the potential to overfly the development area however it is unlikely that the port site represents a significant area of this species' foraging grounds or critical habitat. Impacts from the construction and operation of the port are unlikely to significantly impact on the species. | | | |
| Puffinus tenuirostris = Ardenna tenuirostris | Short-tailed shearwater | Migratory, Marine (EPBC) | Possible that species occurs in the area. This species was not observed during the flora and fauna survey at the site (Jacobs, 2014a) however there is a log describing a group of 15 birds of unknown <i>Puffinus</i> sp. was recorded in 2009, with behaviour described as "feeding near dolphins", approximately 10 km south of the development area (BDBSA 2013). It is possible the <i>Puffinus</i> sp. recorded were Short-tailed Shearwaters that are thought to roost on Lipson Island (Madden-Hallett <i>et al.</i> , 2011). | Entanglement with marine debris; oil spills; pollution (e.g. poisoning from pesticides) | The species is found from Perth to Brisbane in coastal and offshore waters and is classified as abundant in Australia (Simpson and Day, 1999). This species is thought to roost and potentially breed on Lipson Island (Madden-Hallett et al., 2011) 5 km south of the development area. As the species has a large feeding range it has the potential to overfly the development area however it is unlikely that the port site represents a significant area of the species' global foraging grounds or range. Impacts from the construction and operation of the port are unlikely to significantly impact on the species. | N | | |
| Rostratula australis | Australian painted snipe | Endangered Migratory, Marine (EPBC); Vulnerable (NPW) | Unlikely to occur in area. The Australian painted snipe generally inhabits shallow terrestrial freshwater (occasionally brackish) wetlands, including temporary and permanent lakes, swamps as well as inundated or waterlogged grassland or saltmarsh, dams, rice crops, sewage farms and bore drains (SPRAT, 2014), habitat that is not present at the port site. There are no records for this species within 10 km of the development (BDBSA, 2012) and the species was not observed during the flora and fauna survey at the site (Jacobs, 2014a). As there are no records for the species in the wider area and with no suitable habitat at the port site the species is not expected to occur in the area. | Entanglement with marine debris; oil spills; pollution (e.g. poisoning from pesticides) | The species' range includes the Pilbara, Kimberley's, most of NT, QLD, NSW, Victoria and eastern South Australia to the Eyre Peninsula, with an isolated population in south west WA (SPRAT, 2014). Breeding occurs in Victoria, NSW and South Australia's south east its range does not include Eyre Peninsula (Simpson and Day, 2010). As the port site does not include suitable habitat and given the species prefers eastern and northern areas of Australia it is not expected to frequent the development area and therefore impacts from the construction and operation of the port are unlikely to significantly impact on | N | | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|----------------------------|---|--|---|---|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | | the species. | | |
| | | | | | | | |
| Haematopus fuliginosus | Crested tern | Migratory, Marine (EPBC); Rare (NPW) | Likely that species occurs in the area. This species is known to nest on Lipson Island (Madden-Hallett <i>et al.</i> , 2011), and sightings records have been logged at Port Neill and the Lipson Cove area from 1982-2009 (BDBSA, 2012). The species is known to use coastal seas and continental shelf habitats (Simpson and Day 2004) with an extensive range it is likely the species will be found at the port site. | Loss of habitat through clearance or disturbance; entanglement with marine debris; oil spills; pollution (e.g. poisoning from pesticides) | This species is considered common throughout its range in all Australian coastal waters and into South East Asia (Simpson and Day, 1999). This species is known to congregate on coastal shores and intertidal zones. Although the species was not observed during the field survey (Jacobs, 2014a), suitable habitat occurs within the coastal zones of the port site and it is considered likely that this species uses habitat available in the area. The species is highly mobile and is not considered likely to rely solely on habitat present within the port area. Therefore given its extensive range it is expected that impacts from the construction and operation of the port are unlikely to significantly impact on the species. | N | |
| Sterna fuscata = Onychoprion fuscata | Sooty tern | Marine (EPBC); | Unlikely to occur in the area. No records of sightings of this species have been submitted to the BDBSA within 10 km of the | Entanglement with marine | There is no evidence that this species has been sighted within 10 km of the development area | N | |
| | | Rare (NPW) | development area (BDBSA, 2013). This species was not observed during the flora and fauna survey in 2011 (Jacobs, 2014a) or the targeted flora and fauna survey of Lipson Island (Madden-Hallett et al., 2011) where it was thought to breed (EPBC search results Nov., 2013). This species is generally pelagic but also occupies islands. In Australia this species' distribution is concentrated on the northern aspects of the continent, outside of Spencer Gulf or southern Australia (Simpson and Day 2004). | debris; oil spills; pollution | (BDBSA, 2013). The species' range includes coastal and offshore waters of Australia but excludes southern coasts (Simpson and Day, 1999). Its presence in the development area is therefore unlikely. Impacts from the construction and operation of the port are unlikely to significantly impact on the species. | | |
| Sternula nereis nereis | Fairy tern (Australian) | Vulnerable (EPBC); Endangered (NPW) | Possible to occur in the area. This species was not observed during Jacobs' port site surveys in 2011, or the flora and fauna survey of Lipson Island (Madden-Hallett <i>et al.</i> , 2011). However the species is thought to breed on Lipson Island Conservation Park 5 km to the south of the development area despite recent surveys failing to find the species (Madden-Hallett <i>et al.</i> , 2011). There is only one record submitted to the BDBSA from Port Neill, north of | Loss of habitat through clearance or disturbance; entanglement with marine | Although there is potentially suitable habitat at the port site for this species there are other suitable beaches within Spencer Gulf and across Australia. While presence of this species at the site is considered possible and due to the geographic range of the species, the port site does not represent a significant area of critical | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|---|--|---|--|--|--|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | the development area in 1998 (BDBSA, 2013). This species nests on sheltered sandy beaches, sand spits and banks between the high tide line and vegetation along Southern Australia from Victoria and Tasmania to as far up as Dampier in Western Australia (SPRAT, 2014). The habitat type exists at the port site along with several nearby locations within Spencer Gulf. Given the threatened status of this species and suitable habitat in the area it is considered possible the species occurs in the area. | debris; oil spills; pollution; predation by feral animals or dogs | habitat or known breeding location. Impacts from the construction and operation of the port are considered unlikely to significantly impact on the species. | | |
| Thalassarche bulleri | Buller's albatross | Vulnerable, Migratory, Marine (EPBC) | Unlikely to occur in area. This species was not observed during Jacobs port site surveys in 2011 and no records have been submitted to the BDBSA within 10 km of the development (BDBSA, 2012). This species has a large feeding range from New Zealand to the coasts of south eastern Australia (Simpson and Day, 1999). Therefore the range of this species is outside of the development area it is unlikely that the port site would represent critical habitat for this species. | Entanglement with marine debris; oil spills; pollution | This species is endemic to New Zealand where it breeds (SPRAT, 2014) and has a very large foraging range covering the south-eastern Australian coastline including Tasmania. The location of the development does not represent a significant area of the species foraging grounds or range. Impacts from the construction and operation of the port are considered unlikely to significantly impact on the species | N | |
| Thalassarche cauta cauta For marine and migratory listed as Thalassarche cauta (sensustricto) | Shy albatross, Tasmanian shy albatross | Vulnerable, Migratory, Marine (EPBC) | Unlikely to occur in area. This species was not observed during Jacobs' port site surveys and no sighting records have been submitted to the BDBSA within 10 km of the development (BDBSA, 2012). The recent separation of the Shy albatross from other closely related taxa confounds the current understanding of its at-sea distribution; it appears to occur over all Australian coastal waters below 25° S (SPRAT, 2014). However this species is most commonly observed over the shelf waters around Tasmania and south-east Australia, with all breeding occurring around Tasmania (SPRAT, 2014). The development area is unlikely to represent a significant area of its range or critical foraging grounds and it is considered unlikely the species would occur at the port. | Entanglement with marine debris; oil spills; pollution | This species is endemic to south-eastern Australian and Tasmania where it breeds (SPRAT, 2014) and has a very large foraging range covering the south-eastern Australian coastline. The location of the development area of the port is not within the known range and does not represent key habitat type for the species. Impacts from the construction and operation of the port are unlikely to significantly impact on the species. | N | |
| Thalassarche cauta steadi | White-capped albatross | Vulnerable, Migratory, Marine (EPBC) | Unlikely to occur in area. No records have been lodged with the BDBSA within 10 km of the development (BDBSA, 2012) and the species was not observed during Jacobs' 2011 port site surveys. This species is similar in appearance to the Shy albatross, so sightings of the two are difficult to assign (SPRAT, 2014) however there are no records for either species in the area. This species is | Entanglement with marine debris; oil spills; pollution | This species is endemic to New Zealand where it breeds (SPRAT, 2014) and has a very large foraging range covering the Southern Ocean and Australian coastline including Tasmania. The location of the development does not represent a significant area of the species foraging grounds | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | | |
|--|----------------------------|---|--|---|---|-----------------------------|--|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | thought to be common off the coast of south-east Australia (SPRAT, 2014) and has a large range extending from South Africa to New Zealand where it breeds on offshore islands. No breeding occurs in Australian waters (SPRAT, 2014). | | or critical habitat. Impacts from the construction and operation of the port are considered unlikely to significantly impact on the species | | |
| Thalassarche melanophris | Black-browed albatross | Vulnerable, Migratory, Marine (EPBC) | Unlikely to occur in area. This species breeds on sub-Antarctic islands and is confined to the waters surrounding those islands during the breeding season with less than 1% of the global population breeding within Australian jurisdiction (SPRAT, 2014). In the non-breeding season it migrates north and its range expands to be truly circumpolar, and it is a common visitor to the continental shelf and shelf break in southern Australia (SPRAT, 2014). Only one 20 year old record has been submitted to the BDBSA, at Port Neill in 1989 (BDBSA, 2012) indicating the record is for a transient or lost individual. Any individual of this species in the port area would be considered a transient visitor. | Entanglement with marine debris; oil spills; pollution | This species has a large feeding range covering the Southern Ocean (Simpson and Day, 1999) and although it may have the potential to overfly the development area it is unlikely that the port site would make up a significant area of its foraging grounds or represent critical habitat. Impacts from the construction and operation of the port are considered unlikely to significantly impact on the species | N | |
| Thalassarche melanophris impavida For marine and migratory listed as Thalassarche impavida | Campbell albatross | Vulnerable, Migratory, Marine (EPBC) | Unlikely to occur in area. This species is a migratory non-breeding visitor to Australian waters only breeding on the sub-Antarctic Campbell Island south of New Zealand (SPRAT, 2014). Its non-breeding range is very large extending into the sub-tropics including offshore waters from Rockhampton south around to Exmouth (SPRAT, 2014). However there are no records for the species within 10 km of the development area (BDBSA, 2012). This species would be considered a transient visitor to the area. | Entanglement with marine debris; oil spills; pollution | This species has a large feeding range (SPRAT, 2014) and although it may have the potential to overfly the development area it is unlikely that the port site would make up a significant area of its foraging grounds or represent critical habitat. Impacts from the construction and operation of the port are unlikely to significantly impact on the species | N | |
| Thinornis rubricollis rubricollis | Hooded plover (eastern) | Vulnerable, Marine (EPBC); Vulnerable (NPW) | Present. A hooded plover pair was observed on the southern end of the Cape Hardy beach during the port site survey (Jacobs, 2014a). This species has also been recorded on beaches north and south of the development area (BDBSA, 2013; Madden-Hallett et al., 2011). There is suitable habitat for this species in a number of beach areas within Spencer Gulf including Lipson Island. The birds nest at the high tide mark on sandy beaches, laying usually two eggs in a sand scrap and foraging for invertebrates at the water's edge and amongst seagrass wrack. Pairs are known to have breeding territories over 30 ha in size with several nesting beaches within a territory (SPRAT, 2014). This species' range in Australia is generally along coastal sandy beaches from Victoria | Loss of habitat through clearance or exclusion; predation by feral animals or pet dogs; marine debris; off road vehicles, oil spills; | The construction of the wharf and MOF will impact on the head land at Cape Hardy removing intertidal habitat for this species. Given the species preference for remote beaches, mitigation measures to minimise disturbance to the beach habitat during construction of the port will be required, with exclusion zones on the southern extent of the Cape Hardy beach to be included in the CEMP and OEMP to be developed. Given the species ability to relocate to alternate beaches within their territory during construction disturbance | N | |



| Protected Species potentially present at the port | | | Likelihood of Occurrence | Likelihood of significant impacts | | |
|---|-------------------|---|--|---|--|-----------------------------|
| Species Name | Common Name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | | and Tasmania, throughout South Australia and including some inland salt-lakes populations in WA (Simpson and Day, 1999). The Eyre Peninsula population is not considered an important population by the Threatened Species Conservation Committee (SPRAT, 2014). | pollution, noise | and the limited footprint of the MOF, the development is not expected to have a significant impact on this species. | |
| Reptiles | | | | | | |
| Lerista arenicola | Beach slider | Rare (NPW) | Possible to occur in area. There are no known sightings for this species in the area; all sightings have been recorded on the far west coast of the Eyre Peninsula (Caton et al., 2010). However, the species is inconspicuous and easily missed amongst the seagrass wrack and other detritus around the high tide mark where they forage therefore the lack of records cannot be considered confirmation of presence or absence for this species. As this species is known to inhabit the seaweed wrack found on sandy beaches and that habitat is known to occur at Cape Hardy beach the EP Coastal Action Plan 2010 potential habitat for this species (Caton et al., 2010a) | Loss of habitat through clearance or exclusion; predation by feral animals or pet dogs; marine debris; off road vehicles, oil spills; pollution | Given the species is not known from the area and is has been recorded predominantly on the west coast of Eyre Peninsula the development is not considered to represent a significant area of critical habitat and therefore impacts from the development on this species are not considered to be significant. As a precaution to protect this species and similar intertidal foragers such as the Hooded Plover and Bight Coast skink, a CEMP and OEMP will be developed to protect these species. | N |
| Pseudemoia pagenstecheri | Bight Coast skink | Rare (NPW) | Possible to occur in area. There are no known sightings for this species in the area; all sightings have been recorded on the far west coast of the Eyre Peninsula (Caton et al., 2010). However, the species is inconspicuous and easily missed amongst the seagrass wrack and other detritus around the high tide mark where they forage therefore the lack of records cannot be considered confirmation of presence or absence for this species. As this species is known to inhabit the seaweed wrack found on sandy beaches and that habitat is known to occur at Cape Hardy beach the EP Coastal Action Plan 2010 potential habitat for this species (Caton et al., 2010a) | Loss of habitat through clearance or exclusion; predation by feral animals or pet dogs; marine debris; off road vehicles, oil spills; pollution | Given the species is not known from the area and is has been recorded predominantly on the west coast of Eyre Peninsula the development is not considered to represent a significant area of critical habitat and therefore impacts from the development on this species are not considered to be significant. As a precaution to protect this species and similar intertidal foragers such as the Hooded Plover and Beach slider, a CEMP and OEMP will be developed to protect these species. | N |



6.1.2.2 Risks

There is a risk that the introduction of artificial light sources will result in altered behavioural patterns amongst coastal and marine fauna through the attraction of higher levels of marine prey species and marine birds than anticipated. The level of attraction and subsequent effect on resident species as a result of greater utilisation of habitat and foraging resource is largely unknown. It is considered possible that despite design measures to limit artificial light spill (refer Section 4), additional marine fauna will be attracted to the site. However, the port site does not represent a key habitat or large breeding colony for any bird species with resident populations limited to isolated pairs and individuals. As such, the introduction of artificial light sources is considered unlikely to affect the viability of any species. The artificial light sources may however result in a non-significant alteration to the behaviour of fauna at the port site which is considered to be of minor consequence. As such, the overall risk associated with the introduction of artificial light sources at the port site is considered to be low.

Construction and operation of the proposed port will result in the generation of a number of waste streams that may enter the marine environment. Marine debris can significantly affect marine birds and fauna as a result of pollution, injury through collision, entanglement or ingestion of non-biological products. The Silver Gull, Black-faced Cormorant and Nankeen Night Heron along with several other species considered "likely or possible" to occur in the area (refer Table 6-1) have all be highlighted as having the potential to be attracted to the port activities. Due to these species' characteristics of exploiting human activities in the marine environment, they are considered particularly susceptible to debris within the marine environment.

Waste management and handling procedures developed as part of the CEMP and OEMP will control waste streams with the overarching aim of no waste products entering the marine environment. Despite the implementation of control measures, it is considered almost certain that some form of waste / debris will enter the marine environment during construction and operation of the port. As the port site is not a key habitat or breeding area, the consequences of debris entering the marine environment are considered to be minimal; insignificant to the overall viability of marine birds and fauna. As such, the overall risk is considered to be low.

Marine birds and coastal species are also at risk from the unintended discharge of pollution in the form of site run-off, wastewater, hydrocarbons or chemicals. With any marine activity there is potential for waste or spills to enter the marine environment. The implementation of design measures (refer Section 4) will control hazardous pollutants from entering the marine environment. There remains a risk that a major spill of oil or other chemicals may occur as a result of vessel failure or accident. Based on the experiences of other operating ports, the likelihood of this event occurring is considered to be rare. As previously outlined, the port site does not represent a critical habitat or breeding colony for any marine bird. The nearest bird colonies are located on Lipson Island, approximately 5 km south of the port site, and foraging birds from these colonies (e.g. little penguins) are likely to use the waters around the port. The consequences of a major spill event are considered to be moderate, resulting in a long term decrease in the local abundance of fauna. As such, the overall risk of a major spill event affecting marine birds is considered to be low.

6.1.3 Conclusions

- The EPBC and NPW Vulnerable species eastern hooded plover are known to utlise the site. The majority
 of the bird's habitat is preserved by the design of the port which avoids the majority of the sandy beach
 habitat, thereby minimising risks to these species. Hyrdodynamic modelling predicts little change to the
 existing beach environment. CEMP and OEMP control strategies and measurement measures will
 minimise access to the beach
- While it is possible for the two NPW protected lizard species (beach slider and Bight Coast skink) to be present due to suitable beach habitat, the design of the port avoids the majority of the beach habitat thereby mitigating risks to these species
- No EPBC listed flora species were recorded during the field survey



- In total, 56 bird species of conservation significance were identified during the desktop study as potentially occurring in the project area, including species listed under the EPBC Act and the NPW Act. Thirty of these species are considered unlikely to occur in the area due to a lack of suitable habitat, or a known distribution that is distant from the proposed development area. Twelve species have potential to occur in the development area and surrounding region due to the presence of suitable habitat, regional BDBSA records or recent regional studies/biodiversity planning. Fourteen protected bird species were observed during the field surveys
- None of the 12 bird species of conservation significance with potential to occur in the development area are expected to be significantly impacted. If local individuals occur they may be displaced however all of these species are highly mobile and unlikely to be solely reliant on habitat within the study area.
- As several species of marine birds may be attracted to the port operations there is a potential for impacts on individual birds, impacts to these populations are considered low
- Of the 14 bird species of conservation significance recorded in the study area, two are listed as Migratory under the EPBC Act (Caspian tern and common sandpiper) and one is listed as Vulnerable (Hooded Plover). The 11 other species are listed as "Marine" of which two other species (rock parrot and white-bellied sea-eagle) are also listed as threatened under the State NPW Act
- There are no known large breeding colonies within 5 km of the proposed development area and only the one species of shorebird (hooded plover) is known to nest at the site. Therefore this site is not considered an internationally or nationally significant shorebird habitat under the Ramsar Convention

6.2 Intertidal Habitats

The intertidal marine environment of the Spencer Gulf is varied, supporting a wide range of habitats and species groups. Species in the lower reaches of the gulf are generally associated with three key intertidal habitat groups: seagrass meadows, rocky reefs and soft substrates. A clear understanding of which of these habitats and species are present is necessary to assess the biological importance of the site, identify any species that trigger further regulatory requirements, and any species or habitats requiring implementation of specific management or mitigation measures. Reefs and seagrass meadows provide essential habitat to a wide range of marine species such as fish, crabs and sea urchins, while mature seagrass meadow also stabilise the seabed to prevent erosion and sand movement.

Previous studies indicate that within the intertidal zone, Spencer Gulf seagrass meadows are dominated by *Zostera* spp. and *Heterozostera nigricaulis*, with other areas either bare or dominated by macroalgal assemblages including areas of rocky shores (Edyvane, 1999a; 1999b). The intertidal extents of the coastline from Port Neill to Tumby Bay are known to include sandy and rocky substrates, moderate wave exposure, and no mangroves; with estuary and saltmarsh habitat restricted to the Tumby Bay area. Rocky substrates occur predominantly at headlands, but also as discrete boulder areas scattered around sandy beaches. Sandy substrates in the area were reported to support seagrasses of varying density (Nature Maps), between approximately 19 m of depth and the near-shore up to the intertidal zone with areas of macroalgae.

A desktop study characterised the intertidal habitats present at the site to inform the assessment of the range of associated species likely to be present at the site.

6.2.1 Existing Environment and Key Findings

The results of the EPBC PMST found that no Threatened Ecological Communities occurred within 10 km of the study area. Previous literature reviews did not highlight any significant or unique intertidal habitats as being likely at Cape Hardy (Caton *et al.*, 2011a; 2011b). The desktop review suggested that the intertidal area of Cape Hardy is likely to be typical of southern Australia and in particular the Spencer Gulf. As noted in Section 6.1, there are no mangrove communities within the Cape Hardy study area.



The intertidal habitat at Cape Hardy contains both sandy beaches and rocky intertidal zones both as gneiss and gabbro outcroppings on the sandy beaches and rocky intertidal reefs extending from the headlands into the subtidal zone (MES, 2012). Marine flora and fauna of the lower Spencer Gulf is typical of transitional warm to cool temperate waters (i.e. Flindersian Province). In sheltered areas of Spencer Gulf, intertidal flats are typically dominated by *Hormosira banksii* and potentially the seagrasses, *Zostera (Heterozostera) nigricaulis* and *Zostera muelleri*. (Note: *Heterozostera nigricaulis* was distinguished from *Heterozostera tasmanica* in 2005 based on morphological features and distribution (Kuo, 2005) with some taxonomists preferring to retain the subgenus *Zostera* (Jacobs and Les, 2009). For the purposes of this assessment we have used the *Heterozostera nigricaulis* classification). The areas of rocky reef and shore are typically dominated by macroalgal communities including *Scaberia agardhii*, *Osmundaria* spp., *Lobophora variegata* and *Sargassum* spp. in low wave energy areas, and species of *Cystophora* (e.g. *C. expansa*) on moderate-energy coasts. Sandy areas are either bare or dominated by additional algal species such as *Caulocystis* spp., *Cystoseira* spp., *Ecklonia radiata* and *Sargassum* spp., usually with an understorey of coralline algae (Edyvane, 1999a, 1999b).

The macroalgal habitats and rocky areas also support invertebrate demersal species including relatively sedentary chitons and gastropods as well as highly mobile foragers including crabs and fish species. Other invertebrates may exist either permanently attached to surfaces such as barnacles, sponges, bryozoans or as infauna species such as cockles, clams, tube worms or small arthropods. Typical invertebrate species previously identified along the Spencer Gulf Eyre Peninsula coast include *Austrocochlea* spp., *A. unifasciata*, *Bembicum* spp., *Notocamea* spp., *Patellaida latisrigata*, *P. alticostata*, *Patella chapmanii*, *Plaxiphora albida*, *Siphionaria diemenensis* and *S. zelandica* (Golders, 2012).

The EPBC search and desktop literature review did not identify any rare or protected species known to occur in the intertidal zones at Cape Hardy. The seagrass subspecies *Zostera muelleri* ssp. *mucronata* is listed as rare and protected under the NPW Act, however it is considered unlikely to exist at the Cape Hardy site as there are no records of it within 50 km and its preferred habitat is very low-energy intertidal sandy beaches which are not found in the study area.

As the intertidal reef and rocky shore can provide habitats for species the extent of these areas within the marine study area was estimated using a conservative approach (i.e. erring on the side of overestimation) as 2.3 ha, based on the aerial imagery and bathymetry data from the study area. This approach allowed a buffer around these habitat extents and guided design of the port site infrastructure to avoid as much of these habitats as practicable.

6.2.2 Impacts and Risks: Intertidal Habitats

Most of the intertidal rocky shore and sandy beaches in the study area are outside the port infrastructure footprint and not expected to be directly impacted by the development. The area of intertidal habitat to be impacted is all rocky shore and will be permanently covered by the MOF and tug harbour which incorporates the entire footprint including the road landing beginning of the jetty. A conservative estimate of intertidal zone to be lost beneath the MOF is 0.09 ha.

Impacts to intertidal habitats associated with the proposed port facility could occur as a result of:

- Habitat loss or exclusion due to the construction of the rock armouring
- Erosion (from stormwater and vessel wakes/propwash, changes to hydrodynamics)
- Spills (e.g. oil), pesticides, herbicides, or other pollutants
- A build-up of contaminants (e.g. anti-foulants)
- Vehicle traffic on the shore
- Marine debris
- The introduction of IMS
- Vessel run aground



6.2.2.1 Impacts

As the construction of the jetty and wharf, and covering of seabed by the MOF and tug harbour, will extended directly from the headland there is no intertidal zone within the footprint. The existing substrate in the adjacent area is predominantly rocky reef and although the MOF footprint will completely and permanently cover this substrate in the subtidal zone, the material of the MOF will provide alternative hard surfaces and crevices for flora and fauna to colonise. As there are expected to be no direct loss or clearance of intertidal zone the small area of adjacent disturbance and provision of alternate substrates results in a negligible impact to intertidal habitat. Hydrodynamic modelling predicts limited changes to the oceanographic conditions at the port site as a result of the proposed infrastructure (Jacobs, 2014d). The existing site hydrodynamics confine the majority of longshore sediment drift to the two embayments either side of the proposed jetty and MOF structure.

The changes to the hydrodynamic conditions at the site (as discussed in Section 5.2) are generally associated with the construction of the MOF and pylon structure. As indicated by the hydrodynamics modelling (Jacobs, 2014d), alterations to the maximum currents and bed shear across the site are not significant. The predicted increase in sedimentation rate around the MOF is less than 1% (Jacobs, 2014d); and it is anticipated that the existing seagrass beds in the area will be able to adsorb this additional sediment. As such, alterations to the hydrodynamic conditions at the port site are not considered to affect intertidal habitats within the port site, representing a negligible impact.

Construction of the jetty includes impact piling and building the MOF outward from the headland will result in silt and sediment suspension into the water column. Increased turbidity can affect water quality and benthic intertidal communities as a result of decreased light penetration and silting in the intertidal zone. The existing wave climate at the port site regularly transports and resuspends sediment along the coast (Jacobs, 2014d). As such, localised turbidity during construction is considered to represent a low impact, with short term localised impact to intertidal flora during construction activity, but no longer term effects.

Tug vessels will be utilised for all large vessel movements in depths less than 23 m within the port site (including large vessel approaches and turning basins) to minimise propwash and subsequent sedimentation. Despite the utilisation of tugs, the repeated manoeuvring of large vessels will result in ship scour; destabilising sediments and resulting in short term turbidity. The increased turbidity will result in decreased light penetration to intertidal flora following large ship movements. Approximately 1 cargo ship movement per day is anticipated during operation of the proposed port. As such, localised impacts to intertidal flora within the port site as a result of sediment suspension will occur on a daily basis for the life of the port facility, and is considered to represent a medium impact.

Recovery of intertidal habitats is anticipated to occur slowly post-construction, with intertidal rocky shore flora and fauna colonising suitable areas of substrate on the MOF. Many species of macroalgae (and some encrusting invertebrates) have generation times rapid enough to recolonise disturbed areas once construction has ceased. However, recovery will be inhibited by changes in any factor which restricts the growth of macroalgae or encrusting fauna, including increased sediment transport, turbidity or pollution (i.e. contamination from anti-foulants).

6.2.2.2 Risks

The identification of IMS during construction and operation of the proposed port is considered likely despite the implementation of control measures. As Cape Hardy is isolated from a hydrodynamic perspective (refer Section 5.2) IMS that may colonise the area would not be readily transported by currents beyond the study area and into new regions. As such, most IMS would rely on vessel movements for transportation to areas beyond the study area. To minimise the likelihood of spreading IMS in Spencer Gulf waters, measures will be developed and implemented to prevent settlement of IMS, prevent the growth of settled IMS and / or the



removal and disposal of IMS if established. As such, the consequences of IMS are considered to be isolated to the Cape Hardy study area and are not considered likely to affect the marine environment at a regional level. IMS may result in the long term local decrease in abundance of marine species, considered to be a moderate consequence. As such, the overall risk associated with IMS is considered to be high.

Waste management and handling procedures developed as part of the CEMP and OEMP will control waste streams with the overarching aim of no waste products entering the marine environment. Despite the implementation of control measures, it is considered almost certain that some form of waste / debris will enter the marine environment during construction and operation of the port. As the port site is not a key habitat or breeding area, the consequences of debris entering the marine environment are considered to be minimal; insignificant to the overall viability of species utilising the intertidal habitat. As such, the overall risk is considered to be low.

Species in the intertidal habitat are also at risk from the unintended discharge of pollution in the form of site run-off, wastewater, hydrocarbons or chemicals. With any marine activity there is potential for waste or spills to enter the marine environment. The implementation of design measures (refer Section 4) for the control of surface and waste water at the port site will limit the majority of hazardous pollutants from entering the marine environment. There remains a risk that a major spill of oil or other chemicals may occur as a result of vessel failure or accident. Based on the experiences of other operating ports, the likelihood of this event occurring is considered to be rare. As previously outlined, the port site does not represent a critical intertidal habitat for protected species. The consequences of a major spill event are considered to be moderate, resulting in a long term decrease in the abundance of local marine flora and fauna. As such, the overall risk of a major spill event is considered to be low.

6.2.3 Conclusions

The intertidal communities at Cape Hardy are not considered unique with comparable habitats commonly observed in this region of the Spencer Gulf. The EPBC PMST, BDBSA search and desktop literature review did not identify any rare or protected species known to exclusively occur in the intertidal zones at Cape Hardy.

The total area of intertidal zone expected to be removed due to the construction of the MOF is less than 0.1 ha. As the majority of the intertidal zone within the study area will not be directly impacted from the development the 0.1 ha loss below the MOF and tug harbour is not considered significant. The MOF and tug harbour itself will provide additional rocky habitat in the intertidal zone for organisms to colonise.

After taking into consideration the mitigation measures proposed during the construction and operation phases of this development, the introduction of IMS remain a high risk to the intertidal habitat of Cape Hardy.

The majority of the intertidal zone is expected to be conserved and affected areas are expected to recover over time following construction. However the potential impacts from marine debris, chemical spills and IMS will have to be continually managed.



6.3 Subtidal Habitats and Flora

The subtidal marine environment of the Spencer Gulf is varied, supporting a wide range of habitats and species groups. Species in the lower reaches of the gulf are generally associated with three key habitat groups: seagrass meadows, rocky reefs and soft substrates. A clear understanding of which subtidal habitats and species are present at the port site is necessary to assess the biological importance of the site, identify any species that trigger further regulatory requirements, and any species or habitats requiring implementation of specific management or mitigation measures. Reefs and seagrass meadows provide essential habitat to a wide range of marine species such as fish, crabs and sea urchins; while mature seagrass meadows also stabilise the seabed to prevent erosion and sand movement.

Studies by Edyvane in the 1980s and 90s synthesised the available information relating to the marine biogeography and conservation values of Spencer Gulf (Edyvane, 1999a; b). A comprehensive study of the gulf benthos has not occurred since the late 1990's and the overall coverage of gulf benthos is based on calculations from Edyvane's work (DEWNR, 2011; Nature Maps, 2013). There is approximately 552,000 Ha of seagrass within the Spencer Gulf; representing approximately 60% of seagrass in South Australia, and 10% of all seagrass habitat in Australia (Edyvane, 1999a). Within the Gulf, the Cape Hardy study area falls within the 'Southwest Spencer Gulf' bioregion (between Cape Catastrophe and Tumby Bay), which itself supports an estimated seagrass coverage of 137,700 Ha (Edyvane, 1999a).

Typically, Spencer Gulf seagrass meadows are dominated by *Zostera* spp. and *Heterozostera nigricaulis*. A species shift to *Posidonia* spp. occurs in the few metres below the low tide mark, with *P. australis* dominating in shallower waters, while *P. sinuosa* and *P. angustifolia* become more dominant with increasing depth (Seddon, 2000). Other species such as *Amphibolis antarctica*, *A. griffithii*, and *H. nigricaulis*have been found to occupy edges, blowouts, and smaller areas within *Posidonia* spp. meadows (Edyvane, 1999a). The local distribution and abundances of these species are affected by numerous factors, including wave energy, tidal velocity, sediment stability, and light availability (Shepherd and Robertson, 1989).

Existing data for the study area identified the presence of sandy and rocky substrates, at depth ranging from the low water mark in the intertidal to 25 m at the outer extents. Moderate wave exposure, no mangrove or saltmarsh habitat, and no nearby estuaries were identified (Nature Maps, 2013 – National Benthic layer; DEWNR, 2011). Estimates of coverage from the National Benthic layer (Nature Maps, 2014) were based on aerial imagery with limited ground-truthing, and showed that sandy substrates were expected to be inhabited by seagrasses (predominantly *Posidonia* spp.) of varying density, between approximately 19 m of depth and the near-shore up to the intertidal zone (Figure 6-1). Given the physiological requirements *Posidonia* spp. for minimum water depth and light availability it was considered unlikely that dense seagrass would exist from the intertidal to out beyond 20 m.

6.3.1 Existing Environment and Key Findings

The desktop review suggested that the subtidal area of Cape Hardy is typical of southern Australia and in particular the Spencer Gulf. A review of the National Benthic habitat data indicated that the study area was predominantly covered by dense seagrass, though the rocky headlands were likely to provide rocky reef habitat (see Figure 6-1). This mapping was undertaken between the late 1990s and 2000s, using satellite imagery to detail any underwater features discernible on 1:100,000 scale prints and supported in places by a range of other data sources (Nature Maps, 2013). It is unknown whether ground-truthing of the satellite imagery was undertaken for the National Benthic habitat mapping of the study area, so Nature Maps was only used during the desktop portion of the study to inform the likely habitat types.

The only previously existing data which ground-truths against the state/national benthic habitat maps comes from benthic surveys carried out as part of two expired aquaculture lease applications in the area. These leases have since been surrendered (Section 6.9). The surveys undertaken for the lease application showed sparse



biota including small clumps of green and brown algae (Chlorophyta and Pheophyta), with no seagrass or benthic community identified (PIRSA, 2008). Review of the aerial imagery indicates little coverage of seagrass in the nearshore. These finding were in contradiction to the National Benthic Habitat maps (Figure 6-1; Nature Maps, 2013), which predicted dense seagrass, highlighting the need for ground-truthing of the study area, as undertaken using towed video.

An EPBC Protected Matters search was undertaken covering the study area and an additional buffer of 10 km. The PMST did not identify any Threatened Ecological Communities, National Heritage Places or EPBC-listed marine flora. However a number of protected fauna species commonly associated with seagrass beds were identified in the search, including seahorses and pipefish from the family Syngnathidae. These species are discussed in more detail in Section 6.6. The full EPBC PMST search results are presented in Appendix F.

JACOBS




Figure 6-1 State / National benthic habitat maps across the area

JACOBS

Three primary habitat types were recorded in the study area, including (see video stills in Figure 6-2):

- The 'Shallow Benthos' (<16 m depth) is largely inhabited by seagrass meadows (predominately *Posidonia* spp., with some *A. antarctica* patches restricted to shallower waters (<12 m depth)). Figure 6-2 images (A) and (B). The northern nearshore extent of the study area contained the greatest coverage of seagrass with densities over 50%
- The 'Mid-benthos' (depth 16 18 m) comprises predominantly bare fine sand and silty sediment with very sparse mixed small algae, very sparse *Posidonia* sp.(<5 % coverage) and occasional scattered invertebrates Figure 6-2 image (C)
- The 'Deep Benthos' (>20 m Depth), comprises bare silt with clumps of mixed invertebrates (sponges, ascidians and both motile and sessile crustaceans), Figure 6-2 images (D) and (E). Very sparse, mixed small algae were also present along with some evidence of cyanobacterial matting in deeper water Figure 6-2 image (F)
- No dense seagrass beds were observed beyond 19 m depth and macroalgae became uncommon after 21 m depth (Figure 6-3)




Figure 6-2 Study area benthic habitat types

GIS analysis of the towed video survey data generated a benthic habitat map for the study area (Figure 6-3). The map shows that benthic flora followed depth and substrate contours, with a higher density of flora coverage in water less than 16 m deep, where light is generally limited (dependent on water clarity) and the sediment was found to be coarser.



The benthic flora throughout the study area when present was intact with no signs of scouring or removal. No IMS were recorded during the video survey. There were large areas of discontinuous vegetation, with isolated patches of benthic vegetation generally found in waters deeper than 10 m. Within the study area the denser coverage (>50%) of seagrass or macroalgae was confined to the northern areas and shallow waters around the headlands south of the proposed marine infrastructure. These areas of denser seagrass coverage in particular provide habitat opportunities for fauna, and may provide habitat for conservation significant species that occur in the area.

The video survey also observed scattered small patches of rocky reef habitat in shallower waters. The near-shore areas were not surveyed due to the draft clearance requirements of the survey vessel, and safety constraints related to towing the camera over shallow reef. It is expected from analysis of aerial imagery that more complex rocky reef habitat exists along the shoreline of the cape, including the intertidal substrates discussed in Section 6.2. The patches of rocky reef and nearshore rocky substrates are considered to be related to boulder fields or outcroppings of rock that were identified during the nearshore geophysical surveys (MES, 2012).






Figure 6-3 Benthic habitat map of study area developed from towed video



A review of the publicly available literature for the wider area (Edyvane, 1999a; PIRSA, 2008; Golders Associates, 2012; BDBSA, 2013) and benthic species noted during the video tow were used to compile a table of likely habitat-forming marine flora (Table 6-2).

Table 6-2 Habitat-forming marine flora likely to occur at Cape Hardy

| | Scientific Name | Common Name | Legislative Protection* |
|---|-----------------------------------|--------------------|-------------------------|
| Seagrasses | Amphibolis antarctica | Wire weed | NV Act |
| Division: Magnoliophyta | Amphibolis griffithii | Wire weed | NV Act |
| Division: Magnonophyta | Halophila spp. | Paddleweed | NV Act |
| | Heterozostera nigricaulis | Eelgrass | NV Act |
| | Posidonia spp. | Tapeweed | NV Act |
| | Zostera spp. | Garweed | NV Act |
| | Zostera muelleri subsp. mucronata | Garweed | Rare (NPW), NV Act |
| Brown algae | Caulocystis spp. | | NV Act |
| Division: | <u>Corynophlaea cystophorae</u> | | NV Act |
| Heterokontophyta Class: Phaeophyceae | <u>Cystophora expansa</u> | | NV Act |
| Рпиеорпусеие | <u>Cystophora moniliformis</u> | | NV Act |
| | Cystophora polycystidea | | NV Act |
| | <u>Cystophora subfarcinata</u> | | NV Act |
| | Cystophora spp. | | NV Act |
| | Cystoseira spp. | | NV Act |
| | Ecklonia radiata | Common kelp | NV Act |
| | Hincksia spp. | Snot algae | NV Act |
| | Hormosira banksii | Neptune's necklace | NV Act |
| | <u>Pachydictyon paniculatum</u> | | NV Act |
| | Ralfsia verrucosa | | |
| | Sargassum decipiens | | NV Act |
| | <u>Sargassum paradoxum</u> | | NV Act |
| | Sargassum spp. | | NV Act |
| | Scaberia agardhii | | NV Act |
| | Zonaria spiralis | | NV Act |
| Red Algae Division: Rhodophyta | <u>Capreolia implexa</u> | Red turf alga | NV Act |
| | <u>Rhabdonia clavigera</u> | | NV Act |
| Green Algae Division: Chlorophyta | <u>Cladophora feredayi</u> | | NV Act |
| Division. Chlorophyta | <u>Cladophora lehmanniana</u> | | NV Act |

^{*}The Native Vegetation Act regulates actions which may involve 'a plant or plants of a species indigenous to South Australia including a plant or plants growing in or under waters of the sea...'. Whilst algae are not strictly plants, the Native Vegetation Council often includes terrestrial mosses and lichens (which are not plants) as vegetation. List of species likely to occur were developed from literature searches (not underlined) and survey records. **Bold** entries represent species or groups observed during habitat surveys at the site. Underlined entries are taken from the BDBSA search records.



6.3.1.1 Ecological Significance of the Subtidal Habitat Present in the Study Area

The benthic habitat survey of the study area detailed the distribution and range of habitat present at the site. The majority of the inshore (<16 m depth) areas of the site are covered by varying densities of seagrass, which is of high conservation value. Deeper areas are generally not vegetated and involve scattered filter feeders and other invertebrates. While the rocky reef areas within the study area are small, they may still provide niche or refuge areas from which species can forage in the wider area. The findings are consistent with current knowledge about the ecological significance of subtidal habitats within the wider Spencer Gulf.

Seagrass beds (along with all marine benthic flora) are a protected habitat in South Australian waters under the Native Vegetation Act, and are considered a habitat of high conservation value. Seagrass beds are associated with a host of secondary benefits including playing key roles in carbon and nutrient cycling (Duarte and Cebrian, 1996), sediment stabilisation (de Boar, 2007), supporting biodiversity (Edgar *et al.*, 1994) and providing nursery habitats for many commercially fished species (Beck *et al.*, 2001; Jackson *et al.*, 2001, Heck *et al.*, 2003). A wide range of South Australian fish and crustacean species of commercial and recreational fishery importance rely on seagrass habitats, including anchovy (*Engraulis australis*), snapper (*Pagrus auratus*), western king prawn (*Melicertus latisulcatus*) and blue swimmer crab (*Portunus armatus*) (Edyvane, 1999a; b; Bryars, 2003).

Rocky reefs support a diverse food web based on filter feeders and mobile species including commercial fish and crustaceans that use the reef habitat as a refuge to forage from. The weathering pattern of the underlying rock can support a high complexity of habitat types and ecological niches that can be available for species inhabiting the area.

6.3.2 Impacts and Risks: Subtidal habitats

Impacts and risks to subtidal habitats associated with the construction and operation of the proposed port facility involve immediate short-term impacts during construction such as clearance of habitat, and long-term processes during operations, such as hydrodynamic changes, smothering, long-term shading, ship scour, anchor damage, changes in water quality from run-off or stormwater, spills or the introduction of IMS.

Impacts to subtidal habitats associated with the proposed port facility could occur as a result of:

- Habitat loss or exclusion
- Erosion (from stormwater and vessel wakes/propwash, changes to hydrodynamics)
- Spills (e.g. oil), pesticides, herbicides, or other pollutants
- A build-up of contaminants (e.g. anti-foulants)
- Marine debris
- The introduction of IMS
- Vessel run aground or sinking
- Shading effects from vessels, structures and increases in turbidity or spills

6.3.2.1 Impacts

There will be minimal direct loss of seagrass arising from the proposed jetty/wharf structure as seagrass was identified at depths greater than 10 m within the infrastructure footprint. The major disturbance in the proposed jetty/wharf footprint will occur from placement of the jack-up pile drivers, the laying of anchors and chain to hold the barge on location, and from driving piles into the seabed. While these construction activities will occur in depths >10 m, they are not expected to cause permanent loss of seagrass.

A conservative estimate of the area of each benthic habitat type impacted by the proposed port footprint has been calculated, referred to as the construction footprint. The footprint has been estimated based on conservative estimates of total clearance incorporating the areas of jetty/wharfpiling, MOF facility and tug



harbour, the vessel mooring areas, turning basins and anchorage area. All clearance footprints and disturbance areas have been conservatively calculated with an additional 5 m buffer. These estimates are presented in Table 6-3.

Table 6-3 Esimated area of subtidal substrate affected by marine infrastructure in the study area

| | Areas Likely to | b be Disturbed | Areas to be cleared | | |
|---|-----------------|----------------|---------------------|---------|--|
| Substrate Type | Anchorage area | Port Area | MOF & Tug Harbour | Jetty | |
| | (299.9 Ha) | (147.1 Ha) | (2.6 Ha) | (14 Ha) | |
| Bare sediment | 58.9 Ha | 131.6 Ha | 0 Ha | 6.49 Ha | |
| Seagrass habitat | 0 Ha | 14.7 Ha | 2.49 Ha | 0.16 Ha | |
| Macroalgal habitat | 241.2 Ha | 2.98 Ha | 0.51 Ha | 2.1 Ha | |
| Total Subtidal Habitat Impacted (vegetated áreas) | 258. | 7 Ha | 5.26 Ha | | |

Note: All disturbed or cleared areas include a 5 m buffer. Total Subtidal Habitat does not include bare sediment substrates.

The total area of seagrass to be cleared is 2.65 Ha, based on the footprint of the entire MOF, tug harbour and jetty area including vessel loading areas (with an additional 5 m buffer around areas for a conservative estimate). This calculation includes areas where seagrass will be only partly impacted by shading from fixed infrastructure and vessels. Only the area immediately beneath the MOF will suffer permanent loss, with the habitat beneath the jetty considered unlikely to be lost as a direct result of shading due to the jetty alignment. Allowing for the turning basins and anchorage areas associated with the port the total area of impacted seagrass would be 14.7 Ha (Table 6-3). Based on Edyvanne (1999a) the estimated area of impact represents <0.003% of the total known seagrass from within Spencer Gulf.

Calculation for the combined intertidal and near-shore subtidal environment around the headlands in the study area to be cleared have been based on the geophysical surveys (MES, 2012) and aerial imagery. A conservative (i.e. biased toward over-estimation) estimate of 48 Ha of intertidal rocky reef exists in the study area. The combined intertidal and near-shore subtidal environment is dominated by rocky substrates, to a depth of approximately 8 m. Some areas of subtidal rocky reef will be subject to direct impact from the MOF and piling activity with permanent loss of macroalgal habitat (0.51 Ha from Table 6-3). There are expected to be no loss of intertidal rocky reef habitat.

The the majority of the roughly 260 Ha of impacted subtidal habitat are in areas of low density macroalgal or sparse seagrass (14.7 Ha) which will only incur intermittent impacts associated with vessel manouvering or shading. The vast majority of the vegetative habitat within the Cape Hardy study area (2736 Ha) has been avoided.

It is expected that recovery will occur post-construction, with Posidonia spp. seagrass slowly re-colonising suitable areas and Amphibolis spp. colonising suitable areas within 3 years. Recovery will be inhibited by changes in any factor which restricts the growth of seagrass (or macro-algae), including increased sediment transport, scouring or changes in water quality. Macroalgae have generation times rapid enough to recolonise once construction has ceased provided water quality and sediment conditions remain suitable. The clearance of seagrass and sub-tidal habitat as a result of the construction of the proposed port is considered a short term impact (recoverable within 3 years) and restricted to the study area. As such, the overall impact is considered to be low.



While seagrasses are considered to be susceptible to the effects of long-term heavy shading from infrastructure or increased turbidity as a result of ship movements, most species of seagrass, and in particular *A. antarctica* and *Posidonia spp.* which are likely to exist at the site, have some tolerance for partial shading and can survive beneath jetty structures (Duarte, 1991; Dennison *et al.*, 1993; Gordon *et al.*, 1994; Fitzpatrick and Kirkman, 1995; Masini *et al.*, 1995; Bryars and Collins, 2008; Bryars and Rowling, 2009). Similarly, macroalgae are capable of tolerating low-light conditions, with major habitat-forming species such as *E. radiata* occurring to a maximum depth of greater than 40 m (Edgar, 2001).

A variety of factors determine the area of seabed shaded including the dimensions of the jetty / wharf, the duration of vessel mooring and the solar angle. The jetty orientation runs at 331° north-north-west resulting in minimal shading by the jetty due to the approximate east-west sun path. Affected seabeds will only receive reduced light for around 1-2 hours per day. The sun is at its highest during summer at around 79° with the lowest angle in winter approximately 32°. Seasonal changes in the angle of the sun along with the orientation of the jetty structure means that there will be no permanent shading of the seabed beneath the jetty structure.

Shading from impact piling and temporary construction activities (i.e. jack-up barge and support vessels) is not considered likely to affect benthic habitats as the construction activities are only temporary and seagrass can tolerate heavy shading for several weeks due to their ability to draw energy from their root mass (Duarte, 1991; Westphalen *et al.*, 2004).

Shading effects at the port site are expected to be restricted to the areas adjacent to and beneath the jetty and adjacent the MOF. Seagrasses identified in the footprint area include sparse coverage of the tapeweeds *Posidonia* spp. seagrasses. As a result of the jetty orientation over an area with minimal seagrass coverage, and the lack of permanent shading indicates that there will be little to no shading impacts from the jetty on seagrass. As such, shading impacts to subtidal habitats are considered to be negligible.

Hydrodynamic modelling predicts limited changes to the oceanographic conditions at the port site as a result of the proposed infrastructure (Jacobs, 2014d). The existing site hydrodynamics confine the majority of longshore sediment drift to the two embayments either side of the proposed jetty and MOF structure.

The changes to the hydrodynamic conditions at the site (as discussed in Section 5.2) are generally associated with the construction of the MOF and jetty structure. As indicated by the hydrodynamics modelling (Jacobs, 2014d), alterations to the maximum currents and bed shear across the site are not significant. The predicted increase in sedimentation rate around the MOF is less than 1% (Jacobs, 2014d); and it is anticipated that the existing seagrass beds in the area will be able to adsorb this additional sediment. As such, alterations to the hydrodynamic conditions at the port site are not considered to affect intertidal habitats within the port site, representing a negligible impact.

Tugs will be utilised for all large vessel movements in depths less than 23 m within the port site (including large vessel approaches and turning basins) to minimise propwash and subsequent sedimentation. Despite the utilisation of tugs, the repeated manoeuvring of large vessels will result in ship scour; destabilising sediments and resulting in short term turbidity. The increased turbidity will result in decreased light penetration to subtidal habitats following large ship movements. Approximately 1 cargo ship movement per day is anticipated during operation of the proposed port. As such, localised impacts to subtidal habitats within the port site as a result of sediment suspension will occur on a daily basis for the life of the port facility, and is considered to represent a medium impact.



6.3.2.2 Risks

The identification of IMS during construction and operation of the proposed port is considered likely despite the implementation of control measures. As Cape Hardy is isolated from a hydrodynamic perspective (refer Section 5.2) IMS that may colonise the area would not be readily transported by currents beyond the study area and into new regions. As such, IMS would rely on vessel movements for transportation to areas beyond the study area. To minimise the likelihood of spreading IMS in Spencer Gulf waters, measures will be developed and implemented to prevent settlement of IMS, prevent the growth of settled IMS and / or the removal and disposal of IMS if established. As such, the consequences of IMS are considered to be isolated to the Cape Hardy study area and are not considered likely to affect the marine environment at a regional level. Despite the control measures proposed to be implemented, IMS may result in the long term decrease in abundance of seagrass and marine species in the study area, considered to be a moderate consequence. As such, the overall risk associated with IMS to local flora and fauna species is considered to be high.

Construction and operation of the proposed port will result in the generation of a number of waste streams that may enter the marine environment and potentially impact sub-tidal habitats. Waste management and handling procedures developed as part of the CEMP and OEMP will control waste streams with the overarching aim of no waste products entering the marine environment. Despite the implementation of control measures, it is considered almost certain that some form of waste / debris will enter the marine environment during construction and operation of the port. As the port site is not a key habitat and does not support large areas of seagrass, the consequences of debris entering the marine environment to sub-tidal habitats are considered to be minimal; insignificant to the overall viability of species. As such, the overall risk is considered to be low.

Seagrass is also at risk from the unintended discharge of pollution in the form of site run-off, wastewater, hydrocarbons or chemicals. With any marine activity there is potential for waste or spills to enter the marine environment and affect plankton, benthic fauna and habitat forming benthic flora (e.g. seagrass), reducing habitat suitability for fish and marine fauna.

Iron ore is not known to be toxic to seagrass or macroalgal assemblages but otherrun-off or pollution from the port (as discussed in Section 5.4) has the potential to impact on water quality either stimulating algal growth through increase nutrient loading, reducing salinity as a result of increased freshwater discharge during storms, or inputs of hydrocarbons and/or other pollutants. Nutrients from run-off can stimulate algae in the water column or epiphytes that would shade benthic flora while other changes in water quality i.e. chemical contaminants such as biocides in anti-foulant paint or hydrocarbons could have direct toxic effects on benthic flora. Excess nutrients can also encourage more phytoplankton to grow in the water, reducing the amount of light getting to the seagrass, particularly in the deeper regions. Seagrass is commonly lost when nutrient levels in the water increase, these nutrients cause a large number of epiphytes to grow on the seagrass leaves, blocking light or causing the leaves to become too heavy and to break off. The implementation of design measures (refer Section 4) for the control of surface and waste water at the port site will limit hazardous pollutants from entering the marine environment in all but extreme weather events.

There remains a risk that a major spill of oil or other chemicals may occur as a result of vessel failure or accident. Based on the experiences of other operating ports, the likelihood of this event occurring is considered to be unlikely. As previously outlined, the port site does not represent a critical habitat or support large areas of seagrass. The consequences of a major spill event are considered to be minor, as seagrass within the study area is representative of the broader Spencer Gulf. As such, seagrass loss would result in a short term decrease in the local abundance. As such, the overall risk of a major spill event affecting subtidal habitats and flora is considered to be low.



6.3.3 Conclusions

The subtidal habitats and flora identified at Cape Hardy during habitat mapping are consistent with the habitats observed in this region of the Spencer Gulf. Protected seagrass beds, which are likely to support a number of associated protected species, dominate the shallow waters to 10 m depth. No seagrass beds were present in depths greater than 19 m. Flora coverage is denser (greater than 50%) in the northern extents of the study area, whilst some areas along the shoreline to the south have a sparse cover of 5-15%. The presence and density of seagrass beds have been considered in the siting and alignment of marine infrastructure elements to avoid and minimise impacts. The total area of seagrass within the study area is around 930 Ha, of which less than 2.65 Ha (of sparse seagrass) is expected to be cleared by the port development.

6.4 Benthic Fauna

Benthic fauna can be used as an indicator of an area's environmental condition (Llansó, 2002). Most infauna (benthic fauna living within the sediments) are sedentary and respond rapidly to local environmental impacts and disturbances. Similarly, the majority of infauna and epifaunal (benthic fauna attached to the surface of substrates, rocks, vegetation or other benthic fauna) cannot easily move to avoid stressful conditions. Infauna often have short generation times and are important components of aquatic food webs, affecting transport and cycling of nutrients and toxic substances. As such, the abundance and diversity of benthic infauna can be used to indicate habitat health (Llansó, 2002; McConnaughey and Syrjala, 2008).

In general, a high density of diverse infaunal invertebrate species is an indicator of good habitat health. Conversely, low densities or a low diversity of infaunal invertebrate species indicates a lower value habitat, or one which is potentially already impacted (Ward and Hutchings, 1996). A complex and high density of epifauna can be an indicator of good habitat health, provided the species present are endemic species, while many degraded habitats are often dominated by high numbers of only a few taxa. *Zostera* and *Posidonia* beds have been shown to support distinctive epifauna and infauna assemblages (Hutchings *et al.*, 1993) therefore benthic fauna can also act as an indicator of habitat composition as well as system health.

6.4.1 Existing Environment and Key Findings

The physical isolation of the Spencer Gulf along with a lack of oceanic upwelling and low volumes of freshwater inputs has led to increased salinities and resource scarcity within the gulf. These physical conditions have contributed to high levels of endemism within the upper gulf; however the lower gulf has more in common with the southern Australian region (Poore, 1995).

Previous studies of the benthic invertebrate and infauna assemblages within the Spencer Gulf have focused on the northern and upper gulf regions (Shepherd, 1983; Ainslie *et al.*, 1989; Hutchings *et al.*, 1993; Ward and Hutchings, 1996; Gillanders *et al.*, 2013). Hutchings *et al.*, (1993) investigated the infaunal community of marine sediments and seagrass beds in the upper Spencer Gulf (near Port Pirie) and showed polychaetes to be the dominant invertebrate taxa, comprising up to 76% of the infauna species in *Posidonia* seagrass beds and 48% of the infauna species in *Zostera* beds. Other infauna taxa found within Spencer Gulf include crustaceans, molluscs, ascidians and echinoderms.

The distribution of polychaete, mollusc and crustacean infauna in the upper Spencer Gulf has been connected to concentrations of trace metals in sediments (As, Cu, Mn, Pb and Zn), with areas dominated by polychaetes correlated with areas of higher trace metals (Ward and Hutchings, 1996). These studies were generally carried out in areas of high anthropogenic disturbance. Areas with a high level of disturbance can also change sediment composition i.e. distribution of grainsize can lead to differences in taxa between sites with delicate bivalves such as *Tellina* spp. preferring undisturbed coarse sand and opportunistic taxa such as polychaetes colonising areas of high disturbance in either silt or gravel.



The results of the benthic infaunal analysis are presented in Table 6-4. In summary the results indicate:

- The shallowest transect (A) had the greatest infaunal diversity, as well as the highest number of organisms per sample (Table 6-4). This transect was aligned with dense areas of seagrass habit (Figure 6-3) and sandy sediments
- Transect B had the next highest diversity of infauna, this transect had some areas of sparse to medium density seagrass habitat.
- Transect C that had predominantly sparse or no seagrass habitat with large areas of cyanobacterial mat and had the lowest diversity of infauna.
- Transect D was on average the deepest transect and was dominated by mixed small macroalgae, cyanobacterial mats and groupings of sponge and ascidians, its diversity of infauna was only slightly lower than Transect B.

Moving from north to south within the site; the trends of highest infauna diversity again aligned with areas of densest seagrass coverage at the north of the study area (0.1 infauna/cm $^3 \pm 0.002$; density \pm st. dev) and the lowest diversity of infauna was recorded at the southern extent of the study area (0.03 infauna/cm $^3 \pm 0.001$; density \pm st. dev.). The numbers of infauna recorded during survey is comparable to other recent studies of undeveloped sites in Spencer Gulf (Golder Associates, 2012; Sinclair Knight Merz, 2011) and the diversity of species indicates a good overall habitat health for the area.



Table 6-4 Richness and density of macroinvertebrate infauna collected from the marine sediment sampling locations within the Cape Hardy study area

| Family | | Location | | |
|---|--------------------------|-----------------------------|-----------------------------|------------------------------|
| Ī | Transect A (~10 m depth) | Transect B (~15 m depth) | Transect C (~20 m depth) | Transect D (> 22 m depth) |
| Nephtyidae | 2 | | | 1 |
| Syllidae | 2 | | | |
| Cheirocratidae | 3 | 1 | | |
| Amphipoda | 1 | | | |
| Eusiridae | 2 | | | |
| Anaspidacea | 3 | | | |
| Asellota | 2 | | | |
| Isopoda | 4 | | | |
| Kalliapseudidae | 1 | | | |
| Chamidae | 1 | | | 1 |
| Limidae | 1 | | | |
| Littorinidae | 2 | | | |
| Tauraxinus sp. | 1 | | | 1 |
| Veneridae | 1 | | | |
| Patellidae | 1 | | | |
| Armandia sp. | | 1 | | |
| Lysianassidae | | 1 | | |
| Leptostraca | | 2 | | |
| Asteroidea | | 1 | | |
| Nuculana sp. | | 3 | | 1 |
| Patellidae | | 1 | | |
| Astralium sp. | | 1 | | |
| Nemertea | | 1 | | |
| Actiniaria | | | 1 | |
| Echinoidea | | | 1 | |
| Tellina sp. | | | 2 | 2 |
| Terebellidae | | | | 1 |
| Stegocephalidae | | | | 2 |
| Volutidae | | | | 1 |
| Total # infauna spp. | 27 | 14 | 4 | 10 |
| Density of infauna/cm ³ ± st. dev. | 0.11 ± 0.002 | 0.06 ± 0.002 | 0.02 ± 0.00 | 0.04 ± 0.000 |



Analysis of the towed video footage identified a number of epifaunal invertebrates in the study area. These are listed in Table 6-5 below. None of these species are protected under the EPBC or NPW Acts; several of the species are afforded protection in specific (aquatic reserve) areas or seasons under the SA Fisheries Management Act 2007.

Table 6-5 Benthic fauna of interest in the study area

| | Scientific Name | Common Name | Legislative Protection | Commercial/ Recreational Interest |
|-------------------------------|--|--|--|---|
| Annelida - Polychaeta | Multiple unidentified spp. | Unidentified bristle worms, sandworms | Nil | No |
| Porifera - | Multiple unidentified spp. | Unidentified sponges. sea sponge, barrel, branching, chimney, ball | Nil | Minimal |
| Urochordata - Ascidians | Multiple unidentified spp. | Sea squirt | Nil | No |
| Echinodermata - Asteroidea | Coscinatsterias muricata | Eleven-arm seastar | Nil | No |
| | Unknown spp. | 2 spp. unidentified seastars | Nil | No |
| - Echinozoa | Multiple unidentified spp. | Unidentified sea urchins | Nil | Minimal |
| - Holothuroidea | Stichopus ludwigi/ Holothuria hartmeyeri | Sea cucumbers | Nil | Not in SA |
| Mollusca | Haliotis rubra/ laevigata | Abalone (Roe's, Black-lip and Green-lip Abalone) | At all times in intertidal reefs and aquatic reserves* | Yes |
| | Pinna bicolor | Razorfish | Catch limited | Yes |
| Anthropoda | Ibacus alticrenatus, Ibacus peronii | Slipper lobster/bug | When carrying external eggs* | Yes |
| | Jasus edwardsii | Southern rock lobster | Protected from collection May-Nov | Yes |
| | Majoidea (various) | Spider crab | Nil | Minimal |
| | Ovalipes australiensis | Sand crab | When carrying external eggs* | Yes |
| | Portunus armatus | Blue swimmer crab | When carrying external eggs* | Yes |
| | Pseudocarcinus gigas | Giant crab | When carrying external eggs* | Yes |

^{*}Fisheries Management Act, 2007: (PIRSA, 2014). **Bold** entries represent opportunistic sightings during habitat surveys at the site.



6.4.2 Risks and Impact: Benthic fauna

Impacts could occur to the benthic fauna as a result of:

- Habitat loss or exclusion
- Build-up of contaminants, including hydrocarbons and biocides (including antifoulants, herbicides and pesticides)
- Alterations to hydrodynamic or sediment transport conditions
- Removal of habitat-forming species (i.e. seagrass)
- Changes to sediment chemistry
- Marine debris

6.4.2.1 Impacts

Disturbance to benthic fauna is expected to occur in the immediate areas involved in piling for the jetty and under the rockwall related to the MOF (see Section 6.3.2). Additional areas that may be subject to disturbance include areas underneath the jetty, ship berths and vessel manoeuvring areas, which will be subject to occasional shading. It is expected that there will be complete direct disturbance to the benthic habitat within the footprint of port structures and minor disturbance in vessel manoeuvring areas. Conservatively, 2.65 Ha of seagrass and 2.61 Ha of combined intertidal and near-shore subtidal environment will be cleared (refer Table 6-3).

Soft sediment areas directly under the MOF will be permanently lost. However, epifauna from hard substrate areas will be able to colonise the MOF and jetty surfaces, thereby creating new areas of habitat which will further increase habitat diversity as additional organisms colonise the soft/biological surfaces of the encrusting organisms.

Typically, benthic fauna will regenerate following significant disturbance such as that required to support the proposed port development. As such, impacts to benthic fauna within the study area associated with habitat clearance will be resolved in the short term (< 3 years), and are considered to represent a low impact.

Hydrodynamic modelling predicts limited changes to the oceanographic conditions at the port site as a result of the proposed infrastructure (Jacobs, 2014d). The existing site hydrodynamics confine the majority of longshore sediment drift to the two embayments either side of the proposed jetty and MOF structure. The changes to the hydrodynamic conditions at the site (as discussed in Section 5.2) are generally associated with the construction of the MOF and jetty structure. As indicated by the hydrodynamics modelling (Jacobs, 2014d), alterations to the maximum currents and bed shear across the site are not significant. The predicted increase in sedimentation rate around the MOF is less than 1%, and it is anticipated that the existing seagrass beds in the area will be able to adsorb this additional sediment. As such, alterations to the hydrodynamic conditions at the port site are not considered to significantly affect the habitat, density or diversity of benthic fauna, representing a negligible impact.

An assessment of the likelihood of occurrence and impacts to each invertebrate species potentially occurring within the study area is provided in Table 6-6 below.

Table 6-6 Invertebrate species: likelihood of occurrence and potential impacts

| | nvertebrate Species otentially in the are | | Likelihood of Occurrence | | Likelihood of significant impacts | |
|------------------------------------|--|---|--|--|---|--------------------------------|
| Species Name | Common Name | Legislative Protection. EPBC, NPW or Fisheries Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| Jasus edwardsii | Southern rock lobster | Fisheries Act (SA) ICUN least concern | Possible for species to be present in area. This species is of commercial and recreational interest and occurs throughout the Spencer Gulf. To occur in high densities this species requires rocky reef habitat with a high degree of complexity (holes, cracks, overhangs or ledges). The rocky reef at Cape Hardy is granitic, which generally provides less complexity and accordingly a lower density of rock lobster (Sloan and Crosthwaite, 2007). Therefore, while it is possible that southern rock lobsters are present at Cape Hardy, the immediate area is highly unlikely to support a significant population. | Disturbance to habitat; pollution; oil spills; fisheries access/exclusion. | The species are protected from collection May-November under the Fisheries Act (SA) but are not protected under any other legislation. There will be a small area of intertidal and shallow sub-tidal reef impacted by the wharf construction. While this habitat may support some juvenile rock lobster, it is unlikely that Cape Hardy represents a significant site of lobster recruitment or habitat. It is also unlikely that the site represents a significant lobster fishing site. No significant impact to this species will occur as a result of this development. | N |
| Haliotis rubra, laevigata, roei | Abalone (Black- lip, Green-lip and Roe's abalone) | Fisheries Act (SA) | Possible for species to be present in area. This species is not listed for protection but is of commercial and recreational interest, and the collection of this species is under tight control. Abalone occur in rocky reef habitat throughout the Spencer Gulf. It is likely that suitable habitat for abalone occurs in the intertidal and shallow sub-tidal strip along the shore-line at the study area. | Disturbance to habitat; pollution; oil spills; IMS; fisheries access/exclusion | The species are protected from collection on intertidal reefs at all times under the Fisheries Act (SA). There will be a small area of intertidal and shallow sub-tidal reef impacted by the wharf construction. While this habitat may support some abalone, it is unlikely that the development will cause a permanent disturbance, or that Cape Hardy represents a significant site of abalone recruitment or habitat. It is also unlikely that the site represents a significant Abalone fishing site. No significant impact to this species will occur as a result of this development. | N |
| Portunus armatus | Blue swimmer crab | Fisheries Act (SA) | Present. A number of individual crabs were sighted during the video tow survey in range of substrates from bare sand to dense seagrass. Spencer Gulf Blue swimmer crabs are believed to be an isolated sub-population and rely on widespread mangrove creek and mud-flat habitats during recruitment stages. The crabs are found on a range of habitat types from algal or seagrass substrate through to bare sand or muddy substrate. Adult crabs generally migrate to deeper waters up to 50 m during the months of May to August but migrate to shallower waters in warmer months (Dixon et al., 2013). This species occurs at the site but it is unlikely that the | Disturbance to habitat; pollution; oil spills; fisheries access/exclusion. | Females of this species are not to be taken when carrying eggs as they are protected under the Fisheries Act (SA). The species is not listed under any other protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf, across seagrass habitat and muddy or sandy substrates There will be a small area of suitable habitat for this species impacted by the wharf construction, but it is unlikely to have an effect on the species' ability to inhabit the area. The development site does not represent habitat critical for recruitment or growth of the wider population of this species. It is also | N |

| | nvertebrate Species | | Likelihood of Occurrence | | Likelihood of significant impacts | | |
|------------------------------------|-------------------------------------|---|---|--|---|--------------------------------|--|
| Species Name | Common Name | Legislative Protection. EPBC, NPW or Fisheries Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | density of individuals is high. | | unlikely that the site represents a significant Blue swimmer fishing site. No significant impact to this species will occur as a result of this development. | | |
| Ovalipes australiensis | Sand crab | Fisheries Act (SA) | Present. This species was sighted during the video survey on areas of bare sand throughout the study area. It is of commercial and recreational interest and occurs throughout the Spencer Gulf. Sand crabs generally inhabit sandy substrates and seagrass beds. It is likely that these crabs exist in the seagrass areas as well at Cape Hardy, due to the species being known to use seagrass habitats. But it is unlikely that the site supports a large population of these crabs. | Disturbance to habitat; pollution; oil spills. | Females of this species are protected under the Fisheries Act (SA) when carrying external eggs. The species is not listed under any other protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf, across seagrass habitat and muddy or sandy substrates There will be a small area of suitable habitat for this species impacted by the wharf development, but it is unlikely to have a lasting impact on the species' ability to inhabit the area. The development site does not represent habitat critical for the recruitment or survival of this species. No significant impact to this species will occur as a result of this development. | N | |
| FAMILY MAJIDAE | Spider crab | N/A | Present. Spider Crabs were observed on sandy or muddy bare substrate in water greater than 15m depth (generally >20 m) throughout the study area. Due to the habitat type and depth range and size of the crabs observed the species were likely <i>Leptomithrax gairmardii</i> | Disturbance to habitat; pollution; oil spills. | None of the species of Spider Crabs found in South Australia are protected or considered a commercial fisheries and as <i>Leptomithrax gairmardii</i> are generally found in waters deeper than 15m the development will impact on only a small area of suitable habitat and it is unlikely to have a lasting impact on the Spider Crabs' ability to inhabit the area. The development site does not represent habitat critical for the recruitment or survival of this species. No significant impact to this species will occur as a result of this development. | | |
| Ibacus peronii, I. alticrenatus | Slipper lobsters or Slipper bugs | Fisheries Act (SA) ICUN least concern | Present. This species is of commercial and recreational interest and occurs throughout the Spencer Gulf. Slipper lobsters are found in a wide range of habitats including both sandy and muddy substrates. Slipper lobsters are generally taken incidentally as a part of other commercial and recreational fishing efforts (such as fishing for other crustaceans, including prawn trawling). Suitable habitat for this species exists in the area, and one was sighted on the transect video from site surveys. | Disturbance to habitat; pollution; oil spills. | There will be a small area of suitable habitat for this species impacted by the wharf development, but it is unlikely to have a lasting impact on the species' ability to inhabit the area. It is unlikely that the study area supports a high density of slipper lobsters or is a critical area for this species. It is also unlikely that the site represents a significant lobster fishing site. No significant impact to this species will occur as a result of this development. | N | |

| | nvertebrate Species | | Likelihood of Occurrence | | While there may be suitable habitat for adult prawns in the N | | |
|---|--------------------------------|--|--|---|--|--------------------------------|--|
| Species Name | Common Name | Legislative Protection. EPBC, NPW or Fisheries Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Penaeus (Melicertus) latisulcatus | Western king prawn | N/A | Possible that species is present in area. This species is not listed for protection but is of commercial interest and occurs throughout the Spencer Gulf. Coastal mangrove and tidal flat/saltmarsh habitat is critical to population recruitment and occurs throughout Spencer Gulf, especially north of Cowell. None of this recruitment habitat occurs in the study area. Deeper waters with a generally sandy or silty substrate are the preferred habitat for adult prawns, rather than specifically seagrass or reef. It is likely that there is some suitable habitat in the outer extents of the study area. The key areas that prawns inhabit have, however, been identified and are unlikely to include the study area. For example, recently around 50% of prawns have been harvested from the Wallaroo region on the eastern side of Spencer Gulf (Dixon and Sloan, 2007). | Disturbance to habitat; pollution; oil spills. | deeper areas of the study area it is unlikely that the development or shipping activity will affect this habitat. It is also unlikely that the site represents a significant prawn trawling site. No significant impact on this species or | N | |
| Sepioteuthis australis | Southern calamary | N/A | Possible that species is present in area. This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. Southern calamary typically cross habitat boundaries, with larger adults preferring deeper waters while smaller individuals are typically found in shallow waters, not necessarily attracted to seagrass or rocky reef (Dixon and Sloan, 2007). As these habitats exist in the area it is possible the species will be present. | Disturbance to habitat; pollution; oil spills. | It is possible that this species occurs in the study area, but the area does not represent a large area of critical habitat for this species. The development and/or shipping operations are not expected to have any significant impact on this species or its habitat. | N | |
| Sepia apama | Giant Australian cuttlefish | Reef Watch species of SA conservation concern. ICUN Near Threatened EPBC not listed, Fisheries Act (SA) | Present. A single cuttlefish was sighted during the video tow in an area of bare silt substrate at over 20 m. This species is of conservation concern due to its unique behaviour of aggregating annually to breed in a small area near Whyalla (upper Spencer Gulf) but is only protected in the Northern Spencer Gulf, outside of the study area. The species also holds commercial and recreational interest and occurs throughout the Spencer Gulf. This species is considered common across southern Australia, but there are a series of unknowns about the aggregation of cuttlefish including influences on recent declines in numbers (Gaylard et al., | Disturbance to habitat; pollution; oil spill; underwater noise during piling | The species is only afforded protection in the northern waters of Spencer Gulf under the Fisheries Act (SA), it is not EPBC listed. It is likely that minor loss of potential habitat for this species will occur due to the construction of the wharf and MOF. It is expected that ongoing operations at the site will not cause further disturbance to this species, as the species is known to occur around jetties and piers around Southern Australia. While it is likely that cuttlefish utilise the reef and seagrass habitat at Cape Hardy, the area could not be considered of significant importance to the species. There will be no significant impact to this species | N | |

| | nvertebrate Species | | Likelihood of Occurrence | | Likelihood of significant impacts | |
|--|----------------------------|---|---|--|--|--------------------------------|
| po | otentially in the area | a | | | | |
| Species Name | Common Name | Legislative Protection. EPBC, NPW or Fisheries Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | | 2013). It is likely that <i>S. apama</i> are present across the reef and seagrass habitat at Cape Hardy in low numbers. | | as a result of this development. | |
| Pectinidae spp. | scallop (various) | N/A | Possible that species is present in area. Scallops naturally occur in areas with a moderate current flow, supplying generous amounts of plankton on which to feed. Their preferred habitat ranges between | Disturbance to habitat; pollution; oil spills; IMS | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. It is possible that a small number of this species will be disturbed by the development, but it is unlikely that will represent a significant portion of the species range or population therefore it is concluded there will be no significant impact on this species as a result of this development. | N |
| Heliocidaris erythrogramma (and various similar spp.) | Sea urchin | N/A | Possible that species is present in area. This species generally inhabits subtidal rocky reefs throughout SA. It is likely that this species would be found in the reef habitat surrounding the capes at the site, though high densities are unlikely. It is unlikely that Cape Hardy represents a significant site for this species. | Disturbance to habitat; pollution; oil spills; IMS | This species is not listed for protection but is of some commercial and recreational interest and occurs throughout the Spencer Gulf. While a small amount of potential habitat for this species will be impacted by the construction of the wharf and MOF, this will not represent a significant portion of possible habitat for this species. There will be no significant impact on this species as a result of this development. | N |
| Donax deltoides | Pipi, cockle, surf clam | N/A | Unlikely that species is present in area. Pipis prefer to inhabit the swash zone of sandy beaches with consistently high wave energy. Cape Hardy does not have any beaches suitable to support large numbers of pipis (Ferguson, 2013). | Localised altered nearshore hydrodynamic conditions (i.e. vessel scour/propwash); disturbance to habitat; pollution; oil spills; IMS | This species is not listed for protection but is of commercial and recreational interest. However pipis are highly unlikely to occur in the study area due to insufficient suitable habitat. There will be no significant impact to this species as a result of this development. | N |
| Katelysia spp. | Vongole, clams | N/A | Unlikely that species is present in area. The preferred habitat for this species is generally sandy subtidal sediment across tidal flats and estuary mouths. Cape Hardy does not have habitat suitable to support high densities of Vongole, | Pollution; oil spills; IMS | This species is not listed for protection but is of commercial and recreational interest and occurs in the Spencer Gulf with less than 1 tonne of vongole commercially harvested annually across the Spencer Gulf. Vongole are unlikely to occur at Cape Hardy, as their required habitat of tidal flats or estuary mouths does not exist within the study area. There will be no significant impact to this species as a | N |



| | nvertebrate Species | | Likelihood of Occurrence | | Likelihood of significant impacts | |
|----------------------------------|---------------------|---|---|--|---|--------------------------------|
| Species Name | Common Name | Legislative Protection. EPBC, NPW or Fisheries Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | | | | result of this development. | |
| Pseudocarcinus gigas | Giant crab | Fisheries Act (SA) | Unlikely that species is present in area. This species is not listed for protection but is taken by commercial fishers in the Spencer Gulf. This species prefers deeper water (>50 m) habitats and is therefore highly unlikely to utilise the habitat present at Cape Hardy. | Disturbance to habitat; pollution | Females of this species are protected under the Fisheries Act (SA) when carrying external eggs. The species is not listed under any other protection but is of commercial and recreational interest. The giant crab typically inhabits water significantly deeper than the environment at the Cape Hardy study area (>50m). Therefore it is considered unlikely that the study area plays a role in this species' growth, development or survival. There will be no significant impact to this species as a result of this development. | N |
| Pinna bicolor | Razorfish | N/A | Present. Observed during video tow survey. This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. Typically this species prefers lower energy environments with soft substrates, including sparse seagrass beds. This species occurs in the Cape Hardy area, but only low density beds of razorfish were identified. | Localised altered nearshore hydrodynamic conditions (i.e. vessel scour/propwash); disturbance to habitat; pollution; oil spills; IMS | Razorfish occur throughout the Spencer Gulf in higher densities than observed at Cape Hardy. Cape Hardy could not be considered a critical site for razorfish, and are unlikely to be affected by the development, including the construction of the wharf and MOF. There will be no significant impact to this species as a result of this development. | N |
| Zoila friendii ssp. thersites | Black cowry | Reef Watch species of SA conservation concern. | Possible to be present in port area. This species is widely distributed across southern Australia, found on reefs, in seagrass and near artificial structures such as jetty pilings. The shell of this species is popular amongst shell collectors. This species is long-lived (>12 years), and susceptible to overexploitation. It is possible that this species occurs in the project area, as suitable habitat is present. | Loss of habitat; pollution; oil spills; IMS | This species is not listed for protection, though harvesting is controlled under the Fisheries Management Act, 2007. While a small amount of potential habitat for this species will be impacted by the construction of the wharf and MOF, this will not represent a significant portion of possible habitat for this species. The area does not represent critical habitat for this species. This development will not cause significant impact to this species. | N |

Marine debris DEWHA (2009) Threat abatement plan for the impacts of marine debris on vertebrate marine life, May 2009. Prepared for the Department of the Environment, Water, Heritage and the Arts (DEWHA) [Threat abatement plan] available from http://www.environment.gov.au/biodiversity/threatened/publications/tap/marine-debris.html Fish species and family information summarised from: Baker, 2011; Gomon et al., 2008; Edgar, 1997; Kuiter, 1996, unless expressed otherwise.



6.4.2.2 Risks

If large areas of seagrass loss were to occur, a reduced diversity and density of associated infauna and epifauna species would be expected in any resulting unvegetated areas (Stoner, 1980). The loss of macroalgae from reef habitat is expected to cause similar effects for benthic epifauna. The level of impact to epifauna due to loss of seagrass is difficult to predict as communities depend on a complex interaction of variables including but not limited to hydrodynamics, the seagrass species present, patch orientation, edge effects, and the area of sandy 'blowouts' (Edgar and Robertson, 1992; Tanner, 2003). The location of the marine infrastructure avoids area of dense seagrass or macroalgae thereby reducing the likelihood of affects to these habitats. As detailed in Section 6.3, nominated levels of seagrass clearance were based on conservative estimates. As such, the likelihood of further clearance of seagrass and benthic fauna habitat is considered to be unlikely. The benthic fauna and habitats within the port site are typical of those identified throughout the Spencer Gulf and are not identified as a key location supporting endangered or protected species. As such, the consequences are considered to be minor and would result in a local short term decrease in abundance of benthic fauna, but would not result in lasting effects on the local population. As such, the overall risk of unintended additional seagrass clearance on benthic fauna is considered to be low.

A major spill of oil or other chemicals may occur as a result of vessel failure or accident. Based on the experiences of other operating ports, the likelihood of this event occurring is considered to be unlikely. The consequences of a major spill event are considered to be minor, resulting in a short term decrease in the local abundance of benthic fauna. As such, the overall risk of a major spill event affecting benthic fauna is considered to be low.

The distribution of benthic fauna can be driven by differences in sediment biogeochemistry (Reynoldson, 1987). It is considered unlikely that the emissions of iron ore dust or potential spillage of iron ore during the shipping process will lead to any measurable change in sediment chemistry, or the distribution of benthic fauna in any area other than the immediate settling area of an accidental spill (see Section 5.4.2). Iron occurs naturally in marine sediments, but is generally not soluble or bio-available (Canfield, 1989). With this in mind, it is expected that iron ore (magnetite or hematite) which reaches the sediment from dust emissions will not cause elevations in dissolved or bioavailable iron concentrations. Whilst some spillage of iron ore is considered to be almost certain, as the study area is relatively sheltered from a hydrodynamics perspective, the consequences of an ore spillage will be limited to the study area, not result in long term effects on the local population, and are considered to be minor. As such, the overall risk of iron ore spillage on benthic fauna is considered to be medium.

IMS introduction via the port operations poses a high risk to the endemic benthic fauna, as the disturbed areas created during construction along with new settlement surfaces will provide cleared areas which could be exploited by opportunistic IMS. IMS are discussed in more detail in Section 6.8.

6.4.3 Conclusions

The benthic habitats and fauna found at Cape Hardy during the habitat mapping and sediment survey are consistent with the habitats observed in this region of the Spencer Gulf. There are no significant or unique benthic habitats identified within the study area. The limited footprint of the marine infrastructure (< 17ha) will only have direct impacts on the benthic community within the footprint area of the MOF and tug harbour with a permanent loss of subtidal habitat immediately beneath the MOF (<3 Ha from Table 6-3). While there will be a fundamental change to the habitat from the loss of the soft bottom areas, this loss of habitat area is negligible in the context of the study area. It is also expected that hard substrate colonising organisms will be able to utilise the hard surfaces of the MOF and jetty, and over time the colonising organisms will encrust structures and trap particulates enabling more organisms to colonise and increase the complexity of the species utilising the site.



6.5 Marine Megafauna

In South Australia all cetaceans are protected under the State NPW Act, Fisheries Management Act 2007 and the Commonwealth EPBC Act (whether they are listed or not). The following section reviews the range of threatened marine megafauna (whales, dolphins, seals and turtles) that are potentially present within the study area and their conservation status.

6.5.1 Existing Environment and Key Findings

The database and literature reviews identified the protected species that have the potential to occur in waters around the Eyre Peninsula region in Table 6-7. Species are classed as common, uncommon, rare or vagrant based on the frequency which they are seen in the region (Caton *et al.*, 2011a). Table 6-8 summarises the output of the EPBC PMST.

Table 6-7 Protected marine megafauna potentially within study area

| Common name | Scientific name | EPBC listing | NPW listing | Frequency* |
|---------------------------|--------------------------|------------------------|-------------|------------|
| | CET | TACEANS | | - |
| Southern Minke whale | Balaenoptera bonaerensis | Cetacean, Migratory | - | U |
| Bryde's whale | Balaenoptera edeni | Cetacean, Migratory | Rare | R |
| Blue whale | Balaenoptera musculus | Endangered | Endangered | U |
| Pygmy right whale | Caperea marginata | Cetacean, Migratory | Rare | U |
| Common dolphin | Delphinus delphis | Cetacean | | С |
| Southern right whale | Eubalaena australis | Endangered | Vulnerable | С |
| Risso's dolphin | Grampus griseus | Cetacean | Rare | N/A |
| Southern bottlenose whale | Hyperoodon planifrons | Cetacean, Migratory | Rare | R |
| Dusky dolphin | Lagenorhynchus obscurus | Cetacean, Migratory | - | N/A |
| Humpback whale | Megaptera novaeangliae | Vulnerable | Vulnerable | С |
| Killer whale | Orcinus Orca | Cetacean, Migratory | - | С |
| Bottlenose dolphin | Tursiops aduncas | Cetacean | - | С |
| | PII | NNIPEDS | | |
| New Zealand fur seal | Arctocephalus forsteri | Marine | - | С |
| Australian fur seal | Arctocephalus pusillus | Marine | Rare | V |
| Subantarctic fur seal | Arctocephalus tropicalis | Vulnerable | Endangered | V |
| Australian sea lion | Neophoca cinerea | Vulnerable | Vulnerable | С |
| | R | EPTILES | | |
| Leatherback turtle | Dermochelys coriacea | Endangered | Vulnerable | R |
| Loggerhead turtle | Caretta caretta | Endangered | Endangered | R |
| Green turtle | Chelonia mydas | Vulnerable | Vulnerable | R |

^{*}As reported in Caton et al. (2011a) Eyre Peninsula Coastal Action Plan (EPCAP) 2011: C = Common, U = Uncommon, R= Rare, V = Vagrant N/A = not applicable (not reported in EPCAP 2011). **Bold** entries indicate opportunistic sighting during surveys. All marine mammals (and The Great White Shark) are also protected at all times under the Fisheries Management Act 2007.

Common and bottlenose dolphins are known to occur in the Spencer Gulf region year-round (Gibbs and Kemper, 2014). Generally, bottlenose dolphins have a nearshore coastal distribution whereas common dolphins tend to be distributed more evenly across the wider gulf area in open water. A study of common dolphins in Gulf St Vincent established that the species has a preference for depths greater than 14 m (Filby,

2010). Several common dolphins were observed during the field survey (Figure 6-4), trailing the video survey vessel in around 20 m of water and captured on the towed video diving down to inspect the video sled.

Large whale species (Blue, Humpback, Southern right whales and including Killer whales) are migratory in nature and therefore are seasonal visitors to South Australian waters (Gibbs and Kemper, 2014). The Humpback Whale is listed as Vulnerable under both the EPBC Act and the NPW Act. Spencer Gulf is not located on the main Humpback whale migration routes which exist along the eastern and western Australian coasts as depicted in Figure 6-5 (Kemper, 2005; SPRAT, 2014). Humpback whales have been widely recorded in South Australian waters (BDBSA, 2013). With its distinctive humpback fin and large size the species is easily recognisable and regularly reported when it is sighted, as evidenced by the numerous sightings from Port Augusta in 2008 and 2011. However, as the gulf does not represent any known feeding or calving grounds, verified records (assessed for duplication) submitted within the Spencer Gulf are classified as uncommon (Kemper, 2005). There have been concerns raised regarding verification of 'citizen science' and data overlaps from multiple independent databases for whales in Australia (Bannister *et al.*, 1996). This suggests that sightings of Humpback whales (or mother and calf) visiting Spencer Gulf have the potential to be overestimated from multiple sightings of the same vagrants in the gulf.

Blue Whales are listed as Endangered under both the EPBC Act and the NPW Act. Blue whales are migratory in nature, with a widespread oceanic distribution. Blue whale migration paths are unknown but key sighting areas have been identified within South Australia (Figure 6-6) including the Great Australian Bight, near Robe in the south-east through to the open waters of the Bonney Upwelling off western Kangaroo Island (Bannister *et al.*, 1996). Although this species has been recorded within the Spencer Gulf (ALA, 2013), they are normally found in deeper water along the 200 m contour in the Bonney Upwelling area and are unlikely to be observed closer to shore. Being the largest animal on the planet Blue Whales would be highly conspicuous in Spencer Gulf therefore it is unlikely the species frequents gulf waters more than has been reported.



Figure 6-4 Common dolphin photographed trailing video tow survey vessel in study area




Figure 6-5 Known distribution, migration and aggregation areas of Humpback Whales (DEH, 2005a)




Figure 6-6 Recognised aggregation areas of Blue whales in south-east Australia (DEH, 2005b)

The Southern Right Whale (SRW) is listed as Endangered under the EPBC Act and Vulnerable under the NPW Act. The SRW was heavily impacted by the nineteenth-century whaling industry, reducing numbers from an estimated 60,000 individuals to around 300 in 1920 (Bannister, 2001; Bannister, 2007). Since the moratorium on whaling, SRW population numbers have been recovering with numbers worldwide estimated at 7000 in 2007 (NOAA, 2007), with a global population growth rate (~7%/year) approaching the biological maximum for the species (Bannister, 2007). Within Australia the population has been estimated at over 3500 and growing at close to 7% per year (Carroll *et al.*, 2014). The SRW is a circumpolar species living in the sub-Antarctic and open ocean during the summer months and non-calving years. Based on the overwintering sightings of mothers and calves in nearshore areas during nursery aggregation there are four main populations of SRW; in South Africa, Argentina, Australia and sub-Antarctic New Zealand (DSEWPaC, 2012).

Recent studies of the genetics of SRW show that mothers with calves show a high fidelity to the same aggregation areas (NOAA, 2007; Carroll et al., 2011) with the Australian population showing some limited interconnection with the sub-Antarctic New Zealand population (Bannister, 2007, Valenzuela et al., 2009; Carroll et al., 2011; DSEWPaC, 2012; Carroll et al., 2014). Within Australia there are a number of current and known historical aggregation grounds distributed from Western Australia to southern New South Wales as depicted in Figure 6-7 (DSEWPaC, 2012). These different aggregations are now known to be all from the same genetic population (Valenzuela et al., 2009). Historically, there were reported differences between the western and eastern SRW aggregation areas in Australia, likely due to historical separation of the populations during



periods of heavy whaling. Recent genetic studies indicate a relinking of these populations suggesting migration between the areas and inter-breeding (Carroll et al., 2011). Studies also indicate that the Australian distribution of SRW follows a circular, counter-clockwise seasonal migration pattern of a single undivided population (Bannister, 2001; Burnell, 2001). During the winter season whales travel westward along the southern coastline, then south towards summer feeding grounds in the sub-antarctic waters of the Southern Ocean, then eastward in the sub-polar latitudes and then finally north again to their wintering grounds with no difference in distributional movements between males and non-breeding females (Burnell, 2001).

SRW population structure suggests high juvenile and low adult mortality (Bannister *et al.*, 1996). Upon reaching maturity at around six to nine years female SRW generally calve at three-year intervals and mature females are almost never seen in Australian coastal waters in non-calving years, suggesting conception takes place elsewhere (DSEWPaC, 2012). The long interval between calving and fidelity to foraging and nursery grounds means that despite the growth in the SRW population the expansion of the species range is limited by individual animal's fecundity and their offspring's (limited) exploration of new areas.

In Western Australia the SRW population growth rate has been monitored closely and is known to be at or near the maximum population growth biologically possible for the species, however there is little knowledge of the growth rate at other nursery grounds around the country (DSEWPaC, 2012). The SRW that occur in South Australia are closely linked to the western population (Carroll *et al.*, 2011) and therefore are expected to be growing at the same rate. Given that the Australian nursery ground / aggregation populations of SRW have interlinkage (Valenzuela *et al.*, 2009; Carroll *et al.*, 2011)) it can be surmised that the Australian SRW population is growing at around the biological maximum. The population of the SRW in south Western Australia is growing at their maximum biological rate despite the presence of several major port developments in the region. This indicates that there would appear to be no apparent significant impacts from shipping or ports, despite ship strike being listed as a major threat to this species (Bannister, 2007), and that with appropriate management plans in place ports and shipping can coexist with SRW. Similarly, population recovery for SRW in South Africa is occurring despite human caused mortalities (ship strikes and entanglements in fishing gear (Best *et al.*, 2001)), indicating the species population growth rate is not severely impacted by vessel strikes. Within South Australia it has been reported that the greatest threat to SRW mortality from human interaction comes from entanglement rather than vessel strike (Kemper *et al.*, 2008)

SRW are known to be present along the South Australian coast from May to November during their calving season (SPRAT, 2014) although the species tends to aggregate in predictable locations along the coast outside of the Spencer Gulf (Figure 6-7). Within South Australia there are two winter aggregation areas in the Great Australian Bight (GAB) and one aggregation area around Encounter Bay south of Fleurieu Peninsula where mothers come to calve or nurse their young (Bannister, 2007). There are no known current or historical aggregation areas within the South Australian gulfs (Kostoglou and McCarthy, 1991; DSEWPaC, 2012) indicating the gulfs are not part of the species migration path.

The SRW is easily identifiable by the general public and highly conspicuous during their nearshore mother-calve aggregations. Therefore, a single individual, or mother and calf, may be sighted on numerous occasions as they move east to west from one aggregation area to another, as evidenced by SA Whale Centre records (2013-14). The species is considered an occasional visitor to Spencer Gulf (Caton *et al.*, 2010a) with most sightings mainly during the winter months, when females use the shallow waters along Australia's southern mainland for calving (Bannister et al., 1996; Gibbs and Kemper, 2014) (Figure 6-7).

As previously discussed, genetic and isotopic studies of the species indicate that mothers and their offspring return annually to the same feeding and calving grounds (Valenzuela *et al.*, 2009; Carroll *et al.*, 2011), suggesting the species may have limited capacity to explore new feeding grounds (DSEWPaC, 2012). Despite the known sightings of SRW within Spencer Gulf, the gulf is not part of any established or historical migration path. Given the tendency of SRW to show high fidelity to aggregation areas, future high level utilisation of the



Spencer Gulf is considered highly unlikely. Individual SRW, or mother and calf pairs that visit Spencer Gulf are likely moving from one aggregation area to another (Victoria to Encounter Bay to GAB) and not using the area for foraging or nursery grounds.

Records show the SRW rarely strands in South Australia (Kemper and Ling, 1991; Kemper *et al.*, 2008; Caton *et al.*, 2010), with no stranding records along the Eyre Peninsula coast. However, the calves are susceptible to direct disturbance, such as whale watching vessels and/or low flying aircraft, around calving areas. Coastal industrial activities (seismic, drilling), fishing operations (entanglement), pollution and collisions with ships can also impact SRW.




Figure 6-7 Known aggregation locations for southern right whale in South Australia (DSEWPaC, 2012)

Two seal species are considered common in the Spencer Gulf area (Caton *et al.*, 2011a), the Australian sea lion and the New Zealand fur seal. The Australia sea lion is endemic to Southern Australia and only known to occur in South Australia and Western Australia (DSEWPaC, 2013; SPRAT, 2014). The entire Spencer Gulf, including the study area, is a known foraging range for Australian sea lions (Goldsworthy *et al.*, 2007). The species range is vast covering approximately 30% of the 178,000 km² of potential habitat from Western Australia to Kangaroo Island, with individual Sea lions known to travel around 70 km while foraging (DSEWPaC, 2013). Aerial surveys (Shaughnessy *et al.*, 2005) indicated several pupping sites around the western and southern coast of the Eyre Peninsula. A number of pupping sites exist in the southern Spencer Gulf including English Island, Lewis Island and Dangerous Reef (PIRSA, 2008), with the closest site to the study area being in Sir Joseph Banks Conservation Park including Langton Island (Shaughnessy *et al.*, 2011) some 45 km to the south of Cape Hardy. While the species is capable of utilising all waters in South Australia they tend to aggregate in breeding colonies

where they also focus their foraging effort, (see Figure 6-8). Therefore Australian Sea lions are unlikely to be heavily reliant on the waters surrounding Cape Hardy.




Figure 6-8 Geographic distribution of foraging Australian Sea lions across South Australia (high: red, medium: orange, low: blue) (DSEWPaC, 2013)

New Zealand fur seals also breed in South Australia, however all known pupping sites for the species are outside the Spencer Gulf on offshore islands either to the west or south of Eyre Peninsula and Kangaroo Island (Shaughnessy, 1999; Shaughnessy *et al.*, 2005; Goldsworthy and Page, 2009). Outside of the breeding season the South Australian colonies of New Zealand Fur Seals are known to forage from the coastline near Fowlers Bay to southern Tasmania and offshore out to 1000 km south of the continental shelf break. While adult males and juveniles are found in both gulfs adult females tend to use offshore waters (Goldsworthy and Page, 2009). Therefore although the species is likely to occur at Cape Hardy, the study area is unlikely to represent significant habitat for either breeding or foraging.

The three turtle species (leatherback, loggerhead and green) that have been recorded in Spencer Gulf are considered only occasional visitors to the area as none of the species use haul out beaches or breed within Southern Australia. They are rarely seen, with several of the records for turtles in Spencer Gulf based on dead animals washed ashore or found floating (Caton *et al.*, 2010; ALA, 2013). These species generally inhabit tropical waters in Australia from northern New South Wales, Queensland, and Northern Territory around to the Pilbara in Western Australia. All marine turtles are migratory, leading essentially pelagic lives. The males do not come ashore once hatched and tend to have large ranges that can extend into temperate waters as they forage or seek out access to new areas of females. Females are predominantly found in the tropics and only come ashore to dig nests and lay eggs. There are no known nesting sites for any marine turtle species in Southern



Australia (ICUN, 2013; IOSEA, 2013; SPRAT, 2014). Therefore the records of marine turtles in Spencer Gulf are most likely for vagrant males.

6.5.2 Impacts and Risks: Marine Megafauna

Impacts could occur to marine megafauna as a result of:

- Loss of habitat through clearance or exclusion
- Disturbance to breeding colonies or haul-out beaches
- Noise emissions during construction and operation
- Vessel interaction and increased likelihood of ship strike as a result of increased vessel traffic
- Marine debris
- Pollutant build-up or spills including oil
- Introduction of disease

6.5.2.1 Impacts

As previously outlined, the study area of the proposed port does not represent critical habitat, including haulout or nursery beaches, foraging areas, mating area or migration path for any identified marine megafauna. As such, the loss of intertidal and subtidal habitat (as discussed in Section 6.2 and 0) is not anticipated to significantly impact marine megafauna.

The key identified impact to marine megafauna is as a result of noise emissions from the construction and operation of the proposed port. Impacts to marine fauna (including megafauna) from noise are discussed in detail in Section 6.7. In summary, the predominant source of noise is piling during construction. Vessel movements during both construction and operation also represent a key noise source. Noise emissions will alter the behavioural patterns of marine megafauna. Impacts will vary between individuals as an animal's response to construction noise is dependent on the individual animal's sensitivity and curiosity. Animals that initially display behavioural responses such as avoidance when encountering a new noise may quickly become habituated to the noise. For example bottlenose dolphins are commonly observed around the world inhabiting operating ports despite ongoing noise emissions (ADS, 2007).

Soft start piling procedures will be implemented prior to the commencement of piling operations. Soft start piling provides the opportunity for marine fauna to vacate an area and avoid the full impact of noise from piling. Noise modelling for the proposed port facility has taken into account underwater noise from vessels and predicted that there would be no noise impacts to low-frequency cetaceans beyond 10 m from the vessel propeller (Sonus, 2014). Due to the sound propagation through water it is likely that SRW or other megafauna would hear vessels long before damage could occur and move away from the source. As such, noise emissions during construction and operation of the proposed port are not anticipated to result in physical injury to marine megafauna. Behavioural patterns of megafauna are expected to change as a result of ad hoc noise emissions during construction and operation. Noise emissions will predominately occur within the study area with affects beyond this area not anticipated. As such, noise emissions from the proposed port are expected to result in a long term change to behavioural patterns of megafauna within the study area, considered to be a medium impact.

An assessment of the likelihood of occurrence and impacts to each marine megafauna potentially occurring within the study area is provided in Table 6-6 below.

Table 6-8 Marine megafauna: likelihood of occurrence and potential impacts

| Potential mar | rine megafauna at t | he port Site | Likelihood of occurrence | Likelihood of signifi | icant impacts | |
|-------------------------------|---|---|---|---|---|--------------------------------|
| Species name | Common name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | Mammals | | | | | |
| Arctocephalus forsteri | New Zealand Fur seal | Marine (EPBC) | Possible. There are four BDBSA records for this species at Lipson Island (ALA, 2013) and one record at Port Neill, both confirmed by the SA Museum (ALA, 2013). No BDBSA records exist for this species within 10 km of the site (BDBSA, 2012). There are no known haul-out areas or breeding colonies for this species within 100 km of the development area (Shaughnessy et al., 2005). This species may pass in the vicinity of the development site however the size and location of the development will not significantly impact critical habitat for this species. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Oil spill; loss of habitat through clearance or exclusion; disturbance to breeding colonies or haul-out beaches; vessel impacts; noise impacts; entanglement in marine debris introduction of disease | There is no potential for the development to impact on breeding colonies or haul out beaches required by this species as there are none present in the development area. The development does not represent critical habitat or affect a large area of the gulf, therefore, no significant impacts on this species are likely from clearance or exclusion from foraging areas. Therefore impacts on this species from the development are not expected to be significant. | N |
| Arctocephalus pusillus | Australian Fur seal, Australo- African Fur seal | Marine (EPBC); Rare (NPW) | Possible. No live sightings of this species have been recorded within 10 km of the development area (ALA, 2013; BDBSA, 2013). There are no known haul-out areas or breeding colonies for this species within 100 km of the development area (BDBSA, 2012). There are two records for the species from the SA Museum representing preserved specimens collected at Port Neill and Tumby Bay in 2009 and 2011 respectively (ALA, 2013). This species may pass in the vicinity of the development site; however, the size and location of the port site mean it will not disrupt critical habitat for this species. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Oil spill; loss of habitat through clearance or exclusion; disturbance to breeding colonies or haul-out beaches; vessel impacts; noise impacts; entanglement in marine debris introduction of disease | There is potential for this species to pass within the vicinity of the port site. There is no potential for the development to impact on breeding colonies or haul out beaches as there are none present in the area. The development area does not represent critical habitat or a large area of the gulf; therefore no significant impacts to this species are likely from clearance or exclusion from foraging areas. Therefore impacts on this species from the development are not expected to be significant. | N |
| Balaenoptera acutorostrata | Minke whale | Whales and other Cetaceans | Unlikely. This species was not recorded opportunistically during Jacobs site surveys, or from recent marine surveys at Port Spencer (Golder Associates, 2011). No records for this species occur within the Spencer gulf (ALA, 2013; BDBSA, 2013). The species' range | Vessel collision; entanglement in marine debris; oil | Given the species is unlikely to frequent the area (based on historical sighting information (ALA, 2013; BDBSA 2013), has | N |

| Potential mar | rine megafauna at tl | ne port Site | Likelihood of occurrence | Likelihood of signifi | cant impacts | |
|--------------------------|----------------------|---|--|---|---|--------------------------------|
| Species name | Common name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | (EPBC); Rare (NPW) | includes coastal and offshore waters of the Pilbara, southern Australia including Tasmania and up to Cape Yorke in Queensland (SPRAT, 2014). The Common Minke Whale is found in all oceans from 65°S to 80°N (ICUN, 2013). Population estimates for Common Minke whales from the Northern hemisphere exceed 100,000 individuals while estimates for the Southern hemisphere are not available due to sighting data not distinguishing between Common and Dwarf species (ICUN, 2013). During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | spills; noise impacts – seismic, drilling, piling | a large geographic range and a total global population above any threshold for threatened species (ICUN, 2013), impacts on this species from the development are not expected to be significant. | |
| Balaenoptera edeni | Bryde's whale | Migratory, Whales and Other Cetaceans (EPBC); Rare (NPW) | Unlikely. This species was not recorded opportunistically from Jacobs' surveys or recent marine surveys at Port Spencer (Golder Associates, 2012). There are no records of this species within 50 km of the study area (BDBSA, 2012). Two records for this species exist in the Spencer Gulf - one stranding at Corny Point at the tip of Yorke Peninsula in 1971 from the SA Museum and one unverified and undated sighting at Port Augusta (ALA, 2013). The species' is not known to frequent Spencer Gulf and the recorded sightings of this species within South Australia are limited to stranding records (SPRAT, 2014). Historically, Bryde's Whales have been confused with other species including the Sei Whale. Bryde's Whales are likely to be found along either east or west coasts of Australia and less so along the southern coast (Bannister <i>et al.</i> 1996). Any individual or group of this species entering the gulfs would therefore be considered a vagrant visitor. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling | The development site does not coincide with critical habitat for this species. Therefore, although Bryde's whales may enter the Spencer Gulf it is unlikely that the Bryde's whale travels in the vicinity of the harbour as part of its natural range. It is unlikely for the development to have any significant impact on this species | N |
| Balaenoptera musculus | Blue whale | Endangered, Migratory,, Whales and Other Cetaceans, Marine (EPBC); Endangered (NPW) | Unlikely. No records of this species have been submitted to the BDBSA within 50 km of the study area (BDBSA, 2012). The proposed port site is not in the vicinity of known feeding areas, migration paths or breeding grounds. This species is the largest animal to exist on the planet and therefore is highly conspicuous. Given the lack of sightings for the species in Spencer Gulf, it is highly unlikely to frequent the port site. While this species may have the potential to pass through gulf waters, the development area does not represent critical habitat for this species and therefore the presence of the species is unlikely. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling | It is unlikely for this species to pass in the vicinity of the development site as the gulf is not critical habitat for this species. As the port site is not within critical habitat for the species the species is unlikely to frequent the development. Therefore impacts on this species from the development are not expected to be significant. | N |

| Potential mar | ine megafauna at t | he port Site | Likelihood of occurrence | Likelihood of signifi | cant impacts | |
|------------------------|--|---|---|--|--|--------------------------------|
| Species name | Common name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| Caperea marginata | Pygmy right whale | Migratory, Whales and Other Cetaceans (EPBC); Rare (NPW) | Unlikely. No records of this species have been recorded in the BDBSA within 50 km of the area (BDBSA, 2012). The species' range within South Australia does not include Spencer Gulf. This species has the potential to pass in the vicinity or enter the Spencer Gulf; however, the development will not interfere with critical habitat for this species. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling | It is unlikely for this species to pass in the vicinity of the development site as the gulf is not critical habitat for this species. It is unlikely for the development to have any significant impact on this species | N |
| Delphinus delphis | Common dolphin, Short- beaked common dolphin | Whales and Other Cetaceans (EPBC) | Present. This species was sighted during the underwater video tow component of the surveys at the port site (SKM). This species is found globally in tropical, subtropical and temperate waters of the Atlantic, Pacific and Indian Oceans with a population in excess of 3 million (SPRAT, 2014). The extent or population of the Common Dolphin in Australia has not been estimated but it is known to inhabit all coastal and offshore regions of Australia with no fragmentation of the populations (SPRAT, 2014). During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling | This species was observed during the marine survey at the site. As this species is highly mobile with a large global range the development area does not represent a significant area of critical habitat for this species nor would any impacts from the port pose a significant impact to the species. It is unlikely for the development to have any significant impact on this species. | N |
| Eubalaena australis | Southern right whale | Endangered, Migratory, Whales and Other Cetaceans, Marine (EPBC) | Possible. The port site is not considered critical habitat for this species. The known calving and feeding grounds for this species do not include Spencer Gulf. The range of this species in Australia covers southern waters from Exmouth in WA to Harvey Bay in QLD, including Tasmania. There are several calving grounds in Australia: parts of WA, the Great Australian Bight outside of the SA gulfs, Warnambool Victoria, Tasmania and NSW (SPRAT, 2014). The species only frequents Australian coastal waters during the winter months and over summers in the Southern Ocean or Sub-Antarctic waters with populations found in South Africa, New Zealand and Argentina. Kemper (SA Museum) noted that sightings of the species have been made less than 10 times within Spencer Gulf since 1948, including at the port site in 1983 and 1984 (BDBSA, 2012). This species is rather conspicuus and sightings of individual animals are often logged multiple times. Recent sightings around Lipson Island (Golder Associates, 2013) indicate the species may pass within the vicinity of the port site but the development site does not represent critical habitat for this species. The occurrence of this species at the port would be considered transient; as the gulf is not a significant migration path or critical breeding area (Bannister et al. 2005). During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine | Direct disturbance; vessel collision; entanglement in marine debris, oil spills; noise impacts – seismic, drilling, piling | Southern Right Whales may travel within Spencer Gulf sporadically, including in the vicinity of the port site. However, the gulf is not a significant migration path or critical breeding area. It is therefore considered unlikely the species would be present the majority of the time. As the port site does not represent critical habitat for the species the development is not expected to have a significant impact on the Southern Right Whale. It is unlikely for the development to have any significant impact on this species. | N |

| Potential mar | rine megafauna at tl | ne port Site | Likelihood of occurrence | Likelihood of signifi | elihood of significant impacts | |
|----------------------------|-----------------------------|--|--|---|--|--------------------------------|
| Species name | Common name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | | mammals and turtles. | | | |
| Grampus griseus | Risso's Dolphin, Grampus | Whales and Other Cetaceans (EPBC) | Unlikely. No records of this species have been recorded in the BDBSA within 50 km of the area (BDBSA, 2012). The species are considered to be widely distributed around the Australian mainland and are found in tropical and sub-tropical coastal and shallow offshore waters throughout the Indian Ocean, Indo-Pacific and Pacific Ocean (SPRAT, 2014). This species has the potential to pass in the vicinity or enter the Spencer Gulf; however, the development will not interfere with critical habitat for this species. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling | It is likely that suitable habitat for Grampus exists near the study area, and it is possible that this species may occur as a visitor to the area. The habitat near the study area is, however, unlikely to sustain an important population of this species. Given the limited disturbance of marine habitat and the lack of sightings of Grampus in the wider area, unlikely for the development to have any significant impact on this species. | |
| Lagenorhynchus obscurus | Dusky dolphin | Migratory, Whales and Other Cetaceans, Marine (EPBC) | Unlikely. No records of this species have been submitted to the BDBSA within 50 km of the study area (BDBSA, 2012). The only record for this species in South Australia comes from the southern Coorong, representing a specimen collected by the SA Museum in 2003 (ALA, 2013). The species' presence in Australia is only known from 13 reports since 1828. This species has an estimated distribution from Western Australia to Tasmania (SPRAT, 2014). There are well documented populations of the species in New Zealand, South America and South Africa with its global population considered to be moderately abundant across the sub-Antarctic regions (SPRAT, 2014). Due to the paucity of records for this species in Australia, its presence at the port site is unlikely. As the species is thought to only move between Australia and other countries (SPRAT, 2014), the port site is not considered to represent critical habitat for this species. The population size of Dusky Dolphins is unknown; however, the low stranding and sighting rates suggest these dolphins are uncommon in Australian waters (Bannister et al. 1996). Due to the geographic range and low frequency of sightings for this species in South Australia, impacts from the development on this species are not expected to be significant. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling | As the proposed development is not in the vicinity of known breeding or significant foraging areas it is considered unlikely that this species will be present at the site. It is unlikely for the development to have any significant impact on this species. | N |
| Megaptera novaeangliae | Humpback whale | Vulnerable, Migratory, Whales and Other Cetaceans | Unlikely. No records have been submitted to the BDBSA for this species within 50 km of the study area (BDBSA, 2012). Kemper (SA Museum) noted that there were less than 10 sightings of this species within the Spencer Gulf and there are no known calving grounds for the species off southern Australia as the species prefers tropical waters for breeding (Kemper, 2005). The Spencer Gulf does not form part of any migration path, feeding or | Vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, | As the proposed development is not in the vicinity of known breeding areas, migration paths or significant foraging areas, it is considered unlikely that this species will be present. It is unlikely for | N |

| Potential marine megafauna at the port Site | | he port Site | Likelihood of occurrence Likelihood of significant impacts | | | | |
|---|---|--|---|--|--|--------------------------------|--|
| Species name | Common name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | Marine (EPBC); Vulnerable (NPW) | calving ground for this species (Bannister <i>et al.</i> , 2005; SPRAT, 2014). Despite the potential for species to enter the gulf, the port site is not considered critical habitat for the Humpback Whale. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | drilling, piling | the development to have any significant impact on this species. | | |
| Neophoca cinerea | Australian sea lion | Vulnerable, Marine (EPBC) | Unlikely. No BDBSA records of this species have been submitted within 10 km of the area (BDBSA, 2012). There is a record of an opportunistic sighting of a Sea lion taking a Little penguin at Lipson Island (Madden-Hallett <i>et al.</i> , 2011). The most current BDBSA records of the species were a haul-out area at Point Gibbon >100 km from the Study area in 1992 (DEWNR 2013) with less than 10 individuals and no record of the species in the area since 2000. Although the development is within the known range of the species and may provide suitable habitat it is unlikely the species would frequent the beach near the development. The site is not close to any known breeding sites or existing haulout beaches. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Loss of habitat; disturbance; vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling; introduction of disease | The proposed development is not in the vicinity of known breeding or significant foraging areas, it is considered unlikely the species would be present. It is unlikely for the development to have any significant impact on this species. | N | |
| Orcinus orca | Killer whale, Orca | Migratory, Whales and Other Cetaceans, Marine (EPBC) | Unlikely. Orcas (Orcinus orca) are a well-known distinctive species that move freely between southern and northern hemispheres, occurring all around the world though in greater abundant in cooler waters. Orcas have been recorded in all Australian states, but not in the Northern Territory. Concentrations of an Orca population are believed to occur around Tasmania, and sightings occur in South Australia and Victoria. Orcas are not known to be migratory, and seasonal movements are thought to relate to changes in food supply. Orcas are considered a 'data deficient' species by the IUCN, reflecting the level of unknowns about the world population. There have been several records in South Australia, including one reported stranding near Tumby Bay (approximately 30 km south of the study area) in 2000 (ALA, 2013). During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. | Vessel collision; entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling | It is likely that suitable habitat for Orcas exists near the study area, and it is possible that this species may occur as a visitor to the area. The habitat near the study area is, however, unlikely to sustain an important population of this species. Given the limited disturbance of marine habitat and the lack of sightings of Orcas in the area, unlikely for the development to have any significant impact on this species. | N | |
| Tursiops aduncus | Indian Ocean bottlenose dolphin, Spotted | Whales and Other Cetaceans (EPBC) | Possible. Two records of this species were recorded in Spencer Gulf in 1998 and 2000 by the SA Museum as identified by Kemper (ALA, 2013). The species <i>Tursiops aduncus</i> are considered to be widely distributed around the Australian mainland and are found in tropical and sub-tropical coastal and shallow offshore waters throughout the Indian | Vessel collision; entanglement in marine debris; oil spills; noise | The species is highly mobile and may pass in the vicinity of the port site however the development site is not critical habitat for this species. It is unlikely for the | N | |

| Potential marine megafauna at the port Site | | | Likelihood of occurrence | Likelihood of signifi | Likelihood of significant impacts | | |
|---|-----------------------|---|--|--|---|--------------------------------|--|
| Species name | Common name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Tursiops | bottlenose dolphin | Whales and Other | Ocean, Indo-Pacific and Pacific Ocean (SPRAT, 2014). However, recent studies of the genetics of Australian bottlenose dolphin suggest that the population in southern Australia waters described by Kemper (Kemper, 2004) may be a separate, as yet undescribed taxon of dolphin (Möller et al., 2008). Regardless, this species would be protected under the EPBC Act as a cetacean and under State legislation (NPW Act, 1972). As the species has been documented at Tumby Bay (ALA 2013) there is the potential for it to pass in the vicinity of the development area. However as the size and location of the port site does not represent a substantial area of habitat for this species it is not expected that the construction and operation of the port would pose a significant impact to this species. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles. Possible. This species occupies a broader range of habitats than any other marine mammals it can be found in octuaries and coastal waters or open water polarie areas. | impacts – seismic, drilling, piling Vessel collision; | development to have any significant impact on this species. This species is highly mobile and may pass in the vicinity of the port site. However, | N | |
| truncatus s. str. | dolphin | Cetaceans (EPBC) | mammal; it can be found in estuaries and coastal waters or open-water pelagic areas, live in large oceanic pods or small family groups, sometimes with defined territories. This species may pass in the vicinity of the development area or even use the area as a fishing ground since they have been recorded at Tumby Bay (ALA 2013). However, the size and location of the port site does not represent a large area of critical habitat for this species. The species is well adapted to human settlement with well documented pods living in operating ports such as Port Adelaide or frequenting urban waterways in WA and QLD. Given the range and adaptability of this species it is considered unlikely that the operating port would have any significant impacts on the species. During construction of the wharf and MOF a CEMP will be developed to minimise impacts to all species of marine mammals and turtles and an OEMP will be developed for the operation of the port to protect all species of marine mammals and turtles | entanglement in marine debris; oil spills; noise impacts – seismic, drilling, piling | in the vicinity of the port site. However, the development is not critical habitat for this species. It is unlikely that the development will have any significant impact on this species. | | |
| | Reptiles | | | | | | |
| Caretta caretta | Loggerhead turtle | Endangered, Migratory, Marine (EPBC) | Unlikely that this species would be present. Key breeding and foraging habitat occurs in tropical areas of Australia, far away from the study site. The development site is not a known habitat for the species. This turtle is a marine species not capable of using terrestrial area and is unlikely to frequent Cape Hardy. The Spencer Gulf is not a known habitat for this species and as such it is unlikely this species would be found in the gulf. There are no know haul out beaches in SA. No records of this species have been submitted to the BDBSA within 50 km of the area (BDBSA, 2012).No opportunistic | Entanglement in marine debris; vessel impacts; noise impacts. | The development site or gulf is not a known critical habitat for the species. It is unlikely that this species would be present and therefore no significant impact on this species is expected from the development. | N | |



| Potential marine megafauna at the port Site | | | Likelihood of occurrence | Likelihood of occurrence Likelihood of significant impacts | | |
|---|---|---|--|--|--|--------------------------------|
| Species name | Common name | Legislative Protection e.g. EPBC or NPW Status | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | | sightings of this species were recorded during Jacobs surveys. | | | |
| Chelonia mydas | Green turtle | Vulnerable, Migratory, Marine (EPBC) | Unlikely that this species would be present. The key nesting sites for this species are in NT, WA and QLD and females rarely leave the tropics. Males (and young turtles <10 years) may travel through temperate waters however temperate oceanic waters are not critical habitat for this species. The species may pass in the vicinity of Spencer Gulf however it is unlikely to be found in the gulf. As females do not leave the tropics and males do not come ashore, Cape Hardy could not be considered a haul out beach for these turtles. No records of this species have been submitted to the BDBSA within 50 km of the area (BDBSA, 2012).No opportunistic sightings of this species were recorded during Jacobs surveys. | Entanglement in marine debris; vessel impacts; noise impacts. | As the port site is not within the vicinity of known feeding or breeding habitat it is considered highly unlikely that there would be a significant impact to this species from the development. | N |
| Dermochelys coriacea | Leatherback turtle, Leathery turtle, Luth | Endangered, Migratory, Marine (EPBC) | Highly unlikely to occur in the area. This species has no known breeding habitat in Australia and as it is a pelagic feeder it is unlikely that the Leatherback Turtle would frequent the Cape Hardy area. Any individual found in the gulf would be a vagrant and not represent a significant number of the species. No records of this species have been submitted to the BDBSA within 50 km of the area (BDBSA, 2012). No opportunistic sightings of this species were recorded during Jacobs surveys. | Entanglement in marine debris; vessel impacts; noise impacts. | It is unlikely for this species to be found in the gulf and any individual found in the gulf would be a vagrant and not representative of the typical behaviour of the species. As the characteristics of the Spencer gulf provide neither critical habitat or breeding sites for the species it is highly unlikely that the development would have a significant impact on the species. | N |



6.5.2.2 Risks

An increase in boat traffic and manoeuvring during the construction and operation of the proposed port poses a risk of direct contact such as vessel or propeller strike with large marine fauna including whales, pinnipeds, dolphins and turtles. Vessel speeds within the port site will be limited to 6 knots to minimise the likelihood of striking megafauna. Similarly, large cargo vessels will be manoeuvred to the jetty at low speed by tugs. Despite the implementation of vessel management procedures, it is not possible to completely negate the risk of ship strike as whales, pinnipeds, dolphins and turtles can be highly mobile and unpredictable in their movements, often remaining below the surface for substantial periods of time before surfacing.

The potential for ship strike relates to an increase in vessel numbers entering the Spencer Gulf. Currently there are six operating ports in Spencer Gulf (Port Lincoln, Wallaroo, Whyalla, Port Bonython, Port Pirie and Port Augusta) which account for over 4200 annual ship movements (Flinders Ports, 2014). The proposed port at Cape Hardy would accommodate approximately 145 vessels per year (290 vessel movements, or roughly 6 vessel movements per week). The additional 290 vessel movements a year associated with the proposed port would represent a 7% increase to vessel movements in the Spencer Gulf. This figure does not include recreational or commercial vessels based in small harbors within Spencer Gulf and therefore the overall percentage increase in vessel movements is likely far less.

It is generally accepted that the risk of boat strike increases with vessel speed (Hazel *et al.* 2007; Kite-Powell *et al.* 2007; DeAngelis *et al.* 2011) and that the most lethal or severe injuries are caused by vessels greater than 80 m when travelling over 15 knots (Laist *et al.*, 2001). The slower a vessel is moving the more time is available for mobile marine fauna to recognise the danger and take action to avoid a collision (van der Hoop et al., 2008; Gende *et al.*, 2011; van der Hoop et al., 2012). In addition, the force of a strike is reduced at slower speeds. If marine fauna is visible to the vessel master, then action may also be taken by the vessel master to avoid a collision, by reducing speed or changing direction. However, such action is dependent upon the size of the vessel and safety considerations. Generally, small vessels are more manoeuvrable than large vessels, and have greater capacity to be navigated to avoid collisions. Capesize vessels can take several hundred metres to reduce speed or change course, and avoidance actions are rarely practical in nearshore waters where navigation channels are relatively narrow. As vessels entering or exiting the proposed port would be restricted to less than 6 knot by the port operator, marine fauna would typically have time to avoid collision and/or be spotted by the vessel master.

Water depth also plays a role in determining the risk of boat strike. Shallow seagrass habitats used for foraging by some dolphins and pinnipeds are more likely to incur ship strike than deeper areas where there is significant clearance between the vessel hull and the seabed (Hazel *et al.* 2007). Deeper areas also allow large marine fauna to dive to avoid a collision. In this context, the risk of boat strike is likely to be higher within shipping channels surrounded by shallow seagrass habitats than in deeper habitats offshore. As the Cape Hardy site is located in area of the gulf with a width of over 50 km, water depth exceeding 20m and away from dense seagrass, marine fauna are at a reduced risk of collision compared to ports further north in the gulf.

SRW appear to be the primary species involved in vessel collisions in the southern hemisphere. This is in part due to the species use of nearshore waters during certain times of the year and the physical nature of this species in that the dead SRW have a tendency to float. It is a known attribute of the species to be neutrally buoyant and that carcasses float for several days (Kemper et al., 2008; Silber et al., 2010). Therefore numbers of recorded SRW strikes based on carcasses washed ashore, sighted at sea or even struck after they die may be biased by the very nature of these whales physiology. Nevertheless there are low numbers of recorded strikes in Australasian waters (Van Waerebeek et al, cited in SEWPaC, 2012). This observation is supported by the low number of strikes on SRW recorded by the IWC in Australian waters. Two fatal vessel collisions and three non-fatal collisions were recorded in Australian waters in the period 1950-2006 (Kemper et al, 2008). From 2007 to

2011, an additional three ship strikes on SRW, including two deaths, have been reported to the IWC (IWC, 2011). The data set from 1950-2006 shows instances of entanglement (often from fishing, aquaculture and related industries) are nearly 3 times more common than boat strikes (5 vessel collisions, 13 entanglements) (Kemper et al, 2008). Within South Australia the main areas of vessel traffic are into Port Adelaide with vessels entering or exiting Backstairs passage and Investigator Strait (Figure 6-9). Backstairs passage and Investigator Strait are part of the SRW migration route, yet there has been only one confirmed fatal strike attributable to a large vessel (vehicle and passenger ferry) which occurred at night while the vessel was travelling between Kangaroo Island and the mainland at a speed of 15 knots (Kemper et al, 2008). It should be noted that most of recorded incidents are thought to relate to smaller fast moving vessels.




Figure 6-9 Map of shipping routes in and adjacent to Spencer Gulf, based on received Automatic Identification System data. Areas of high shipping traffic are shown in red. Map generated from http://www.marinetraffic.com/en/

The IWC recognises that many vessel strikes to whales may go unreported or undetected due the majority of collisions occurring in open (international) waters. Vessel masters are not required to report all known collisons, however the IWC maintains a register of recorded ship strikes. The Spencer Gulf is not part of any known whale migration route and has substantially less vessel traffic than Gulf St Vincent (Flinders Ports, 2014). Due to the hydrodynamic nature and prevailing winds of Spencer Gulf it is highly likely whales killed by vessel strike in or near the mouth of the gulf would be readily found within the gulf (Kemper *et al*, 2008). Therefore as the number of vessel movements within Spencer Gulf already annually exceed 4000 (Flinders Ports, 2014), the low number of recorded ship strikes on whales in South Australia it is considered that the likelihood of a vessel collision in Spencer Gulf is low.

As expected, wherever possible, most megafauna will avoid contact with moving vessels. However, dolphins, porpoises and some pinipeds often actively seek out moving vessels and swim close alongside in the bow wave which may make them vulnerable to injury from collision. Generally, the risk of collisions with marine mammals is greater for recreational craft and dolphin-watching boats, and subsequently guidelines have been developed for minimising the disturbance to dolphins and whales from these activities (DEH, 2006).



Given the implementation of vessel management procedures, the historical likelihood of ship strike in Australian waters and the location of the proposed port outside of known critical habitat, foraging areas, breeding ground or migration paths for megafauna, the likelihood of ship strike associated with construction and operation of the proposed port is considered to be possible. Should an individual be struck, the consequences are considered to be minor, given that it will not affect the overall viability of the species. As such, the overall risk of ship strike is considered to be low.

Construction and operation of the proposed port will result in the generation of a number of waste streams that may enter the marine environment. Marine debris can significantly affect megafauna as a result of pollution, injury through collision, entanglement or ingestion of non-biological products.

Waste management and handling procedures developed as part of the CEMP and OEMP will control waste streams with the overarching aim of no waste products entering the marine environment. Despite the implementation of control measures, it is considered almost certain that some form of waste / debris will enter the marine environment during construction and operation of the port. As the port site is not a key habitat or breeding area, the consequences of debris entering the marine environment are considered to be minimal; insignificant to the overall viability of megafauna. As such, the overall risk is considered to be low.

Megafauna are also at risk from the unintended discharge of pollution in the form of site run-off, wastewater, hydrocarbons or chemicals. With any marine activity there is potential for waste or spills to enter the marine environment. The implementation of design measures (refer Section 4) for the control of surface and waste water at the port site will limit the majority of hazardous pollutants from entering the marine environment. There remains a risk that a major spill of oil or other chemicals may occur as a result of vessel failure or accident. Based on the experiences of other operating ports, the likelihood of this event occurring is considered to be unlikely. As previously outlined, the port site does not represent a critical habitat or breeding colony for megafauna, although some species (e.g. seal) may forage in the area. The consequences of a major spill event are considered to be minor, due to the low level of dependence of megafauna on the study area, resulting in a short term decrease in the local abundance of fauna. As such, the overall risk of a major spill event affecting megafauna is considered to be low.

6.5.3 Conclusions

A wide range of marine megafauna may be present in the area at different times or for limited periods; however a number of species have a higher likelihood of occurring in the waters around Cape Hardy. Australian sea lions, New Zealand fur seals, bottlenose dolphins and common dolphins are likely to be present in the area on a year-round basis. The majority of the marine megafauna potentially in the area are expected to be infrequent visitors to the site or the Spencer Gulf. One whale species, the SRW, may occur in low numbers on a seasonal basis.

Piling and underwater noise is considered to represent the greatest impact to marine megafauna, however this impact is for a limited duration and through the implementation of management procedures (such as soft start piling and observation zones) impacts are expected to be medium.



6.6 Protected Fish Species

The Spencer Gulf supports a diverse range of fish species due to the diversity of habitat niches extending from deepwater rocky reefs, to sandy beaches, seagrass beds and mangroves in the upper gulf. Many areas within the gulf provide nursery and spawning areas for protected or commercially important fisheries species. This section identifies which fish species of conservation value are present within the study area. Commercially important fish species are addressed in Section 6.9.

6.6.1 Existing Environment and Key Findings

The Spencer Gulf supports a high diversity of fish species due to its range of marine ecosystems, which provide suitable habitats for both endemic and migratory fish species. In the Cape Hardy area the predominant habitats are nearshore rocky reef, bare sediments and seagrass beds, which provide nursery and feeding area for fish, including commonly observed species of pipefish, toadfish and leatherjackets (Bryars, 2003).

During the towed video surveys several fish were observed including a leafy seadragon (protected under the Fisheries Management Act and an EPBC listed marine species) within a patch of dense *Amphibolis* and *Posidonia* seagrass meadow.

The desktop study identified 77 fish species as potentially occurring in the study area, with 29 fish species protected under either the Fisheries Act, EPBC Act, or both. Several additional fish species were also opportunistically sighted during the video-tow survey. With the exception of the leafy seadragon, none of the species observed in the area are protected under legislation. As survey transects were not designed to detail the presence, absence, abundance, density or species composition of fish in the area, this is simply a record of incidental observations.

The EPBC protected species were predominantly pipefish or seahorse species with the other species of note considered likely to occur in this region the great white shark (*Carcharodon carcharias*), shortfin mako (*Isurus oxyrinchus*) and porbeagle (*Lamna nasus*). The great white shark is protected under the EPBC Act and listed as 'Vulnerable' and the mako and porbeagle are listed as 'Migratory'. Great white sharks are also protected under the Fisheries Act. The great white shark is known to feed in the Eyre Peninsula region, especially in the vicinity of seal pupping colonies. It is also thought that great white sharks may breed in the Spencer Gulf and inshore coastal waters may be important nursery grounds (DEWHA, 2007).

6.6.2 Impacts and Risks: Protected Fish species

There is potential for a number of fish species to be present within the seagrass, reef and soft bottom areas of the site however fish species in the port area are unlikely to suffer high impacts from the ongoing operation of the port.

Impacts to protected fish could occur as a result of:

- Loss of habitat through clearance, exclusion or degradation of the environment
- Pollution from run-off, oil or chemical spills
- Marine debris
- Noise and physical impacts during construction piling
- Artificial light spill
- IMS



6.6.2.1 Impacts

As previously outlined, the study area of the proposed port does not represent critical habitat for any identified protected fish species. Loss of habitat during the construction phase has been discussed in Section 6.1, 6.2 and 0. Construction of the MOF will result in the loss of some soft bottom habitat and covering of existing reef area that may provide habitat for fish species. Recovery of lost habitat is anticipated following construction, with total clearance conservatively estimated at less than 3 Ha (Table 6-3). As such, the disturbance of intertidal and loss of subtidal habitat is considered to represent a low impact; a short term negative change to fish species that can utilise alternate habitat within the study area and areas subject to regeneration.

The key identified impact to fish species is as a result of noise emissions from the construction and operation of the proposed port. Impacts to marine fauna (including fish species) from noise are discussed in detail in Section 6.7. In summary, the predominant source of noise is piling during construction. Vessel movements during both construction and operation also represent a key noise source. Noise emissions will alter the behavioural patterns of fish species.

Soft start piling procedures will be implemented prior to the commencement of piling operations. Soft start piling provides the opportunity for fish species to vacate an area and avoid the full impact of noise from piling. Underwater noise modelling indicates that any physical damage caused by noise impacts from impact piling will be predominantly limited to fish with swim bladders and cephalopods within 30 m of the piling, while non-swim bladder fish will most likely only suffer behavioural changes (avoidance) (Sonus, 2014).

Behavioural patterns of fish species are expected to change as a result of ad hoc noise emissions during construction and operation. Noise emissions will predominately occur within the study area with affects beyond this area not anticipated. As such, noise emissions from the proposed port are expected to result in a long term change to behavioural patterns of fish species within the study area, with physical damage only caused within a few metres of impact piling and likely to be avoided through the implementation of soft start procedures, considered to be a medium impact.

The introduction of artificial light sources during construction and operation of the proposed port will be required for safe night-time operations and way finding. Artificial light sources can attract a number of fish species, thus affecting the abundance of resources for resident species. As light spill will be minimised through the use of directional lighting (i.e. oriented to a specific area) and the study area does not critical habitat for any fish species, no change to fish distribution or abundance is anticipated and the overall impact is considered to be negligible.

The likelihood of each species occurrence and potential impacts on that species are detailed in Table 6-9.

Table 6-9 Fish species: likelihood of occurrence and potential impacts

| Fish species potentially at the port Site | | | Likelihood of occurrence | Likelihood of significant impacts | | |
|---|--------------------------------|---|---|---|---|--------------------------------|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | | SHARKS | | | |
| Alopias vulpinus | Thresher shark | Reef fish of conservation concern in South Australia (Baker, 2011) | Unlikely for species to be present in area. This species has a very wide distribution, but is most commonly found in offshore waters. It is unlikely that this species occurs in the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris | This species is not listed for protection. Given the large range of this shark species the port site development area is not considered critical habitat and therefore the port will not significantly impact this species or its preferred habitat | N |
| Carcharhinus brachyurus | Whaler shark, Bronze Whaler | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species is most abundant between Albany (WA) and Bass Straight, occasionally entering large bays and inshore areas, but spending the majority of time along continental margins. Despite a lack of records submitted to the ALA and BDBSA, it is considered likely that whaler sharks are occasional visitors in the lower Spencer Gulf. Accordingly, it is possible that this species occurs in the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling | This species is not listed for protection but is of conservation, commercial and recreational interest and occurs throughout the Spencer Gulf. Given the large range of this shark species the port site development area is not considered critical habitat and therefore the port will not significantly impact this species or its preferred habitat | N |
| Carcharhinus obscurus | Dusky Whaler | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species ranges from the surf zone to well offshore, to a depth of 400 m, and is found in all Australian coastal and continental shelf waters. Despite a lack of records submitted to the ALA and BDBSA, it is considered that Whaler Sharks are occasional visitors in the lower Spencer Gulf. Accordingly, it is possible that this species occurs in the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling | This species is not listed for protection but is of conservation, commercial and recreational interest and occurs throughout the Spencer Gulf. Given the large range of this shark species the port site development area is not considered critical habitat and therefore the port will not significantly impact this species or its preferred habitat | N |
| Carcharodon carcharias | Great White Shark | Vulnerable, Migratory, Marine (EPBC); Fisheries Act | Possible. No sighting records of the species have been submitted within 50 km of the area (BDBSA, 2013). The species range is large and may therefore pass in the vicinity of the wharf on route to hunt in areas north or south of Cape Hardy. The species' habits include long migrations by individuals and gatherings around seabird, seal and sea lion colonies. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; vessel impacts | The major fishing grounds for this species tend to centre on seal or sea lion colonies, which are not present within the port site. Therefore while it is possible for the species to be present at the port, given the large global range of White Sharks and the development area is not considered to represent a significant hunting area, the port will not significantly impact this species or its preferred habitat | N |

| Fish species potentially at the port Site | | | Likelihood of occurrence | Likelihood of significant impacts | | | |
|---|---------------------------------|--|--|---|---|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Heterodontus portusjacksoni | Port Jackson shark | N/A | Present. Several individuals sighted during video tow survey, including one juvenile, sighted on predominantly sandy substrate. The species is well known to due to its distinctive appearance, and is common throughout South Australia | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling | This species is not considered threatened or listed for protection. It occurs throughout the Spencer Gulf and southern Australia. Given the large range of this shark species the port development area is not considered critical habitat and therefore the port will not significantly impact this species or its preferred habitat. | N | |
| Isurus oxyrinchus | Shortfin Mako | Migratory, Marine (EPBC) Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species ranges from close inshore in surface waters to 150 m depth, though generally restricted to waters warmer than 16 °C (Australian Museum, 2013). Fisheries catch records logged on the ALA indicate the presence of the Shortfin Mako in waters offshore of South Australia in 2001-2002 and earlier records in the 1980s from the SA Museum for inside the Spencer Gulf (ALA, 2013). It is therefore possible that this species could pass by the study area, although due to the paucity of records for the species in SA any individuals in the Cape Hardy area would be considered vagrants. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling | This species is not listed for protection but is of conservation, commercial and recreational interest and occurs throughout the Spencer Gulf. Given the large range of this shark species the port development area is not considered critical habitat and therefore the port will not significantly impact this species or its preferred habitat. | N | |
| Lamna nasus | Porbeagle, Mackerel Shark | Migratory, Marine (EPBC) | Possible . No records of the species have been submitted to the BDBSA within 50 km of the area (BDBSA, 2013). However this species has a large range and may therefore pass in the vicinity of the wharf to forage in the Cape Hardy area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling | Given the large range of this shark species the port development area is not considered critical habitat and therefore the port will not significantly impact this species or its preferred habitat. | N | |
| Mustelus antarcticus | Gummy shark | N/A | Possible. This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. It is possible that suitable habitat exists in the project area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling | Given the large range of this shark species the port development area is not considered critical habitat and therefore is will not significantly impact this species or its preferred habitat. | N | |

| Fish species potentially at the port Site | | | Likelihood of occurrence Likelihood of significant impacts | | | |
|--|------------------------------|---|---|--|--|--------------------------------|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| Orectolobus halei, O. maculatus, Sutorectus tentaculatus | Wobbegong shark (various) | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species is not listed for protection but is regarded as a species of conservation concern by Reef Watch. It is possible that suitable habitat exists in the project area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling | Given the large range of this shark species the port development area is not considered critical habitat and therefore the port will not significantly impact this species or its preferred habitat. | N |
| Urolophus orarius | Coastal Stingaree | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species is endemic to inshore areas of South Australia, usually inhabiting sandy substrates between 14-50 m depth. The range of this species is between Beachport and Ceduna, in South Australia. The majority of records for this species come from the 20 to 50 m depth range as almost all records of the species are from prawn trawl by-catch, which is restricted to depths of more than 10 m (SPRAT, 2014) The species may utilise shallower habitats as well, however there has been no sampling in shallow waters. Suitable habitat exists in the study area, and while no records have been submitted to the ALA in Spencer Gulf, it is possible that this species is present or occasionally visits the area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling | The species is classified as Endangered by the ICUN red list but was rejected for nomination of listing under the EPBC Act in 2011 (SPRAT, 2014). The species is not listed for protection but is of conservation concern It is likely that this species occurs in the area as suitable habitat is present. A small amount of potential habitat for this ray species will be affected by the development however given the species large range the port development area is not considered critical habitat and therefore the port will not significantly impact this species or its preferred habitat. | N |
| | | | FINFISH | | | |
| Achoerodus gouldii | Western blue groper | Fisheries Act; Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. Individuals of this species often form residency in an area, usually preferring complex reef habitats. Juveniles of these fish are female and appear yellow-green in colour. This colour changes to a deep blue later in life after they become male. This species is iconic, growing to 1.75 m in length and often surviving for up to 70 years. It is this species' long life-span and slow growth combined with its curiosity towards humans that makes this species sensitive to fishing pressure. This species is regarded as a species of conservation concern by Reef Watch SA. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is protected at all times within Spencer Gulf under the Fisheries Act (SA) (PIRSA, 2014) as it is of conservation and recreation interest. As it occurs throughout remote areas of southern Australia it is possible that this species may occur in the wider area, but it is considered that only limited suitable reef habitat for this species occurs in the project area. Due to the small footprint of the MOF etc. no significant amount of available habitat will be impacted by this development. As the presence of the port will prevent recreational fishing in the port area the port infrastructure may overtime provide a suitable refuge area for this species that prefers complex habitats. The development will not significantly impact this species or its preferred habitat. | N |

| Fish species pot | entially at the po | rt Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|--------------------------|--------------------------------------|--|---|--|--|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | | | | |
| Aldrichetta forsteri | Yelloweye mullet | N/A | Possible. This fish species is widespread around southern Australia. This species prefers sheltered sand and seagrass habitat of < 20 m depth (. Sheltered sand and seagrass suitable for this species is widespread across the Study Site. This species is likely to occur in the area, as several sightings have been logged with the ALA near populated areas north and south of the site (ALA, 2013)^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. It is likely that this species occurs in the area as suitable habitat is present. This species is widespread. A small amount of potential habitat for this fish will be affected by the development. The development will not significantly impact this species or its available habitat. | N | |
| Argyrosomus japonicus | Mulloway | N/A | Possible. This species' distribution includes the coasts of S.E. Asia, India, Africa, Madagascar and the southern coastline of Australia from Shark Bay (WA) to Brisbane (QLD). Fisheries targeting these fish typically operate in major estuaries in SA. It is possible that this species occurs in the project area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. While it is possible that this species could occur in the study area, the area does not represent critical habitat for this species. This development will not significantly impact on this species. | N | |
| Arripis georgianus | Tommy ruff, Australian herring | N/A | Likely. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom, seagrass and reefs (Bryars, 2003; Baker, 2004). This small fish species frequently schools in the open water of sheltered bays around southern Australia. It is likely that this species uses suitable habitat within the study area. This species migrates to the south-western corner of Australia to spawn in February and March^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. The study area does not represent critical habitat for this species. This development will not significantly impact on this species. | N | |
| Arripis truttaceus | Australian salmon (western) | N/A | Likely. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom, seagrass and reefs (Bryars, 2003; Baker, 2004). This large fish species is common on southern Australian coasts, often schooling in shallow water, leading to popularity amongst anglers. This species migrates westward to spawn after 5-6 years of growth. It is likely that this species passes by Cape Hardy, but the area does not represent any particular significance to this species' life cycle^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil | This species is not listed for protection but is of commercial and recreational interest and occurs throughout southern Australian waters. The study area does not represent critical habitat for this species. This development will not significantly impact on this species. | N | |

| Fish species pote | entially at the po | rt Site | Likelihood of occurrence | Likelihood of significant impacts | | |
|-------------------------------|---------------------|---|---|--|--|--------------------------------|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | | | spill; pollution | | |
| Cheliodactylus spectabilis | Banded morwong | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This fish occurs in Australia between NSW and SA, mostly on high-energy shallow coastal reefs. This species reaches a large size at maturity, and can reach 95 years of age. It is likely that their approachable behaviour leads them to be sensitive to exploitation by fishing activities. It is possible that this species occurs in the reef habitat present at the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of conservation concern, commercial and recreational interest and occurs throughout southern Australian waters. The study area does not represent critical habitat for this species. This development will not significantly impact on this species. | N |
| Dactylophora nigricans | Dusky morwong | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This large fish prefers seagrass and reef habitat in shallow waters. This species is common, sighted between NSW and WA in southern waters. It is likely that this species utilises habitat present in the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of conservation concern (Baker, 2011), recreational interest and occurs throughout southern Australian waters. The study area does not represent critical habitat for this species. This development will not significantly impact on this species. | N |
| Dinolestes lewini | Long-finned Pike | N/A | Possible. This species is common on reefs around southern australia and is likely to occur in the study area^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of recreational interest and occurs throughout the Spencer Gulf (Baker, 2011). While it is likely that this species occurs in the study area, the habitat present is unlikely to represent a significant area for the species. This development will not significantly impact on this species. | N |
| Enoplosus armatus | Old wife | N/A | Present. One individual was observed over bare silt at over 23 m during the video tow survey. The species depth range is 0-100 m and commonly found as pairs or large individuals in exposed areas but as shoals of smaller fish amongst seagrass or sheltered jetties (Edgar, 1997). | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil | This species is not listed for protection but is of recreational diver interest and occurs throughout the Spencer Gulf (Baker, 2011). While this species is known to occur in the study area, the habitat present is unlikely to represent a significant area for the species and given the species ability to utilise man-made structures for shelter of smaller fish, this species will not be significantly impacted by this development. | N |

| Fish species pote | entially at the por | t Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|-------------------------------|-----------------------------------|---|--|--|--|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | spill; pollution | | | |
| Hyporhamphus melanochir | Southern Sea garfish | N/A | Likely. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom and seagrass (Bryars, 2003; Baker, 2004). This species is most commonly found swimming near the surface in areas of seagrass meadow. It is likely that this species occurs in the project area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf (Baker, 2011). The proposed development does not represent a significant portion of this species' range, and thus this species will not be significantly impacted by this development. | N | |
| Hypoplectrode s nigroruber | Black-banded seaperch | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible . This species generally inhabits reefs, but has also been recorded on artificial reefs, shipwrecks and jetties. Knowledge of this species is limited, though these fish are widely distributed between WA and NSW. It is possible that this species occurs in the project area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of conservation interest and occurs throughout the Spencer Gulf (Baker, 2011). The proposed development does not represent a significant portion of this species' range, and thus this species will not be significantly impacted by this development. | N | |
| Kathetostoma laeve | Eastern or common stargazer | N/A | Likely. The species has been regularly reported along the Tumby Bay coastline in areas of sandy bottom and jetties (Baker, 2004). The species is common across southern Australia in waters 0-60 m, it is an ambush predator that prefers sandy or shelly bottoms but is also found around manmade structures. It is likely that this species is present in the study area [^] . | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of some commercial and recreational interest and occurs throughout the Spencer Gulf (Baker, 2011). As the species is already commonly sighted around jetties, the proposed development will potentially benefit this species and thus this species will not be significantly negatively impacted by this development. | N | |
| Kyphosus sydneyanus | Silver drummer | N/A | Possible. This species is common across southern Australia, especially nearby rocky reef habitat, including reef in urbanised areas. It is likely that this species is present in the study area [^] . | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf (Baker, 2011). The proposed development does not represent a significant portion of this species' range, and thus this species will not be significantly impacted by this development. | N | |

| Fish species pote | entially at the port | Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|-------------------------------|--|---|--|--|--|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | spill; pollution | | | |
| Nelusetta ayraudi | Ocean jacket | N/A | Present. This species was sighted on a number of occasions during the video tow survey over bare sand; it is widespread in the Southern ocean continental shelf waters between central WA and Brisbane QLD, including SA gulf waters. This species prefers deep coastal waters as adults, but juveniles utilise bays and estuaries. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of some commercial and recreational interest and occurs throughout the Spencer Gulf (Baker, 2011). While it is possible that this species could occur in the study area, the area is does not represent a large area or critical habitat for the species. This development will not significantly impact on this species. | N | |
| Nemadactylus valenciennesi | Southern blue morwong/ Queen snapper | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species occurs in complex reef habitat around southern Australia [^] . It is likely that this species relies on shallower waters and seagrass habitat during juvenile stages. It is possible that this species occurs in the project area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest, and is regarded as a species of conservation concern by Reef Watch SA (Baker, 2011). The project area does not contain a significant area utilised by this species. This development will not significantly impact on this species. | N | |
| Notolabrus fucicola | Purple wrasse | Reef fish of conservation concern in South Australia (Baker, 2011) | Unlikely This species is common in south-east Australia, but rarely sighted in the waters of Spencer Gulf, which is considered the Western-most edge of its range (Gomon <i>et al.</i> , 2008). This species prefers low rocky reefs, which have limited coverage in the study area. It is possible that this species occurs in the study area but given its rarity in Spencer Gulf its presence is considered unlikely. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is regarded as a species of conservation concern (Baker, 2011). The project area does not contain an area critical to the life cycle of this species, or likely to support a significant population. This development will not significantly impact on this species. | N | |
| Notolabrus parilus | Brown/ orange -spotted wrasse | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species occurs frequently in reef habitat around southern Australia^. There is some indication that juveniles of this species utilise seagrass habitat. It is possible that this species utilises habitat within the project area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil | This species is not listed for protection but is of conservation concern (Baker, 2011). The project area does not contain an area crucial to the life cycle of this species, or likely to support a significant population. The development will not significantly impact on species. | N | |

| Fish species pote | entially at the port | Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|-----------------------------|------------------------|---|---|--|---|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | spill; pollution | | | |
| Notolabrus tetricus | Blue-throated wrasse | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species occurs frequently in reef habitat around southern Australia^. There is some indication that juveniles of this species utilise seagrass habitat. It is possible that this species utilises habitat within the project area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest, and is regarded as a species of conservation concern (Baker, 2011). The project area does not contain an area critical to the life cycle of this species, or likely to support a significant population. This development will not significantly impact on this species. | N | |
| Othos dentex | Harlequin fish | Reef fish of conservation concern in South Australia (Baker, 2011) | Unlikely. This species inhabits reef habitat around South Australia, particularly reefs with higher habitat complexity, and remote areas with lower historic fishing pressure. It is possible that this species is present in the study area, but it is unlikely that the habitat present supports a high population^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and is regarded as a species of conservation concern by Reef Watch SA, mostly due to its sensitivity to fishing pressure (Baker, 2011). The study area does not represent a significant area of suitable habitat for this species. This development will not significantly impact on this species. | N | |
| Paraplesiops meleagris | Southern blue devil | Reef fish of conservation concern in South Australia (Baker, 2011) | Possible. This species is iconic and rare in South Australia, though distributed between the Abolhos (WA) and Port Philip Bay (VIC). These fish are territorial and highly site-attached. Individuals and pairs inhabit caves, ledges and crevices in rocky reef, between 5 and 45 m depth. It is possible that suitable habitat exists for the Blue Devil at the rocky capes in the study area^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of recreational interest and is regarded as a species of conservation concern 'in peril' by Reef Watch SA (Baker, 2011). The study area does not represent a significant area of suitable habitat for this species. This development will not significantly impact on this species. | N | |
| Paristiopterus gallipavo | Brown-spotted boarfish | Reef fish of conservation concern in South Australia (Baker, 2011) | Unlikely. This species prefers reef habitat between 55-200 m depth^. This species occurs commonly in reef habitat around southern Australia between SA gulf waters and the central coast of WA. With water depth shallower than 30 m and no significant reef area in the study area it is considered unlikely the species would frequent the area. | Entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest, and is regarded as a species of conservation concern (Baker, 2011). The project area does not contain an area critical to the life cycle of this species, or likely to support a significant population. This development will not significantly impact on this species. | N | |

| Fish species pote | entially at the po | rt Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|--|-----------------------|---|--|--|--|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Pagrus auratus / Chrysophrys auratus | Snapper | N/A | Likely for the species to be present. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom, seagrass and reefs (Bryars, 2003; Baker, 2004). This species occurs widely across southern Australia, generally around reefs and ledges, or other distinct underwater structures such as drop-offs, artificial reefs and wrecks. Snapper can live to 35 years of age, and are a prized fish for recreational and commercial anglers. Key breeding areas for this fish have been identified, and are subject to defined closures from fishing. It is possible that Snapper are present in the study area, though the study area does not represent a key area for breeding or the life cycle of this species. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf (Baker, 2011). While it is possible that this species is present in the study area, it is not a critical area to the species. This development will not significantly impact on this species. | N | |
| Pegasus Iancifer | Sculptured seamoth | N/A | Likely for the species to be present. The species has been regularly reported along the Tumby Bay coastline in areas of sandy bottom and jetties (Baker, 2004). The species is common across southern Australia in waters 0-55m, it is bottom feeder which searches for small infauna crustaceans and polychaetes, it prefers sandy or muddy bottoms but is also found around manmade structures such as jetties and near seagrass. It is likely that this species is present in the study area*. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of (limited) commercial and recreational interest and occurs throughout the Spencer Gulf (Baker, 2011). While it is possible that this species is present in the study area, it is not a critical area to the species. This development will not significantly impact on this species^. | N | |
| Pentaceropsis recurvirostris | Longsnout boarfish | Reef fish of conservation concern in South Australia (Baker, 2011) | Present. Boarfish were sighted during the video tow survey in an area of sparse seagrass. This species occurs from Botany Bay (NSW) to Rottnest Island (WA), including Tasmania. The Longsnout Boarfish occurs frequently on rocky reef habitat around South Australia, to a depth of 260 m. Suitable habitat exists in the project area but it is unlikely that the habitat present supports a high population ^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of recreational interest and is regarded as a species of conservation concern 'in peril' by Reef Watch SA (Baker, 2011). The study area does not represent a significant area of suitable habitat for this species. This development will not significantly impact on this species. | N | |
| Platycephalus spp. | Flathead (various) | N/A | Likely for the species to be present. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom and seagrass (Bryars, 2003; Baker, 2004). This species prefers to inhabit sandy bottoms and shallow sandy ledges. It is likely that flathead are present at the site, though it is not likely to be an area of high significance for the species^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. Flathead species are a common target for recreational anglers in South Australia.Cape Hardy is not likely to be a key area for these species, which occurs widely throughout southern | N | |

| Fish species pot | entially at the por | t Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|--|------------------------|--|---|--|--|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | impacts within 10m of piling; oil spill; pollution | Australia. This development will not significantly impact on this species. | | |
| Pseudocaranx georgianus (Previously known as P. dentex but now distinguished from the tropical species) | Silver trevally | N/A | Likely for species to be present in area. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom, seagrass and reefs (Bryars, 2003; Baker, 2004). This species is thought to be distributed between NSW and southern WA. Generally, this species prefers to school in shallow shelf waters. It is possible that suitable habitat occurs in the study area, and therefore that this species is present in the area^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. While it is possible that this species utilises habitat in the study area, the area is not a known spawning location for this species. This development will not significantly impact on this species. | N | |
| Scorpis aequipinnis | Sea sweep | N/A | Possible. This species is common in southern Australian waters, especially in reef habitat and shallow coastal waters to 25m (Gomon et al., 1994). This species often associates with artificial structures including jetties. Due to the small about of suitable habitat it is likely that this species will be present in the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. It is likely that this species will be present in the study area, but considering it feeds well off the bottom almost exclusively on plankton and associates with artificial structures (Gomon et al., 1994), this development will not significantly impact on this species. | N | |
| Seriola lalandi | Yellowtail kingfish | N/A | Possible. This species is circumglobal, including the coastal waters of southern Australia. This species is fast moving and pelagic, but is native to Spencer Gulf and seen in coastal areas at times. The species is farmed commercially in Spencer Gulf with aquaculture leases for both land based breeding and seacage growout. It is possible that wild fish of this species may be found in the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. Due to the large range of this species, the study area does not represent critical habitat for this species. This development will not significantly impact on this species. | N | |
| Sillaginodes punctatus | King George whiting | N/A | Likely. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom, seagrass and reefs (Bryars, 2003; Baker, 2004). This species is not listed for protection but is of commercial and recreational interest and occurs | Loss of habitat through clearance or exclusion; entanglement in | Impacts to suitable habitat for this species will be restricted to the area of wharf and MOF construction, and will not greatly affect the ability for this species to inhabit the wider area. The development site does not | N | |

| Fish species pote | entially at the po | ort Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|--|------------------------------------|--|---|--|---|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | throughout the Spencer Gulf. The King George Whiting is an iconic South Australian table fish, targeted by commercial and recreational anglers. This species prefers sandy substrates, especially around seagrass beds or low profile reef habitat. Suitable habitat is present at Cape Hardy, so it is likely that King George whiting are found at the site^. | marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | represent an area critical to the recruitment or ongoing survival of this species. This development will not significantly impact on this species. | | |
| Sardinops sagax, S. neopilchardus. | Pilchard, Australian sardine | N/A | Possible. This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. This species is known to inhabit lower Spencer Gulf, with the majority of fisheries harvest occurring in the lower Spencer Gulf area (Shanks, 2005). Generally, fishing occurs in deeper waters at the mouth of the gulf, but it is possible that schools of pilchards may enter the study area. The area is not considered a critical area for the pilchard population ^A . | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution, disturbance to recruitment; disturbance to prey species (e.g. Anchovy) | The study area is not a known fishing ground for pilchards, although it is possible that schools pass through the area, as suitable habitat exists. It is unlikely that the development site is an important area for the species. This development will not significantly impact on this species. | N | |
| Sphyraena novaehollandia e | Snook | N/A | Likely for the species to be present. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom and seagrass (Bryars, 2003; Baker, 2004). This species is widely distributed across southern Australia, including SA gulf waters and offshore to the continental shelf margin. It is likely that this species occurs in the project area^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. While it is possible that this species could occur in the study area, the area is of no significance to the species. This development will not significantly impact on this species. | N | |
| Sillago schomburgkii | Yellowfin whiting | N/A | Likely. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom, seagrass and reefs (Bryars, 2003). This species occurs between central WA and the Fleurieu Peninsula in SA, including SA gulf waters. The preferred habitat for this species includes inshore sandbanks, bars, spits and estuaries. It is likely that this species utilises habitat at the study | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise | This species is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. While it is possible that this species could occur in the study area, the area is of no significance to the species. This development will not significantly impact on this species. | N | |

| Fish species potentially at the port Site | | Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|---|---|--|---|--|---|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | area^. | impacts within 10m of piling; oil spill; pollution | | | |
| Tilodon sexfasciatus | Moonlighter, six-banded coral fish, six- banded sweep, butterfish | N/A | Likely. The species has been regularly reported along the Tumby Bay coastline in areas of sandy bottom and jetties (Baker, 2004). The species is common across southern Australia. It is likely that this species is present in the study area*. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of recreational interest and occurs throughout the Spencer Gulf. While it is likely that this species occurs in the study area, the size of the development area is not a significant portion of the species range. This development will not significantly impact on this species. | N | |
| Upeneichthys vlamingii | Red mullet, blue-spotted goatfish, southern goatfish | N/A | Likely. The species has been regularly reported along the Tumby Bay to Port Neill coastline in areas of sandy bottom and seagrass (Bryars, 2003; Baker, 2004). The species is widespread and very common across southern Australia including Tasmania – inhabiting sheltered coastal waters, bays, estuaries usually on soft sandy or loose gravel near reefs but is also found in and around seagrass meadows. It is likely that this species is present in the study area^. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This species is not listed for protection but is of recreational and some commercial interest and occurs throughout the Spencer Gulf. While it is likely that this species occurs in the study area, the size of the development area is not a significant portion of the species range. This development will not significantly impact on this species. | N | |
| Vincentia conspersa | Southern cardinalfish, gobbleguts | N/A | Likely. The species has been regularly reported along the Tumby Bay coastline in areas of sandy bottom and jetties (Baker, 2004). The species is common across South Australia and parts of Victoria and Tasmania, it is nocturnal carnivore which feeds on both zooplankton and benthic invertebrates, it prefers reefs hiding in caves or under overhangs by day but is also found around manmade structures such as jetties and breakwaters. It is likely that this species is present in the study area*. | Loss of habitat through clearance, shading, anchoring impacts, epiphyte blooms or smothering by ore spill; oil spills; pollution; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | The species is not protected under any current National or State legislation. They are not species targeted by fisherman but are of some interest to recreational divers. There will be a small area of benthic habitat removed due to the construction of the wharf and MOF. However given that this species distribution covers all of South Australia and its habitat preference is for reef or hard structures, the construction of the MOF and jetty will provide additional habitat for this species. This development will not significantly impact on this species. | N | |
| FAMILY | Wrasse | N/A | Likely for the species from this Family to be present. "Wrasse" have | Loss of habitat | With the exception of A. gouldii, discussed above, this | N | |

| Fish species pot | entially at the port | Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|-----------------------------|------------------------------------|--|--|--|---|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| Labridae | (various) | | been regularly reported along the Tumby Bay to Port Neill coastline in areas of reef (Bryars, 2003; Baker, 2004). This family is also widespread across a variety of inshore habitats in southern waters. It is highly likely that species from this family occur in the study area^. | through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | fish family is not listed for protection but is of some commercial and recreational interest. Occurring throughout the Spencer Gulf and much of southern Australia the proposed development does not represent a significant portion of the range of this family, and thus this family will not be significantly impacted by this development. | | |
| FAMILY Sparidae | Bream | Fisheries Act | Possible. This Family occurs between central WA and Qld, including SA gulf waters. The preferred habitat for these fish includes sheltered inshore sandbanks, bars, spits and estuaries, over seagrass to open water deep reefs (Edgar, 1997). It is therefore likely that fish in this family utilises habitat in the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | This Family of fish are keenly sort by anglers both commercial and recreational. There will be a small area of seagrass and benthic habitat removed due to the construction of the wharf and MOF. Given that this family's distribution covers all of Southern Australia and parts of international waters, the small area of clearance associated with the development is not considered to represent a significant area of crucial habitat for this species. This family will not be significantly impacted by this development. | N | |
| FAMILY Aracanidae | Temperate boxfish or Cowfish | N/A | Present. Observed on a number of occasions throughout the study area in areas of bare sand or silt or patchy to dense seagrass or amongst sponge groups. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | The Family are not protected under any current National or State legislation. They are not species targeted by fisherman but are of interest to recreational divers. There will be a small area of seagrass and benthic habitat removed due to the construction of the wharf and MOF. Given that this family's distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species This family will not be significantly impacted by this development. | N | |
| FAMILY Monacanthida e | Leather- jacket (various) | N/A | Present. Several species of Leatherjacket were observed during the video tow survey, including the Mosaic leatherjacket (<i>Eubalichthys mosaicus</i>). This family is present in all major oceans globally, including tropical and temperate waters and a range of coastal | Loss of habitat through clearance or exclusion; entanglement in | This fish family is not listed for protection but is of commercial and recreational interest and occurs throughout the Spencer Gulf. The proposed development does not represent a significant portion of | N | |



| Fish species pote | entially at the port | Site | Likelihood of occurrence | Likelihood of significant impacts | | |
|------------------------------|--------------------------------|--|--|--|---|--------------------------------|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) |
| | | | habitat types. It is highly likely that fish species in this family utilise each of the habitat types identified in the study area. Juveniles of this family utilise seagrass beds and estuaries. Reefs of all description support this family, along with shipwrecks and artificial structures. It is likely that this family is present in the study area. | marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | the range of this family, and thus this family will not be significantly impacted by this development. | |
| FAMILY Tetraodontida e | Pufferfish and toadfish | N/A | Possible. This Family occurs between central WA and Qld, including SA gulf waters. The preferred habitat for these fish includes sheltered inshore sandbanks, bars, spits and estuaries, over seagrass to open water deep reefs and wrecks (Edgar, 1997). It is therefore likely that fish in this family utilises habitat in the study area. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | The Family are not protected under any current National or State legislation. They are not species targeted by fisherman as many species are highly poisonous but they are of interest to recreational divers due to the fish's highly decorative and inquisitive nature. There will be a small area of seagrass and benthic habitat removed due to the construction of the wharf and MOF. Given that this family's distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. The construction of the MOF and jetty will also provide additional habitat niches for members of this Family. This family will not be significantly impacted by this development. | N |
| | | | SYNGNATHIDS PIPEFISHES | | | |
| Acentronura australe | Southern pygmy pipehorse | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N |
| Campichthys tryoni | Tryon's pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This | N |

| Fish species pote | entially at the port | Site | Likelihood of occurrence | Likelihood of significant impacts | | | |
|-------------------------------|---|--|---|---|---|--------------------------------|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | underwater noise impacts within 10m of piling; oil spill; pollution | development will not significantly impact on this species. | | |
| Filicampus tigris | Tiger pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species is rarely found in Spencer Gulf with less than 30 speciemens recorded. Its preferred habitat is known to be sand and mud (Inshore Fish Group, 2008), and given the amount of suitable substrate at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of sparse seagrass on sandy sediment lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Heraldia nocturna | Upside-down pipefish, eastern upside-down Pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution. | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Hippocampus breviceps | Short-head seahorse, short-snouted seahorse | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Histiogamphel us cristatus | Rhino pipefish, Macleay's, crested pipefish, ring- back, pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This | N | |

| Fish species potentially at the port Site | | | Likelihood of occurrence | Likelihood of significant impacts | | | | |
|---|---|--|--|--|--|--------------------------------|--|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | | |
| | | | | underwater noise impacts within 10m of piling; oil spill; pollution | development will not significantly impact on this species. | | | |
| Hypselognathu s rostratus | Knifesnout pipefish, knife- snouted pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance, shading, anchoring impacts, epiphyte blooms or smothering by ore spill, chemical spill or runoff impacts. | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |
| Kaupus costatus | Deepbody pipefish, deep- bodied pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |
| Leptoichthys fistularius | Brushtail pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |
| Lissocampus caudalis | Australian smooth pipefish, smooth pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this | N | | |

| Fish species potentially at the port Site | | | Likelihood of occurrence | Likelihood of significant impacts | | | |
|---|--|---------------------------------|--|--|--|--------------------------------|--|
| Species name | | | egislative protection PBC, or Fisheries Act (including MOF and wharf) | | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | | impacts within 10m of piling; oil spill; pollution | species. | | |
| Lissocampus runa | Javelin pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Maroubra perserrata | Sawtooth pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Notiocampus ruber | Red pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Phycodurus eques | hycodurus Leafy Marine (EPBC); Present. This species was observed during the marine video tow Lo | | | | There will be a small area of benthic vegetation lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. Furthermore the species has a preference for | N | |

| Fish species potentially at the port Site | | | Likelihood of occurrence | Likelihood of significant impacts | | | |
|---|---|---------------------------------|--|--|--|--------------------------------|--|
| Species name | Species name Common Legislative protection EPBC, or Fisheries Act | | Marine environment (including MOF and wharf) | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | |
| | | | specimen was sighted within seagrass beds during the video tow survey and there is coverage of suitable macro-algal habitat at Cape Hardy it is likely the species would be widespread across the site. | impacts within 10m of piling; oil spill; pollution | protected reef habitat including Rapid Bay Jetty where divers regularly come across these seadragons. Therefore the completed marine infrastructure for this development would potentially increase the amount of suitable habitat at the site. This development will not significantly impact on this species. | | |
| Phyllopteryx taeniolatus | Common seadragon, weedy seadragon | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Pugnaso curtirostris | Pugnose pipefish, pug- nosed pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution. | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Solegnathus robustus | Robust pipehorse, robust spiny pipehorse | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | |
| Stigmatopora argus | Spotted pipefish, gulf pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the | Loss of habitat through clearance or exclusion; | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the | N | |

| Fish species potentially at the port Site | | | Likelihood of occurrence | Likelihood of significant impacts | | | | |
|---|--|---------------------------------|--|--|--|--------------------------------|--|--|
| Species name | Common Legislative protection name EPBC, or Fisheries Act | | | | Likelihood of Impact | Significant Impact (Y/N) | | |
| | | | species could inhabit the site. | entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | | | |
| Stigmatopora nigra | Widebody pipefish, wide- bodied pipefish, Black pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |
| Stipecampus cristatus | Ringback pipefish, ring- backed pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |
| Urocampus carinirostris | Hairy pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |
| Vanacampus margaritifer | Mother-of- pearl pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the | Loss of habitat through clearance or exclusion; | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the | N | | |

| Fish species potentially at the port Site | | | Likelihood of occurrence | Likelihood of significant impacts | | | | |
|---|--|---|--|--|--|--------------------------------|--|--|
| Species name | Common name | Legislative protection EPBC, or Fisheries Act (including MOF and wharf) | | Potential Impacts | Likelihood of Impact | Significant Impact (Y/N) | | |
| | | | species could inhabit the site. | entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution. | small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | | | |
| Vanacampus phillipi | Port Phillip pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |
| Vanacampus poecilolaemus | Longsnout pipefish, Australian long-snout pipefish, long- snouted pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |
| Vanacampus vercoi | Verco's pipefish | Marine (EPBC); Fisheries Act | Possible. No records for this species have been submitted to the BDBSA (BDBSA, 2012). The species in known to inhabit seagrass and given the coverage of seagrass at Cape Hardy it is possible that the species could inhabit the site. | Loss of habitat through clearance or exclusion; entanglement in marine debris; underwater noise impacts within 10m of piling; oil spill; pollution | There will be a small area of seagrass lost due to the construction of the wharf and MOF. Given that this species' distribution covers all of Southern Australia, the small area of clearance is not considered to represent a significant area of crucial habitat for this species. This development will not significantly impact on this species. | N | | |



6.6.2.2 Risks

Construction and operation of the proposed port will result in the generation of a number of waste streams that may enter the marine environment. Marine debris can significantly affect fish species as a result of pollution, injury through collision, entanglement or ingestion of non-biological products.

Waste management and handling procedures developed as part of the CEMP and OEMP will control waste streams with the overarching aim of no waste products entering the marine environment. Despite the implementation of control measures, it is considered almost certain that some form of waste / debris will enter the marine environment during construction and operation of the port. Although the port site contains areas of seagrass, it is not a known key habitat or breeding area for any fish species. The consequences of debris entering the marine environment are therefore considered to be minimal; insignificant to the overall viability of fish species. As such, the overall risk is considered to be low.

Fish species are also at risk from the unintended discharge of pollution in the form of site run-off, wastewater, hydrocarbons or chemicals. With any marine activity there is potential for waste or spills to enter the marine environment and affect plankton, benthic fauna and habitat forming benthic flora (e.g. seagrass), reducing habitat suitability for fish species.

The implementation of design measures (refer Section 4) for the control of surface and waste water at the port site will limit the majority of hazardous pollutants from entering the marine environment. There remains a risk that a major spill of oil or other chemicals may occur as a result of vessel failure or accident. Based on the experiences of other operating ports, the likelihood of this event occurring is considered to be rare. As previously outlined, the port site does not represent a critical habitat or isolated breeding grounds for fish species. The consequences of a major spill event are considered to be minor, due to the low level of dependence of fish species on the study area, resulting in a short term decrease in the local abundance of fauna. As such, the overall risk of a major spill event affecting fish species is considered to be low.

6.6.3 Conclusions

Due to the highly mobile nature of fish a wide range of species may be present in the Cape Hardy area at different times or for limited periods, however there are a number of species that have a higher likelihood of being observed in the waters around Cape Hardy. Leafy seadragon, ornate cowfish, Port Jackson shark, magpie perch, leatherjackets and toadfish were all sighted during the video survey of the study area. There is also the potential for a large number of other fish species to be present within the seagrass, reef and soft bottom areas of the site. EPBC listed Marine species are protected past the 3 nautical mile limit in Commonwealth Waters. Spencer Gulf lies completely within State waters and therefore is not subject to this protection status. Syngnathids are also protected under the Fisheries Management Act 2007 along with the great white shark as are the blue groper at all times within Spencer Gulf.

As with marine megafauna, piling and underwater noise is considered to represent the greatest impact to fish species, however these impacts are of short duration and through the implementation of management procedures (such as soft start piling and observation zones) impacts are expected to be medium.



6.7 Noise and Vibration

While it is common knowledge that cetaceans utilise sounds for navigation and communication, fish, invertebrates, and even planktonic larval stages of many marine biota also use sound for a range of important functions. Marine fauna use sounds as a part of normal behaviour when foraging, to detect prey, predators or other hazards, for navigation, communication or when locating mates. Worldwide, studies into the effects of noise on marine fauna have been undertaken since the 1950s, ranging between the effects of high-intensity and low-level noise, and a diverse range of biota from cetaceans to larvae.

Early research observed the effects of high-intensity pulsed and continuous sound from construction, explosives, sonar, shipping, and geological survey on the behaviour and physiology of marine mammals, fishes, marine turtles and other species (Richardson *et al.*, 1986; Richardson *et al.*, 1990; Richardson and Würsig 1997). High-intensity sound can cause physical trauma to sensitive species, stress responses, altered behaviour, and disorientation which has been linked to numerous stranding events (Goold and Fish, 1998; Fernandez *et al.*, 2005; Goldbogen *et al.*, 2013). Fish can also be injured by high-intensity sounds at close range, though this is dependent on the physiology of the species, sensitivity and the sound profile (Wardle *et al.*, 2001). On long time-frames however, whales have not been seen to change migratory habits or stop visiting key areas after being disturbed by noisy exploration or construction (DEWHA, 2008).

A wide range of human activities emit noise and vibration, including recreational boating, commercial shipping, coastal construction, and seismic surveys, which can be disruptive to marine biota. The impact of noise and vibration on marine biota depends on the type of noise or vibration, frequency, wavelength and duration, and the mobility and physiology of the organism. The organisms considered most sensitive to disturbance from noise and vibration are generally free-swimming, especially those with more developed auditory structures, including marine mammals and some fish (Myrberg, 1990). Fish with swim bladders are also potentially susceptible regardless of their hearing capability as high intensity sound can physically damage their swim bladder resulting in death. The following assessment of impacts and risks to marine biota from noise and vibration is undertaken based on modelling of the propagation of sound in the marine environment during the construction phase, and the proposed level of shipping activity at the port.

Noise and vibration levels will depart from normal conditions as a result of construction and operations at the port. The construction process will involve pile driving, which poses a risk of injury or disturbance to marine mammals and fish in the immediate area. The elevated ambient or background noise and vibration during standard shipping operations at the site poses a risk of masking the sounds required for normal behaviour of large marine fauna including marine mammals or other smaller marine biota including fish and invertebrates (Slabbekoorn *et al.*, 2010). The full effects of masking sounds is not understood as most studies of hearing loss or masking are based on small sample sizes or individual animals under experimental conditions, and do not account for variations in hearing range between individuals or reflect animals natural stress responses (Southall, 2005).

Habituation of marine animals to areas of high vessel traffic and noise has been shown to occur readily around the globe. Several resident populations of marine mammals, birds and fish species are observed at existing large ports in Australia e.g. Port Adelaide Port River (SA) dolphins and bird rookeries; Port Phillip Bay (VIC) dolphin and seabird populations; Swan River (WA) bottlenose dolphins. Therefore, while marine animals are known to be highly sensitive to underwater noise impacts and display either behavioural or physiological changes due to these disturbances, impacts from commercial vessel and port operations are not as clearly understood, with cohabitation of the marine environment a demonstrated outcome.



6.7.1 Existing Environment and Key Findings

Ambient underwater noise measurements within the marine study area were not taken as part of the investigations for the proposed port. Ambient noise levels differ based on sea conditions, water depths, and natural and main made noise sources. As previously outlined, the marine study area is a shallow, moderate energy coastal environment meaning that ocean noise (wind and wave noise) will be higher than in deep water.

No significant man made noise sources were identified in the vicinity of the proposed port. The coastal townships of Port Neill and Tumby Bay are respectively located north and south of the port. Low levels of recreational and commercial vessels utilise these townships and would generate low levels of ambient noise. East of the port site is the shipping channel to support the movement of larger commercial vessels to Whyalla and Port Augusta. Ship movements within the channel would also contribute to low levels of ambient noise.

When considering the impact that noise has on the marine fauna, consideration is given to a range of factors such as the sensitivity of the species to noise, the distance between the noise source and fauna, and the duration of exposure to the noise.

A number of peer reviewed studies have assessed the impact of noise on marine mammals. The most contemporary information is provided in the recommendations for marine mammals of the United States National Oceanic and Atmospheric Administration (NOAA), *Draft Guidance for Assessing the Effects of Anthropogenic Sound on Marine Mammals* (NOAA, 2013).

There is limited information regarding the effects of noise on fish. The studies that have been conducted to date only consider a tiny fraction of the species in existence and in environments which typically do not represent wild fish in their natural habitats (Popper and Hastings 2009). Fish are able to detect sounds, with the majority of species regarded as "hearing generalists" that have a narrow hearing bandwidth. A small number of fish species are classified as "hearing specialists" and have a greater hearing bandwidth and sensitivity due to a coupling between gas filled organs (such as the swim bladder) and inner ear (Hastings and Popper 2005). The hearing range for the different types of fish is provided below. Fish sensitivity to noise also depends on the mass of the fish. It has been found that tissue damage from noise will increase as the mass of the fish decreases (Carlson *et al.*, 2007).

Information regarding the known effects of underwater noise on turtles is limited. Independent studies by Ridgway and Bartol (cited in Bartol, 2008) confirm that turtles can hear and that the hearing range of turtles is approximately between 200Hz and 1000Hz. There are no recommended noise criteria for turtles, however based on behavioural response studies by O'Hara and Wilcox (cited in Bartol, 2008) and Moein (cited in Bartol, 2008), and the hearing range of turles, the low frequency Cetaceans noise criteria is considered appropriate.

There is very limited information known about the effects of underwater noise on penguins (penguins may utilise the port site for foraging purposes). Studies indicate that the hearing range of penguins is best between 2000Hz and 5000Hz in air and is likely to reduce to frequencies below 4000Hz in water (Dooling and Therrien, 2012). Based on the hearing range, and the lack of specific objective criteria, the low-frequency Cetaceans noise criteria have also been applied to penguins.

There is very limited information known about the effects of underwater noise on Cephalopods (cuttlefish). Studies indicate that Cephalopods can perceive low-frequencies. It is not known if Cephalopods can "hear" or if they are sensitive to particle velocity (Mooney *et al.*, 2012). Notwithstanding, it has been shown that they can perceive sounds with frequencies of up to 1.5kHz, but as they do not have any gas filled bladders there is no possibility for sound amplification and therefore have a hearing capacity comparable to fish without swim bladders (Hu *et al.*, 2009). Based on the above, the noise criteria for fish have been conservatively applied to Cephalopods for the assessment of impacts presented here.



The level of noise resulting in auditory injury changes for different species and types of noise is provided in Table 6-10 below, with the threshold level being reached if either the peak level or SEL_{cum} (cumulative sound exposure level), is exceeded (NOAA, 2013).

Table 6-10 Summary of relevant noise limits

| Species | Hearing Range | Impulsive Noise criteria | Non-impulsive noise criteria (includes continuous noise) |
|---|---------------------------|---|---|
| Low-frequency Cetaceans (Baleen whales) | Between 7 Hz and 30 kHz | Peak level: 230 dB _{peak} SEL _{cum} : 187 dB | Peak level: 230 dB _{peak} SEL _{cum} : 198 dB |
| Mid-frequency Cetaceans | Between 150 Hz and 160 | Peak level: 230 dB _{peak} | Peak level: 230 dB _{peak} |
| (Dolphins) | kHz | SEL _{cum} : 187 dB | SEL _{cum} : 198 dB |
| High-frequency Cetaceans | Between 200 Hz and 180 | Peak level: 201 dB _{peak} | Peak level: 201 dB _{peak} |
| (Porpoise) | kHz | SEL _{cum} : 161 dB | SEL _{cum} : 180 dB |
| Phocid Pinnipeds (Earless | Between 75 Hz and 100 kHz | Peak level: 235 dB _{peak} | Peak level: 235 dB _{peak} |
| Seals) | | SEL _{cum} : 192 dB | SEL _{cum} : 197 dB |
| Otariid Pinnipeds (eared sea lions and fur seals) | Between 100 Hz and 40 kHz | Peak level: 235 dB _{peak} SEL _{cum} : 215 dB | Peak level: 235 dB _{peak} SEL _{cum} : 220 dB |
| Fish (hearing generalist) | Between 50 Hz and 500- | Peak level: 206 dB _{peak} | Peak level: 206 dB _{peak} |
| | 1500 Hz | SEL _{cum} : 187 dB | SEL _{cum} : 187 dB |
| Fish (hearing specialist) | Between 50 Hz and 3-100 | Peak level: 206 dB _{peak} | Peak level: 206 dB _{peak} |
| | kHz | SEL _{cum} : 187 dB | SEL _{cum} : 187 dB |
| Turtles | Between 200 Hz and 1000 | Peak level: 230 dB _{peak} | Peak level: 230 dB _{peak} |
| | Hz | SEL _{cum} : 187 dB | SEL _{cum} : 198 dB |
| Penguins | Between 2000 Hz and 4000 | Peak level: 230 dB _{peak} | Peak level: 230 dB _{peak} |
| | Hz (in water) | SEL _{cum} : 187 dB | SEL _{cum} : 198 dB |
| Cephalopods (cuttlefish) | Unknown | Peak level: 206 dB _{peak} SEL _{cum} : 187 dB | Peak level: 206 dB _{peak} SEL _{cum} : 187 dB |

The underwater noise assessment for the proposed port development at Cape Hardy (Sonus, 2014) indicates that separation distances between some noise sources (i.e. piling) and specific marine fauna during construction will be required to minimise noise impacts. The impact criteria for single strike sound exposure level (SEL_{ss}) were taken from the DPTI Underwater Piling Noise Guidelines, and the criterion for all species is 150 dB (DPTI 2012). The full results of the underwater noise modelling are presented in Sonus (2014). The minimum separation distance between different species and noise sources required to achieve noise criteria (refer Table 6-10) is provided in Table 6-11.

Table 6-11 Separation distance required to meet requirements of noise criteria

| - | Separation distance required to achieve noise criteria (m) | | | | | | | | | | |
|------------------|--|-------------------------|---------------|--------------------------|-------------------------|-------------------|--------------------------|-------------------------|---------------|--------|---------------|
| | Low-frequency Cetaceans | | | Mid-frequency Cetaceans | | Otariid Pinnipeds | | | Fish | | |
| Noise Source | SELcum (NOAA 2013) | SELss (DPTI 2012) | Peak level | SELcum (NOAA 2013) | SELss (DPTI 2012) | Peak level | SELcum (NOAA 2013) | SELss (DPTI 2012) | Peak level | SELcum | Peak level |
| Impact Piling | 240 | 470 | 0 | <10 | 30 | 0 | <10 | 470 | 0 | 450 | 30 |
| Drilling | 0 | - | - | 0 | - | - | 0 | - | - | 10 | - |
| Vessels | 10 | - | - | 0 | - | - | 0 | - | - | 80 | - |



6.7.2 Impacts and Risks: Noise and Vibration

Noise and vibration associated with the proposed development could cause impacts to the marine environment from:

- Construction activities (e.g. piling and drilling)
- Vessel movement
- Operations activities (e.g. ship loader and conveyor)

6.7.2.1 Impacts

The construction phase of the proposed port will result in the highest level of noise emissions that may impact marine biota, due to the extent of piling works required. The underwater noise modelling concluded that the effects of other noise sources, such as vessels or equipment, will not increase cumulative noise levels above the peak levels produced through impact piling, as sound waves are not additive (Sonus, 2014).

The Sonus (2014) report identifies that a minimum separation distance of 470 m from piling to marine biota will avoid physical impacts to all species (based on DPTI / NOAA criteria). Physical trauma is anticipated to affect marine biota where:

- Baleen whales and seals are within 470 m
- Cephalopods are within 450 m
- Fish are within 30 m

The most sensitive fishes will be those with swim bladders, while fishes without swim bladders and cephalopods will most likely only suffer behavioural changes (avoidance of the piling site). Through the implementation of soft start procedures, safety observation zones to monitor movement of marine biota and shut down procedures where marine mammals are identified within 500 m of piling activities, it is not anticipated that any marine biota will be significantly affected by underwater noise emissions. Marine biota that may be affected are highly mobile and will likely vacate the study area during soft start procedures. As such, widespread physical trauma to marine biota is not anticipated.

Construction of the port will occur over an approximate 18 month period. Underwater noise emissions during this time will result in behavioural changes to marine fauna, with avoidance of the study area likely as a result of noise emissions. Following construction, marine fauna are expected to return to the study area. These altered arrangements represent a short term alteration to existing behavioural patterns of marine fauna within the study area, and are considered to be a low impact.

There will be some operational noise from the ore conveyor, ship loaders and equipment located on the jetty and wharf. As these noise sources do not have any direct connection to the water there will be limited noise propagation into the water from these sources. Some structure-borne vibration energy will travel through the jetty/wharf and into the water column. However, Sonus (2014) predict the structure borne propagation would only comprise low levels of sound that would be quickly attenuated to levels below the ambient underwater noise level due to the bathymetry of the site and will not impact marine fauna.

Propeller cavitation and flow noise around the hull are the primary causes of vessel noise and will occur throughout operation of the proposed port. While not causing direct physical harm, an increase in anthropogenic noise from sources such as industry and shipping can mask the sounds required for normal behaviour in marine biota (Codarin *et al.*, 2009). At low ship speeds, machinery noise dominates and is primarily low-frequency, although main gearboxes and gas turbines may produce higher-frequency sound. The predicted underwater noise at the port during operation will be dominated by the low frequencies as vessel speeds will be restricted in the port with cargo ships being manoeuvred into position by tugs. Due to the bathymetry of the site, modelling predicts sound propagation in the port area will be restricted (Sonus 2014).



There are no particularly sensitive receptors in the study area (i.e. no marine mammal breeding colonies or bird rookeries), and operational noise will result in negligible impacts to the behaviour and physical wellbeing of marine fauna with an insignificant increase in background noise levels within the study area.

Outside of the study area, increasing vessel numbers in the gulf associated with the port operations will increase the ambient noise levels in the marine environment, as low-frequency sound can travel great distances in the open water environment. This noise can mask communication and navigational sound used by many marine species. The proposed port will result in approximately one additional ship movement per day through the Spencer Gulf (an overall increase in vessel traffic of approximately 7% based on 290 additional vessel movements or 145 vessels per year). Sonus (2014) predicts that separation distances of 80 m for fish and 10 m from whales would be required to prevent physical damage from low-frequency vessel noise. In deeper water low-frequency sound propagates further than high frequency sound enabling most fish and larger marine fauna to avoid any physical damage caused by noise impacts by moving away from high—intensity, low-frequency sounds before physical damage occurs. As such, the affect to ambient noise levels within the gulf as a result of additional ship movements is considered to be insignificant and result in negligible impacts to marine fauna.

6.7.2.2 Risks

A risk associated with underwater noise and vibration in the study area is incorrect assumptions regarding the presence or absence of marine fauna in the area. If additional fauna are present within the area than identified by the review undertaken here, potential impacts to marine fauna from noise and vibration could be greater than predicted. Given the modelling has included all marine fauna types, and exclusion zones calculated from the modelling are conservative, the application of management measures such as soft starts and observation zones for larger fauna should mitigate additional impact. As such, the consequence of additional fauna being present in the area is considered to be minor. The likelihood of additional fauna than those identified here being present is considered to be possible. As such, the risk to marine fauna as a result of additional species being present than identified here is considered to be low.

In the event that the underwater noise and vibration modelling is incorrect, and sound propagates further and at higher amplitudes through the marine environment than predicted, impacts to marine fauna could be more widespread and severe. As above, the exclusion zones proposed which dictate management strategies for shut downs are conservative and already well exceed the predicted impact area based on the modelling, meaning additional impacts are considered unlikely. Consequences of greater than modelled impacts are considered to be minor and therefore overall risk as a result of underestimation of modelled impacts is considered to be low.

6.7.3 Conclusions

Combining the knowledge gained from the underwater noise modelling, publically available reports and legislative guidelines, it is considered that the noise impacts at the Cape Hardy site during piling will represent the largest impact to marine fauna (which is short term), while operational noise from the proposed facility is considered to have no detrimental impacts to the marine environment.

Through the implementation of management procedures (such as soft start piling and observation zones) all impacts associated with noise to marine fauna have been reduced to low or negligible.

6.8 Invasive Marine Species

IMS are marine organisms that are not native to an area but have been introduced by human activities such as shipping. Australia has over 250 introduced marine species. Most of these have little impact. Some however, have the potential to establish large populations and spread geographically to the detriment of native species and/or human uses. IMS are globally recognised as a significant threat to the marine environment (GISD, 2014). Several crabs, mussels, seastars and macroalgae have become aggressive pests in a number of locations. Once



established, IMS can either directly prey upon, displace or outcompete indigenous species. IMS also have the potential to carry diseases which can eliminate native species.

Potential methods of IMS introduction at ports include organisms being attached to cargo imported to sites, within ships' ballast water as cysts, larvae or juveniles, or on the hulls of vessels or vessel infrastructure as biofouling (encrusting organisms) that can break off and reattach from asexual growth or release spawn.

6.8.1 Existing Environment and Key Findings

No IMS were identified as known to exist within the study area during the course of the desktop investigations (Ditmann *et al.*, 2010; Wiltshire *et al.*,2010; NIMPIS, 2014), video surveys or site visits. Within South Australia and specifically Spencer Gulf several IMS have been detected or established themselves and are discussed below. Species such as *Salmo salar* (Atlantic salmon), *Crassostrea gigas* (Pacific oyster) and *Ostrea edulis* (European flat oyster) are not considered to be IMS as they are not on the National or State list of declared pests (NIMPIS, 2014) and in the case of Atlantic salmon are an unlikely species to be transported via port operations.

Caulerpa taxifolia is an invasive marine alga that is widely used as a decorative plant in aquaria. Once in a suitable marine environment it is extremely invasive and smothers other algal species, seagrasses and sessile invertebrate communities forming dense monocultures that exclude much marine life. It does this by either out-competing species for nutrients and light or due to a toxic compound it produces (caulerpenyne; that has almost no known grazers). The primary effects on humans are mostly related to the reduction of catches for commercial fishermen due to the elimination of fish habitat by *C. taxifolia*. This IMS can also foul nets, ropes and moorings. In South Australia it is only known to have established itself within Port Adelaide and its presence is thought to be related to the nutrient loading and warmer waters of that estuary (Wiltshire, 2010).

The Asian paddle crab or lady crab, *Charybdis japonica*, has only been identified in South Australia from one male specimen. It can vary in colour from red / purple / orange to pale green and off-white; it is similar in appearance to the native blue swimmer crab. No other specimens have been found since its initial discovery and it is not currently considered to be present in Australia. It is an aggressive crab that can out-compete native species for food and habitat. It is also known to carry the white spot syndrome, a virus that can impact native and farmed prawns, crabs and lobsters (NIMPIS, 2014).

Codium fragile ssp. tomentosoides is an alga that has been introduced around the globe through shellfish aquaculture, recreational boating, and transport on ship hulls. This IMS exhibits various modes of reproduction, which is a common trait of many IMS. It can reproduce sexually, parthenogentically, and vegetatively which makes eradication difficult. It can propagate from small pieces of the parent plant that can be broken off by propeller wash or hull cleaning (from recreational vessels) and then carried by water currents over long distances, introducing it to new locations. This IMS is also tolerant of a variety of salinity and water temperature levels and thrives in sheltered habitats. This IMS can cause a nuisance to humans when it accumulates on beaches and rots, producing a foul odour. Its main threats to the marine environment come from its tendency to overgrow and smother shellfish beds, often attaching to the shells of oysters, mussels, scallops, and clams. The attached algae can hinder the movement and feeding of the shellfish. There are several direct and indirect effects of this attachment including: smothering shellfish by preventing opening of the valves, clogging aquaculture nets, fouling moorings, interfering with the collection of shellfish and where the attached alga is relatively large and wave exposure is high, the shellfish can be swept away with the alga. This species has been reported as occurring in the Spencer Gulf and is listed as a threat to aquaculture in the area by the global invasive species database (GISD, 2014).

The Asian date or bag mussel, *Musculista senhousia*, is a small, short-lived mussel native to East Asia. This IMS has been identified within Port Adelaide (Wiltshire and Deveney, 2011) and more recently near Lipson Cove during a benthic habitat survey (Golder Associates, 2012). This IMS can grow rapidly and is capable of marked



habitat alteration through reaching high densities (>2000 individuals/m²) on the surface of soft sediments. It can form continuous byssal carpets that smother most other benthic habitat-forming organisms (GISD, 2014). As with many IMS it is an opportunistic species capable of fouling wharf pilings and artificial structures and can be found from intertidal to subtidal soft bottom habitats (to a depth of 20 m). The species is a filter-feeder and due to its high densities can have dramatic impacts on plankton abundance and reduce the densities of native bivalves. High densities of this IMS can dramatically alter the natural benthic habitat, changing both the local physical environment, and the resident macroinvertebrate assemblage and the growth of nearby seagrass (GISD, 2014).

Polysiphonia brodiei (red macroalgae) is a branched filamentous red alga that has become one of the most globally widespread IMS. It has been introduced via ships to Australia, New Zealand and Japan. It has been confirmed around Robe and American River, but not within the Spencer Gulf. It is a hull-fouling organism on slow-moving vessels such as barges, and also fouls ropes, buoys and harbour structures such as pylons and boat ramps. Apart from biofouling there are currently no other known impacts from this IMS (GISD, 2014).

Sabella spallanzanii (the European fan worm), is a filter-feeding tube worm which has the potential to alter native marine ecosystems and compete with native organisms for food and space. It may also inflict economic damage by competing with mussels and oysters in aquaculture farms. These IMS readily settle on shellfish grow-out lines and may reduce shellfish growth by altering water flow around the lines and competing with the shellfish for suspended food. Some species of seagrass can be impacted by the settlement of worms on their fronds (CSIRO, 2001). The European fan worm is considered a major threat to benthic assemblages in both hard and soft bottom habitats and potentially affects nutrient cycling processes in soft sediments (O'Brien et al., 2006). It has been estimated that S. spallanzanii can filter around 12 m³/d/m² of habitat (Stabilia et al. 2006). This rate of filter-feeding can drastically impact plankton abundance including larval fish, and therefore the impacts of this species go beyond smothering and biofouling. The species has been identified in South Australia in Port Augusta and upper Spencer Gulf (BHP Billiton, 2009), Port Lincoln and Port Adelaide (including Westlakes waterways where it has established itself (Conservation Council SA 2011). The species can be transported by cargo vessels (either on the hull, in ballast water or attached to cargo), fisheries and aquaculture (by vessels, accidental translocation through aquaculture farm activities and movements), recreational vessels (hulls and moorings) or through natural dispersal once it has established in an area. As the species is already established in Spencer Gulf it is likely that this IMS will reach the port site and most areas of the Gulf over time.

Watersipora spp. are encrusting bryozoans. The risk from these IMS are their tolerance to copper-based antifouling coatings on ships' hulls which can then facilitate the fouling and spread of other IMS. This IMS is considered widely invasive in temperate ports (GISD, 2014). They are tolerant of copper-based antifoulants and can provide a non-toxic surface for other fouling species to settle. In addition to assisting in the spread of other IMS, Watersipora can also compete with native bryozoans and other encrusting organisms and once established are often the most common intertidal bryozoan. In Australia W. arcuata is declared a medium priority pest and is widespread in South Australia, including the southern Spencer Gulf (NIMPIS, 2014; GISD, 2014).

The European or green shore crab (*Carcinus maenas*) is a voracious predator that feeds on many types of organisms including shellfish and other crabs. This IMS has great potential to impact aquaculture when present in high densities as it is able to crush mussels, and can threaten mussel farms. In invaded regions, this species has been considered responsible for significant impacts on epibenthic and infaunal species, such as bivalves, other molluscs, and crustaceans, through predation, competition, and burrowing (GISD, 2014). This species is a potential facilitator of another IMS, *Styela*, which is an invasive club tunicate in some areas (GISD, 2014). The crabs could facilitate *Styela* invasions by preying on tunicate predators, enabling *Styela* to establish. The European shore crab is known to consume prey from at least 158 genera and has been widely documented to decrease the diversity and biomass of estuarine communities. The species has been previously identified in



Port Adelaide; however recent surveys have failed to detect this crab in South Australia (Wiltshire and Deveney, 2011).

Alexandrium minutum is a toxic, planktonic dinoflagellate (single-celled alga) responsible for toxic algal blooms in warm, temperate, coastal and estuarine waters. It has been reported over a wide geographical range and in a wide range of coastal hydrographic regimes (Hallegraeff et al., 1988; Yoshida et al., 2000; Usup et al., 2002). This IMS produces compounds which are toxic to some zooplankton and fish. The toxins can also bioaccumulate in zooplankton and shellfish. The consumption of contaminated shellfish can lead to paralytic shellfish poisoning in humans and other mammals. The toxins responsible for this disease are neurotoxins, which in humans can cause muscular paralysis, neurological symptoms and, in extreme cases, death (Hallegraeff, 1993). Due to the potential for disease outbreaks the occurrence of algal blooms near shellfish farms usually results in their closure. This IMS is classed as a high priority species for temperate water ports (NIMPIS, 2014)

Alexandrium minutum is restricted to coastal, nutrient-enriched sites, particularly harbours, estuaries and lagoons (GISD, 2014). Blooms are often associated with low salinities and nutrient-rich freshwater inputs (Vila et al., 2005). The growth rate of A. minutum increases with increasing temperature and light (Lim et al., 2006); nevertheless it can reach relatively high growth rates (up to 0.5 divisions/day) at 12 °C (Cannon, 1996). The algae produce cysts in bloom areas, which can accumulate in fine estuarine and coastal sediments, and persist for years (Garcés et al., 2004; Bravo et al., 2006). Given that the environmental conditions linked to outbreaks of this IMS are not dominant in the Cape Hardy area its introduction at Cape Hardy is considered unlikely. The coastline is not enclosed nor are there regular inputs on nutrient-rich freshwater required to support this IMS.

6.8.2 Impacts and Risks: Invasive Marine Species

IMS are opportunistic species that use either marine operations as vectors to transfer between areas, or disturbances to the marine environment or changes in the local environment along with artificial surfaces to colonise new areas. As a port area is likely to provide all three of those scenarios during both construction and the ongoing operation of the port, introduction of IMS has the potential to significantly affect the marine environment at Cape Hardy.

6.8.2.1 Impacts

The introduction of IMS to the study area as a result of the construction and operation of the proposed port is not planned. As such, IMS are considered to be a risk to the project rather than a known impact and are discussed below.

6.8.2.2 Risks

IMS could be transported to the study area as biofouling on vessel hulls, jack-up barge legs, anchors, anchor chains, mooring lines, internal boat compartments, sediment transported in or on vessels, or in any seawater onboard vessels/barges including ballast water, in bilge, inside pipes or pumps. IMS are not selective and will take advantage of any vessel regardless of size. Therefore the jack-up equipment or support vessels used during construction, as well as cargo ships during operation all have the potential to transport IMS to the study area.

IMS can exist in low numbers or persist as cysts in an area and can rapidly increase in numbers after a disturbance to the environment or removal of competitive indigenous species. Therefore construction activities have the potential to either release cysts or propagules of IMS as well as provide cleared surfaces for IMS to colonise. Due to the opportunistic traits of IMS the disturbed areas and new hard surfaces of the MOF, tug harbour and jetty will create suitable areas for IMS to colonise.



All vessels utilising the port site will be required to comply with the national guidelines relevant to biofouling and ballast water. However, the identification of IMS during construction and operation of the proposed port is considered likely despite the implementation of control measures. As Cape Hardy is relatively isolated from a hydrodynamic perspective, IMS that may colonise the area would not be readily transported by currents beyond the study area and into new regions. As such, most IMS would rely on vessel movements for transportation to areas beyond the study area. Therefore, of the effects of water bourne IMS are considered to be isolated to the Cape Hardy study area and are not considered likely to affect the marine environment at a regional level, considered to be of moderate consequence. As such, the overall risk associated with IMS to local flora and fauna species is considered to be high.

Although the introduction of IMS is a possibility, the likelihood of introducing an algal IMS species that effects water quality is considered unlikely due to the Commonwealth requirements for vessels to discharge ballast water in open waters and the requirement for operators to adhere to a CEMP and OEMP for the port. The consequences of introducing an algal IMS species would be considered minor (localised to the marine study area and algal blooms are typically short term in duration). As such, the risk to water quality from IMS algae is considered to be low.

Some IMS such as mussels have the potential to build up biological reefs consequently changing the bathymetry and therefore ultimately the hydrodynamics of an area. Seagrass has also been shown to influence the hydrodynamics of coastal areas by physically slowing currents. Therefore if a reef-building seagrass pest or smothering IMS such as *Caulerpa taxifolia* were introduced to the site it could cause seagrass loss, which could in turn lead to alterations in hydrodynamics and sediment dynamics. The consequences of such an introduction will be limited to the local study area and will likely be present for a long term, and are therefore considered moderate. As with any operating port the introduction of IMS is a possibility, however the likelihood of introducing a specific reef building or seagrass smothering IMS is considered unlikely. Therefore, the risk of IMS to hydrodynamics and bathymetry is considered medium.

6.8.3 Conclusions

Due to the ability of IMS to utilise multiple modes of transport within the marine environment (i.e. any vessel big or small, from small fishing boats to commercial ships to aquaculture operations) and the interconnected nature of marine environments the risks from IMS are already present within the Spencer Gulf and South Australia.

No IMS were identified within the Cape Hardy area during the course of the investigation. IMS have, however, been detected in several locations around South Australia and the presence of a port will provide both suitable substrate for IMS to colonise and to establish as well as vectors for IMS to spread in Spencer Gulf waters. As such, risks associated with the establishment of IMS at the study area to local flora and fauna species are considered to be high. IMS can also affect bathymetry and therefore ultimately the hydrodynamics of an area, however this is considered unlikely to occur, with an overall risk of IMS to hydrodynamics and bathymetry considered medium.



6.9 Fisheries and Aquaculture

Spencer Gulf is an important fisheries and aquaculture area, with a number of commercial fisheries throughout the gulf and designated Aquaculture Zones. Commercial fishery species in the gulf include Abalone, Blue swimmer crab, Garfish, Western King Prawn, King George Whiting, Australian Salmon, Sardines, Snapper and Calamari. In recent years there have been a number of restrictions on fisheries including suspension of trawling and the introduction of Marine Park Sanctuary Zones that prohibit fishing. Aquaculture in the gulf is a growing industry, with farmed marine species including Pacific Oysters, Kingfish, Mulloway, Mussels and Bluefin Tuna. Currently 670 ha of aquaculture is permitted within restricted locations of the Spencer Gulf. Aquaculture has been excluded from certain areas for water quality, navigation, conservation, historic, recreation or commercial fishing reasons.

6.9.1 Existing Environment and Key Findings

Fishing and aquaculture have a significant commercial role in the Spencer Gulf region. Other than agriculture, these industries provide the bulk of income and employment in the region. Port Lincoln (~100 km south of Cape Hardy) is the major centre of fisheries and aquaculture production in South Australia. Arno Bay (~50 km north of Cape Hardy) is another centre of fisheries and aquaculture.

The proposed port site lies within the Port Neill Aquaculture Exclusion zone as delineated by the Land Not Within a Council Area (Coastal Waters) Development Plan. There were two exemptions within the Port Neill Aquaculture Exclusion Zone for registered finfish aquaculture leases numbered LA00071 and LA00072 (Aquaculture Zones, 2008), however these were surrendered to PIRSA in 2013 (personal comms. E. Kaese, PIRSA, 2013). As these leases have been surrendered, no aquaculture activity is permitted within the Port Neill Aquaculture Exclusion Zone.

Currently the closest active aquaculture leases lie 30 km to the south of the study area off Cape Euler where there is a land based lease for abalone, calamari, greenback flounder, blue and pen mussels, native and pacific oysters, scallops, sea urchin, cockles, razorfish, pot-bellied seahorse and leafy seadragon. These commercially reared species are all held in onshore tanks. There is also an offshore lease in the historic Tumby Offshore Aquaculture Zone, with offshore seacages for three abalone species. Thirty three km to the north of the study area in the Arno Aquaculture Zone are several leases with the majority of the licenses for finfish sea cages of bluefin tuna, yellowtail kingfish, mulloway, snapper and black bream. There are also land based operations with license for King George whiting, knifefish, calamari, abalone and pacific oyster at Arno Bay.

It should be noted that while the Port Neill Aquaculture Exclusion Zone provides a buffer from potential aquaculture activities, the Port Neill Aquaculture zone extends to within 2 km of the proposed port location at Cape Hardy (see Figure 6-10). There are currently no active leases within that area however it is possible that at some point in the future the area will be utilised by some form of aquaculture.

During the towed video surveys several commercial fish and invertebrate species were observed. Numerous blue swimmer and sand crabs were sighted throughout the survey area along with several slipper lobsters near seagrass beds and one giant cuttlefish. The Australian Giant Cuttlefish and all cephalopods are protected by the Fisheries Act in the upper Spencer Gulf by the temporary cuttlefish closure area until (at least) February 2015. The protected area extends from the top of the gulf down to Arno Bay and Wallaroo; approximately 35 km north of Cape Hardy.




Figure 6-10 Port Neill aquaculture exclusion zones and Port Neill aquaculture zone

E-F-34-RPT-0039_A (Marine Environmental Tech Report_Rev1a - compared doc 24/09/2015



Spencer Gulf is also an important area for a number of commercial fisheries, including those targeting (western) king prawn, blue swimmer crab, abalone, rock lobster, sardines and marine finfish such as snapper and King George whiting (Knight and Tsolos, 2012).

The northern end of the gulf is an important spawning and nursery area for king prawns and supports an economically valuable fishery. The Spencer Gulf has been subdivided into management areas for the prawn fishery and the Cape Hardy area falls within the Cowell zone which encompasses Cowell to the north down to Tumby Bay in the south. The mean Catch per Unit Effort (CPUE) for the Cowell zone from 2005-11 was approximately 80 kg/h (Dixon et al., 2012). Generally, within the gulf higher CPUEs are observed to the north and west. The annual catch for the Cowell zone was 80 and 17 tonnes in 2009 and 2010 respectively (Dixon et al., 2012), while, annual catches for the upper gulf have been as high as 150 tonnes. Prawn fishing in the Spencer Gulf is restricted to between the months of November to June and generally during nighttime hours, in waters over 10 m in depth (Dixon et al., 2012). Cape Hardy falls within the prawn fishing block 107. An independent survey of block-specific data for this fishery shows there has been no historic trawling effort in block 107 (Currie et al., 2009) however there may be some interaction with the fishery by the vessels accessing the port within the deeper water of the gulf. Currie et al. (2009) show that the heaviest trawling effort is in shipping channel areas of the upper gulf, indicating that the prawn fishery can coexist with large cargo vessel movements. The proposed vessel approach vectors (Figure 3-3) and the port itself, represents only a small portion of the overall fishery and it is not anticipated that the construction or operation of the port will restrict access to the deeper water areas trawled for this species.


A blue swimmer crab (*Portunus armatus*) fishery operates in the Spencer Gulf and is managed under the Fisheries Management (Blue Crab Regulations, 1998) which closes the fishery from 21 December to 19 February. Cape Hardy falls within the fishing block 59 of the Spencer Gulf Blue Crab fishery which includes all waters north of Tumby Bay. No block-specific data is available for the block 59 blue crab fishery. The dominant fishery for this species is greater than 100 km from Cape Hardy in the upper Spencer Gulf fishery north of Cowell (Dixon *et al.*, 2013).

The blue swimmer crab is distributed throughout coastal waters of tropical regions of the western Indian Ocean through to the southern temperate waters of Australia and into the eastern Pacific Ocean. In South Australian waters the species has altered its life cycle to have increased growth and reproduction during the warmer months of the year when water temperatures increase to those similar to tropical regions. The species' habitat range is also large with the crabs found on algal and seagrass habitats on both sandy and muddy substrates from intertidal to at least 50 m depth. Within South Australia there is a seasonal movement of adult crabs from deeper waters into inshore shallow waters during the warmer months of September to April during the breeding season and into deeper water offshore during the colder months of May to August. Observations made during the video survey in November 2011 recorded the species as widely distributed in areas of both sandy bottom and seagrass beds. It is therefore likely the species utilises the Cape Hardy area. However due to the large range of this species, the location of the study area 100 km south of the dominant fishery for this species and the size of the port footprint, the benthic habitat at Cape Hardy is not expected to be on critical habitat for this species or its fishery.

The Northern Zone rock lobster fishery operates in Spencer Gulf between November and May. Cape Hardy falls within Marine Fishing Area (MFA) 29 for this fishery. The species distribution is typically around offshore islands and rocky reefs. In 2011 the total catch estimates for the species indicated that 19% of the catch came from the southern Spencer Gulf (SSG) with the majority of the catch coming from the far west coast of South Australia and Kangaroo Island (80% combined) (Linnane *et al.*, 2012). The catch for this species across South Australia have been declining since 2000 despite increased effort; attributed to low juvenile recruitment. The species has a four-year settlement and recruitment cycle and in 2007-2012 there was low settlement (Linnane *et al.*, 2012). This indicates that recruitment from 2011-2016 may be low and catch numbers may be reduced for several

years for this species in the SSG. The historical catch from MFA 29 represents only 0.02% of the total Northern Zone catch between 1993 and 2009 (Ward and Burch, 2012), indicating the area around Cape Hardy is not a key area for this fishery. Given the low percent catch attributable to MFA 29 and the limited subtidal reef areas present within the Cape Hardy study area the interaction with rock lobster fishing is expected to be minimal.

Green-lip and Black-lip Abalone are also fished in the Spencer Gulf with permits restricted to a number of licensed divers. Cape Hardy Falls within Abalone fishing block 20c. There is no detailed fishing activity or catch data available for 20c, however the relative catch numbers for the State indicate 20c is not a significant fishery area for the species (Figure 6-11). It is anticipated that the bulk of the limited catch from 20c has historically come from near the offshore islands of the Sir Joseph Banks group. Abalone is typically found on rocky substrate and subtidal low reef areas which are not extensive within the study area. Based on the known environment surrounding Cape Hardy and the relative catch data for this fishery, it is not anticipated that the proposed port area is heavily utilised by Abalone divers.



Relative importance (% of catch) of each blacklip (black bars) and greenlip (green bars) SAU. Note each SAU is ranked twice, once for blacklip and once tor greenlip. Red text and dotted lines indicate SAU importance and division. Note abbreviations for cape (Cp), island (Isl), Joseph Banks (JB) and Unass RG A or Unass RG B (unassigned Region A or Region B, respectively).

Figure 6-11 Relative catch percentage of Abalone for Western Zone Region A, taken in 2010/11.

Note: red ovals indicate relative total catch from the Sir Joseph Banks SAU which includes the study area (Adapted from Stobart et al., 2012)

There are also three main finfish fisheries that operate in the Spencer Gulf; the Sardine, King George whiting and Snapper. Cape Hardy lies within the SSG fishery and the Marine (finfish) Fishery Area (MFA) 29 which is over 4250 km².

The SSG is an important area for the Sardine fishery providing the largest catch for the past 15 years (Ward *et al.*, 2012). The bulk of the catch is from deep water areas of MFAs 39, 30 and 29 (Knight and Tsolos, 2012). Sardines are a small pelagic schooling fish and are generally fished in South Australia using purse seines.



The commercial catch for King George whiting in the SSG has predominantly used hand lines since net closures in 2005. Recorded commercial catches for the SSG are approximately 73 tonnes in 2008, 82 tonnes in 2009 and 80 tonnes in 2010. Only the far west coast of Eyre Peninsula recorded higher catches within South Australia during this period (Fowler *et al.*, 2011). There is no commercial fishery data available for MFA 29 as data from all the MFA (29, 30, 31, 32 and 33) in SSG have been pooled together (Fowler *et al.*, 2011). While the SSG is a substantial fishery for the King George whiting, the fishery is widespread throughout South Australia waters, and the Cape Hardy area does not represent any key nursery grounds therefore little interaction with this fishery is expected.

There is no dedicated giant crab fishery in South Australia as most of the crabs are taken as bycatch in the lobster fisheries (within depths shallower than 120 m; Levings *et al.*, 2001; Currie, 2011; Stobart, 2014). The giant crab is endemic to southern Australia and found between southern Western Australia and central New South Wales (Kailola *et al.*, 1993) inhabiting waters between 18 and 400 m in depth and in temperatures between 10 and 18 ° C. The highest population densities occur at the shelf break, where the continental shelf and slope meet (~200 m in depth). Size distribution is stratified by depth (Levings *et al.*, 2001). Circumstantial evidence (based on catch records) suggests that females move onto soft muddy substrate when they moult while males are captured across a broader depth range than females (Levings *et al.*, 2001). As this species' preferred depth range and habitat generally falls outside of the footprint of the study area, direct effects on commercial fishing of this species from the development are unlikely.

Commercial fishing of Garfish in the SSG falls into the MFA 29 and had an average catch of less than 0.5 tonnes for 2010/11, with previous years 2004-2010 between 0.5 -3.0 tonnes (Steer *et al.*, 2012). Recreational fishing for Garfish fell from 5 tonnes in 2001 to less than 0.5 tonnes in 2008. The reduction in catch for both recreational and commercial tonnage has been attributed to area closures for haul netting and a shift to dab netting in the SSG. It is unlikely that Garfish would be present in large numbers within the Cape Hardy study area. The species diet is composed significantly of *Zostera muelleri* (commonly called Garweed) and *Heterozostera nigricaulis* (Steer *et al.*, 2012). There are no large stands of these seagrass species identified in the study area. The nearshore areas of the study area were not surveyed with video tow and may contain patches of *Zostera muelleri* or *Heterozostera nigricaulis*, however the aerial imagery indicates little coverage of seagrass in the nearshore. Therefore the total geographic area of either *Zostera muelleri* or *Heterozostera nigricaulis* distribution in the study area would not be significant. As such, any impacts on the seagrass areas within the study area due to the construction and operation of the port are unlikely to impact on the distribution of Garfish.

There are also a number other fisheries that operate within the wider Spencer Gulf, these species along with species that may be taken recreationally or as bycatch are included in Table 5-5 and Table 6-9 which assess their likelihood of occurrence within the study area as well as the likelihood of potential impacts from the development.

6.9.2 Impacts and Risks: Fisheries and Aquaculture

Impacts to fisheries and aquaculture could occur as a result of:

- Loss of habitat through clearance, exclusion or degradation of the environment
- Exclusion from fishing grounds
- Pollution from run-off, oil or chemical spills
- Marine debris
- Noise and physical impacts during construction piling
- Safety and navigation
- IMS



6.9.2.1 Impacts

Construction and operation of the proposed port will restrict commercial fisheries from operating within the marine study area. As previously discussed, noise emissions during construction and operation of the proposed port may result in the temporary displacement of fish species from the study area. The entire marine study area comprises approximately 1% of MFA 29 and does not contain any habitat identified as critical for commercially fished species. As such, restricting commercial fisheries from operating within the study area will not significantly impact commercial fishing operations and is considered to represent a negligible impact.

6.9.2.2 Risks

As fisheries are dependent on availability of fish the risks to fisheries are in general the same as those to fish species and are discussed in Section 6.6.

As previously outlined, the introduction of IMS is considered likely and may result in the long term decrease in abundance of seagrass and marine species, and subsequently affect the the distribution of fish species. As detailed in Section 6.8, IMS is considered to represent a high risk to fish species. Although IMS may affect the distribution of fish species, the marine study area does not contain any nursery grounds or other habitat identified as critical for commercially fished species. As such, the introduction of IMS is considered to be of minimal consequence to fisheries and aquaculture, representing a low risk.

Additional risks to commercial fisheries could occur as a result of the release of waste products into the marine environment. Construction and operation of the proposed port will result in the generation of a number of waste streams that may enter the marine environment. Marine debris can significantly affect fish species as a result of pollution, injury through collision, entanglement or ingestion of non-biological products.

Waste management and handling procedures developed as part of the CEMP and OEMP will control waste streams with the overarching aim of no waste products entering the marine environment. Despite the implementation of control measures, it is considered almost certain that some form of waste / debris will enter the marine environment during construction and operation of the port. As the port site is not a key habitat or breeding area, the consequences of debris entering the marine environment are considered to be minimal; insignificant to the overall viability of fish species. As such, the overall risk is considered to be low.

Fish species are also at risk from the unintended discharge of pollution in the form of site run-off, wastewater, hydrocarbons or chemicals. With any marine activity there is potential for waste or spills to enter the marine environment and affect plankton, benthic fauna and habitat forming benthic flora (e.g. seagrass), reducing habitat suitability for fish species.

The implementation of design measures (refer Section 4) for the control of surface and waste water at the port site will limit the majority of hazardous pollutants from entering the marine environment. There remains a risk that a major spill of oil or other chemicals may occur as a result of vessel failure or accident. Based on the experiences of other operating ports, the likelihood of this event occurring is considered to be unlikely. As previously outlined, the port site does not represent a critical habitat or breeding colony for fish species. The consequences of a major spill event are considered to be minor, due to the low level of dependence of fish species on the study area, resulting in a short term decrease in the local abundance of fauna. As such, the overall risk of a major spill event affecting fish species is considered to be low.

6.9.3 Conclusions

The port is located within the Port Neill Aquaculture Exclusion zone, with the Port Neill Aquaculture zone located 2 km from the nearest point of the jetty. There are currently no aquaculture leases or licenses located within the Port Neill Aquaculture zone (PIRSA, 2014). Aquaculture leases with licenses for yellowtail kingfish



that were previously approved within the Port Neill Aquaculture Exclusion zone, have been withdrawn or surrendered and are no longer permitted in the Exclusion zone.

Cape Hardy lies within a number of areas designated for specific shellfish and finfish fisheries. It is anticipated that no interactions are likely in regard to the Prawn, Rock lobster and Abalone fisheries. It is anticipated that due to depth requirements for fishing methods the utilisation of the port site infrastructure areas by commercial finfish fisheries would be limited. The deep-water offshore areas of southern Spencer Gulf which have historically provided high catches of snapper, King George whiting and sardines are considered to be outside of the proposed port site and only the vessel approaches and anchorage areas would potentially interact with these fisheries during only limited times.

Although the introduction of IMS is considered a high risk to the distribution of fish species, the marine study area does not contain any habitat identified as critical for commercially fished species. As such, the introduction of IMS is considered to be on minimal consequence to fisheries and aquaculture, representing a low risk. Similarly, the unintended discharge of waste or pollution would not affect critical commercial fishery habitat or aquaculture areas and are considered to represent a low risk.



7 References

Adelaide Dolphin Sanctuary (2007) *Dolphin Reference Paper 1*. Prepared for the Department of Environment, Water and Natural Resources, South Australia

Ainslie RC, Johnston DA, and Offler EW (1989) *Intertidal communities of Northern Spencer Gulf, South Australia,* Transactions of the Royal Society of South Australia, Adelaide.

Atlas of Living Australia (2013) online database accessed at http://www.ala.org.au/species-by-location/ in November 2013. The Atlas of Living Australia is the Australian node of the Global Biodiversity Information Facility (GBIF).

Albert S, O'Neil JM, Udy JW, Ahern KS, O'Sullivan CM and Dennison WC (2005) *Blooms of the cyanobacterium Lyngbya majuscula in coastal Queensland, Australia: disparate sites, common factors,* Marine Pollution Bulletin.

Atkinson CA, Jolley DF, Simpson SL (2007) *Effect of overlying water pH, dissolved oxygen, salinity and sediment disturbances on metal release and sequestration from metal contaminated marine sediments,* Chemosphere

Atlas of Eh-pH diagrams. Intercomparison of thermodynamic databases. Geological Survey of Japan Open File Report No.419. National Institute of Advanced Industrial Science and Technology Research Center for Deep Geological Environments. Naoto TAKENO. May 2005

Appelo CAJ, and Postma D (2005) *Geochemistry, groundwater and pollution*. 2nd edition, Balkema, Leidan.

Aquaculture Zones (2008) Aquaculture (Zones-Port Neill) Policy 2008 under the Aquaculture Act 2001 available online at http://www.legislation.sa.gov.au/LZ/C/POL/AQUACULTURE (ZONES - PORT NEILL) POLICY 2008.aspx

Australian Museum (2013) http://australianmuseum.net.au/Australian-Pelican website accessed November 2013

Australian Quarantine and Inspection Service Australian Ballast Water Management Requirements (2001) and the Australian Quarantine Regulations (2000)

Baker JL (2004) *Towards a System of Ecologically Representative Marine Protected Areas in South Australian Marine Bioregions - Technical Report* Department for Environment and Heritage South Australia

Baker JL (2011) *Reef Fishes of Conservation Concern in South Australia - A Field Guide* Booklet produced with support from the Adelaide and Mt Lofty Ranges Natural Resources Management Board, South Australia.

Bannister, J.L. (2001) Status of southern right whales (Eubalaena australis) off Australia Journal of Cetacean Research and Management

Bannister JL, Kemper CM and Warneke RM (1996) *The Action Plan for Australian Cetaceans* [Online]. Canberra: Australian Nature Conservation Agency. Available from: http://www.environment.gov.au/coasts/publications/cetaceans-action-plan/pubs/whaleplan.pdf.

Bartol, M, (2008) A Review of auditory function of sea turtles. *Bioacoustics: The International Journal of Animal Sounds and it Recording*, vol. 17, no. 1-3 pp 57-59.

Beck MW, Heck KL, Able KW, Childers DL, Eggleston DB, Gillanders BM, Halpern B, Hays CG, Hoshino K, Minello TJ, Orth RJ, Sheridan PF, and Weinstein MR (2001) *The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates*, Bioscience.



Best PB, Brandao A and Butterworth DS (2001) *Demographic parameters of Southern Right Whales of South Africa* Journal of Cetacean Research and Management

BHP Billiton (2009) *Olympic Dam Expansion. Draft Environmental Impact Statement*. Prepare by Arup Pty Ltd and ENSR Australia Pty Ltd (Arup/ENSR) on behalf of BHP Billiton Olympic Dam Corporation Pty Ltd.

Biological Database of South Australia (2013) Biological Database of South Australia search results for study area with a 10 km buffer, accessed 2013

BOM (2014) Bureau of Meteorology website http://www.bom.gov.au/cgi-bin/oceanography/tides

Bowie AR, Maldonado MT, Frew RD, Croot PL, Achterberg EP, Mantoura FC, Worsfold PJ, Law CS, Boyd PW (2001) *The fate of added iron during a mesoscale fertilisation experiment in the Southern Ocean*, Deep-Sea Research II.

Bravo I, Garcés E, Diogène J, Fraga S, Sampedro N and Figueroa R I (2006). *Resting cysts of the toxigenic dinoflagellate genus Alexandrium in recent sediments from the Western Mediterranean coast, including the first description of cysts of A. kutnerae and A. peruvianum*

Bryars S (2003). An inventory of important coastal fisheries habitats in South Australia. Fish habitat Program, Primary Industries and Resources South Australia.

Bryars S and G Collins (2008). Causes of Adelaide's seagrass loss. In: S. Murray-Jones (Ed.) 'Proceedings of the Second Seagrass Restoration Workshop. Adelaide. April 2008.' Department for Environment and Heritage, Adelaide, pp. 18-37.

Bryars S and K Rowling (2009). *Benthic habitats of Eastern Gulf St Vincent: major changes in benthic cover and composition following European settlement of Adelaide*. Transactions of the Royal Society of South Australia 133(2): 318–338.

Bullock DA (1975) *The general water circulation of Spencer Gulf, South Australia, in the period February to May,* Transactions of the Royal Society of South Australia.

Burnell, S.R. (2001) Aspects of the reproductive biology, movements and site fidelity of right whales off Australia Journal of Cetacean Research and Management

Calvert SE and Peterson ML (1993) *Geochemistry of Recent oxic and anoxic marine sediments: Implications for the geological record,* Marine Geology.

Canfield DE (1989) Reactive iron in marine sediments, Geochimica et Cosmochimica Acta.

Cannon J A (1996) Competition between the dinoflagellates Alexandrium minutum and Prorocentrum micans in the Port River, South Australia

Carlson T J, Hastings M C and Popper A N (2007) *Update on recommendation for revised interim sound exposure criteria for fish during pile driving activities*. California Department of Transportation

Carroll E, Patenaude N, Alexander A, Steel D, Harcourt R, Childerhouse S, Smith S, Bannister J, Constantine R and Baker C (2011) *Population structure and individual movement of southern right whales around New Zealand and Australia*. Marine Ecology Progress Series 432, 257-268.



Carroll E L, Jackson J A, Paton D and Smith T D (2014). Two Intense Decades of 19 th Century Whaling Precipitated Rapid Decline of Right Whales around New Zealand and East Australia. PLoS ONE 9, e93789.

Caton B, Detmar S, Fotheringham D, Laurence S, Quinn J, Royal M, Rubbo N and Sandercock R (2011a) *Eyre Peninsula Coastal Action Plan and Conservation Priority Study, Volume 1*, Eyre Peninsula NRM Board and Department of Environment and Natural Resources, Adelaide.

Caton B, Detmar S, Fotheringham D, Laurence S, Quinn J, Royal M, Rubbo N, Sandercock R (2011b) *Eyre Peninsula Coastal Action Plan and Conservation Priority Study, Volume 2*, Eyre Peninsula NRM Board and Department of Environment and Natural Resources, Adelaide.

Codarin A, Wysocki LE, Ladich F, Picciulin M, (2009) *Effects of ambient and boat noise on hearing and communication in three fish species living in a marine protected area (Miramare, Italy)*, Marine Pollution Bulletin 58.

Collier CJ (2006) Characterising responses of the seagrass *Posidonia sinuosa* to changes in light availability. Ph.D. thesis, Faculty of Computing, Health and Science. Perth, Edith Cowan University.

Collier CJ, Lavery PS, Masini R and Ralph PJ (2007) *Morphological, growth and meadow characteristics of the seagrass* Posidonia sinuosa *along a depth-related gradient of light availability*. Mar.Ecol. Prog. Ser. 337, 103–115.

Collier, C.J., P.S. Lavery, R.J. Masini and P.J. Ralph, 2009. *Shade-induced response and recovery of the seagrass* Posidonia sinuosa. J. Exp. Mar. Biol. Ecol. 370: 89–103

Collings, G., Miller, D., O'Loughlin, E., Cheshire, A., and Bryars, S. (2006). *Turbidity and reduced light responses of the meadow forming seagrasses Amphibolis and Posidonia, from the Adelaide metropolitan coastline*. ACWS Technical Report No. 12 prepared for the Adelaide Coastal Waters Study Steering Committee. South Australian Research and Development Institute (Aquatic Sciences) Publication No. RD01/0208-17, Adelaide.

Commonwealth Scientific and Industrial Research Organisation (2001) Marine Pest Information Sheet: giant fanworm (Sabella spallanzanii)

Condon HT (1966) *The Western Whipbird. Preliminary notes on the discovery of a new subspecies on southern Yorke Peninsula, South Australia*, South Australian Ornithologist, Adelaide.

Conservation Council SA (2011) SA Marine Pests - Pest Species Introduced to and Established in SA. Accessed 15/01/2015 URL: http://feralperil.ala.org.au/bdrs-core/feralperil/fieldguide/taxon.htm?id=7729

Cornell RM, Schwertmann U (2003) *The iron oxides: Structure, properties, reactions, occurrences and uses.* Second Edn. Wiley-VCH: New York.

Costanza R, dArge R, deGroot R, Farber S, Grasso M, Hannon B, Limburg K, Naeem S, Oneill RV, Paruelo J, Raskin RG, Sutton P, vandenBelt M (1997) *The value of the world's ecosystem services and natural capital*, Nature.

Currie DR (2011) South Australian Giant Crab (Pseudocarcinus gigas) Fishery 2011/12. Fishery Status Report 2010/11. Status Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide.



Currie, D.R., Dixon, C.D., Roberts, S.D., Hooper, G.E., Sorokin, S.J. and Ward, T.M. (2009). *Fisheryindependent by-catch survey to inform risk assessment of the Spencer Gulf Prawn Trawl Fishery*. Report to PIRSA Fisheries. South Australian Research and Development Institute (Aquatic Sciences), Adelaide.

DeAngelis M, Fahy C and Cordaro J (2011) *Reducing vessel strikes of large whales in California. Report from a workshop held in Long Beach, California*; May 19-20, 2010. US Department of Commerce, National Oceanic and Atmospheric Administration, Mational Marine Fisheries Service, Southwest Regional Office. 36p DeBoar WF (2007) *Seagrass-sediment interactions, positive feedbacks and critical thresholds for occurrence: a review,* Hydrobiologia, Springer, Netherlands.

Dennison WC, Orth RJ, Moore KA, Stevenson JC, Carter V, Kollar S, Bergstrom PW and Batiuk RA (1993) *Assessing water quality with submerged aquatic vegetation*. Bioscience 43: 86-94.

Department of Agriculture, Forestries and Fishery (2011) Australian Ballast Water Management Requirements. Available from: http://www.daff.gov.au/biosecurity/avm/vessels/quarantine-concerns/ballast/australian-ballast-water-management-requirements

Department of Environment and Heritage (2003) *Focus: A Regional Perspective of Spencer Gulf.* Coast and Marine Branch, Department of Environment and Heritage, Adelaide, South Australia.

Department of the Environment and Heritage (DEH) (2005a). *Humpback Whale Recovery Plan 2005 - 2010*. [Online]. Department of the Environment and Heritage. Canberra, Commonwealth of Australia. Available from: http://www.environment.gov.au/resource/humpback-whale-recovery-plan-2005-2010

Department of the Environment and Heritage (DEH) (2005b). *Blue, Fin and Sei Whale Recovery Plan 2005 - 2010*. [Online]. Department of the Environment and Heritage. Canberra, Commonwealth of Australia. Available from: http://www.environment.gov.au/biodiversity/threatened/publications/recovery/balaenoptera-sp/index.html.

Department of the Environment and Heritage (DEH) (2006). Australian National Guidelines for Whale and Dolphin Watching 2005. [Online]. Department of the Environment and Heritage. Canberra, Commonwealth of Australia. Available from: http://www.environment.gov.au/marine/publications/australian-national-guidelines-whale-and-dolphin-watching-2005

Department of the Environment (2002) *National Ocean Disposal (NOD) Guidelines for Dredged Material,* Australian Government, Canberra.

Department of the Environment (2013) Significant Impact Guidelines 1.1 – Matters of National Environmental Significance. Accessed 20/01/2015. URL: http://www.environmental-significance

Department of the Environment, Water, Heritage and the Arts (2007) 2007 South-west Marine Bioregional Plan Bioregional Profile, Australian Government, Canberra.

Department of the Environment, Water, Heritage and the Arts (2008) *EPBC Act Policy Statement 2.1 – Interaction between offshore seismic exploration and whales*, Australian Government, Canberra.

Department of the Environment, Water, Heritage and the Arts (2009) *Threat abatement plan for the impacts of marine debris on vertebrate marine life, May 2009. Prepared for the Department of the Environment, Water, Heritage and the Arts (DEWHA).* available from

http://www.environment.gov.au/biodiversity/threatened/publications/tap/marine-debris.html



Department for Environment, Water, Heritage and the Arts (2010) Survey guidelines for Australia's threatened birds – Guidelines for detecting birds listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999, Australian Government, Canberra.

Department of Environment, Water and Natural Resources (2011) Native Vegetation (Floristic) - NVIS Statewide GIS data layer.

Department of Sustainability, Environment, Water, Population and Communities (2012) *Conservation Management Plan for the Southern Right Whale: A Recovery Plan under the Environment Protection and Biodiversity Conservation Act 1999, 2011-2021* Australian Government, Canberra.

Department of Sustainability, Environment, Water, Population and Communities (2013) *Issues Paper for the Australian Sea Lion (Neophoca cinerea) prepared for the Australian Government Department of Sustainability, Environment, Water, Population and Communities,* Australian Government, Canberra.

Ditmann, S. Cameron, S. Conlon, K (2010) *Increasing knowledge of introduced species in the marine environments of the Eyre Peninsula.* – Flinders University School of Biological sciences, Bedford park, SA

Dixon CD, Hooper, GE, and Burch P (2012) *Spencer Gulf Prawn Penaeus (Melicertus) latisulcatus Fishery 2010/11. Fishery Assessment Report to PIRSA Fisheries and Aquaculture.* South Australian Research and Development Institute (Aquatic Sciences), Adelaide.

Dixon CD, Noell CJ, Hooper GE, and Ward TM (2013) *Blue Crab (Portunus armatus) Fishery 2011/12. Fishery Assessment Report to PIRSA Fisheries and Aquaculture.* South Australian Research and Development Institute (Aquatic Sciences), Adelaide.

Dixon C D and Sloan S (2007) Management Plan for the South Australia Gulf St Vincent Prawn Fishery

Dooling RJ and Therrien SC (2012) Hearing in Birds: What Changes From Air to Water Springer New York

Duarte CM (1991). Seagrass depth limits. Aquatic Botany 40: 363-377.

Duarte, CM, (2002) The future of seagrass meadows, Environmental Conservation.

Duarte CM, Cebrian J (1996) The fate of marine autotrophic production, Limnology and Oceanography.

Duce RA, Tindale NW (1991) Atmospheric transport of iron and its deposition in the ocean, Limnology and Oceanography.

Edgar GJ (2001) Australian Marine Habitats in Temperate Waters, Reed Natural History of Australia.

Edgar GJ, Robertson AI (1992) *The influence of seagrass structure on the distribution and abundance of mobile epifauna: pattern and process in a Western Australian* Amphibolis *bed*. Journal of Experimental Marine Biology and Ecology.

Edgar GJ, Shaw C, Watson GF, Hammond LS (1994) *Comparisons of species richness, size-structure and production of benthos in vegetated and unvegetated habitats in Western Port, Victoria.* Journal of Experimental Marine Biology and Ecology

Edgar G (1997) Australian Marine Life: The Plants and Animals of Temperate Waters



Edyvane KS (1999a) Conserving marine biodiversity in South Australia, Part I: background, status and review of approach to marine biodiversity conservation in South Australia, South Australian Research and Development Institute, South Australia.

Edyvane KS (1999b) Conserving marine biodiversity in South Australia, Part II: identification of areas of high conservation value in South Australia, South Australia Research and Development Institute, South Australia.

Environment Protection Authority (2004) authors Corbin T and Wade S. *Heavy Metal concentrations in razorfish* (*Pinna bicolour*) and sediments across northern Spencer Gulf, Adelaide, South Australia.

Environment Protection Authority (2009) author Sam Gaylard. *Ambient water quality of Boston and Proper Bays, Port Lincoln 1997-2008.* Adelaide, South Australia.

Environment Protection and Biodiversity Conservation (2012) EPBC Act 1999 Protected Matters Search Tool located at the Australian Government Department of Sustainability, Environment, Water, Population and Communities website. http://www.environment.gov.au/epbc/pmst/index.html

Environment Protection and Biodiversity Conservation (2013) Western Whip-bird recovery plan accessed November 2013 http://www.environment.gov.au/system/files/pages/bb8ed384-16cb-411e-bd22-b1d380751030/files/recovery-outline-taxon-sum-birds-w.pdf

Ferguson G J (2013) *Pipi (Donax deltoides) Fishery.* Fishery Stock Assessment Report for PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences) Adelaide. SARDI Publication No. F2007/000550-1

Fernández A, Edwards JF, Rodríguez F, Monteros AE. de los Herráez P, Castro P, Jaber JR, Martín V, Arbelo M, (2005) "Gas and Fat Embolic Syndrome" Involving a Mass Stranding of Beaked Whales (Family Ziphiidae) Exposed to Anthropogenic Sonar Signals. Vet Pathology.

Filby NE, Bossley M, Sanderson KJ, Martinez E and Stockin KA (2010) *Distribution and population demographics of common dolphins (Delphinus delphis) in the Gulf of St. Vincent, South Australia.* Aquatic Mammals.

Fitzpatrick J and Kirkman H (1995) Effects of prolonged shading stress on growth and survival of seagrass Posidonia australis in Jervis Bay, New South Wales, Australia. Marine Ecology Progress Series, 127, 279-289.

Flinders Ports (2014) 12 month vessel records from December 2013 to December 2014, http://portmis.flindersports.com.au/ accessed 19 December 2014.

Fowler AJ, McGarvey R, Burch P, Feenstra JE, and Jackson WB (2010) *Snapper (Chrysophyrs auratus) Fishery.* Fishery Assessment Report to PIRSA. South Australian Research and Development Institute (Aquatic Sciences), Adelaide.

Fowler AJ, McGarvey R, Burch P, and Feenstra JE (2011) *King George Whiting (Sillaginodes punctatus) Fishery.* Fishery Assessment Report to PIRSA Fisheries and Aquaculture. South Australia Research and Development Institute (Aquatic Sciences), Adelaide.

Fowler AJ, McGarvey R, Steer MA and Feenstra JE (2012). South Australian Marine Scalefish Fishery – Stock Status Report 2011/12. Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide.



Gaillard JF, Webb SM (1998). *Biogeochemical Aspects of Metal-Microbes Interactions*. In: Transport, Fate and Effects of Silver in the Environment, 5th International Argentum Conference Proceedings, A. W. Andren and T. W. Bober, Eds. University of Wisconsin Sea Grant Publication No. WISCU-W-97001, 77-86.

Gammons CH and Frandsen AK (2001). Fate and Transport of metals in H2S-rich waters at a treatment wetland. Geochemical Transactions 2001 2:1.

Garcés E, Bravo I, Vila M, Figueroa R I, Masó M and Sampedro N, (2004) *Relationship between vegetative cells and cyst production during Alexandrium minutum bloom in Arenys de Mar harbour (NW Mediterranean)*

Gaylard S, Nelson M and Noble W (2013) *The South Australian monitoring, reporting and evaluation program* for aquatic ecosystems: rationale and methods for the assessment of nearshore marine waters Environment Protection Authority Adelaide

Geering A, Agnew L and Harding S (eds.) (2007) Shorebirds of Australia. CSIRO Publishing, Melbourne.

Gende S M, Hendrix A N, Harris K R, Eichenlaub B, Nielsen J and Pyare S (2011) A Bayesian approach for understanding the role of ship speed in whale-ship encounters. Ecological Applications 21, 2232-2240.

Gibbs S E and Kemper C M (2014) Whales and dolphins in Spencer Gulf. In Natural History of Spencer Gulf (Shepherd, S. A., Madigan, S. M., Gillanders, B. M., Murray-Jones, S. & Wiltshire, D. J., eds.), pp. 242-253. Adelaide: Royal Society of South Australia Inc.

Gillanders BM, Z Doubleday, P Cassey, SD Connell, M Deveney, S Dittman, S Divecha, M Doubell, S Goldsworthy, B Hayden, C Huveneers, C James, S Leterme, X Li, M Loo, J Luick, W Meyer, J Middleton, D Miller, L Moller, T Prowse, P Rogers, BD Russell, P van Ruth, JE Tanner, T Ward, SH Woodcock, M Young, (2013) Spencer Gulf Ecosystem & Development Initiative. Report on Scenario development, Stakeholder workshops, Existing knowledge & information gaps. Report for Spencer Gulf Ecosystem and Development Initiative. The University of Adelaide, Adelaide. 94 pages.

Global Invasive Species Database (2014) http://www.issg.org/database/welcome/ accessed on 12 February 2014

Gobler CJ, Donat JR, Consolvo JA and Sanudo-Wilhelmy SA (2002) *Physicochemical speciation of iron during coastal algal blooms.* Marine Chemistry.

Goldbogen JA, Southall BL, DeRuiter SL, Calambokidis J, Friedlaender AS, Hazen EL, Falcone EA, Schorr GS, Douglas A, Moretti DJ, Kyburg C, McKenna MF, Tyack PL (2013). *Blue whales respond to simulated mid-frequency military sonar*. Proceedings of the Royal Society 280, 20130657.

Golder Associates (2012) Winter 2011: Port Spencer Marine Baseline Quantitative Surveys. Prepared on behalf of Centrex Metals Ltd by Golder Associates Pty Ltd

Goldsworthy SD, and Page B (2009). *A Review of the Distribution of Seals in South Australia*. South Australian Research and Development Institute (Aquatic Sciences), Adelaide, SARDI Publication Number F2009/000368-1, 21 pp. SARDI Research Report Series No. 373.

Goldsworthy SD, Peters KJ and Page B (2007). <u>Foraging ecology and diet analysis of Australian sea lions.</u> Final Report to the Department of Environment and Water Resources. SARDI Aquatic Sciences Publication Number F2007/001024-1. SARDI Research Report Series No. 251

Gomon M, Bray D and Kuiter R (2008) *Fishes of Australia's southern coast*. Museum Victoria, Melbourne
E-F-34-RPT-0039_A (Marine Environmental Tech Report_Rev1a - compared doc 24/09/2015 Page 183 of 204



Goold JC, Fish PJ, (1998) *Broadband spectra of seismic survey air-gun emissions, with reference to dolphin auditory thresholds.* The Journal of the Acoustical Society of America 103, 2177–2184.

Gordon DM, Grey KA, Chase SC and CJ Simpson (1994) *Changes to the structure and productivity of a* Posidonia sinuosa *meadow during and after imposed shading*. Aquatic Botany, 47, 265-275.

Gostlin VA, Halis JR and Belperio AP (1984) *The sedimentary framework of northern Spencer Gulf, South Australia*. Marine Geology, 61(2-4):111-138.

Government of South Australia (1998) Fisheries Management (Blue Crab Fishery) Regulations 1998 [CEASED] South Australian Attorney-General's Department, Adelaide.

Hallegraeff G M, Steffensen D A and Wetherbee R (1988) *Three estuarine Australian dinoflagellates that can produce paralytic shellfish poison*

Hallegraeff G M (1993) A review of harmful algal blooms and their apparent global increase

Harvey N, Caton B (2010) Coastal Management in Australia, ed. Dragovitch and Maude p28

Harvey N, Clouston B, Carvalho P (1995) *Coastal Vulnerability Study: Northern Spencer Gulf, South Australia*, Mawson Graduate Centre for Environmental Studies, University of Adelaide, South Australia.

Hastings, M and Popper, A (2005) Effects of sound on fish. California Department of Transportation.

Hazel J, Lawler IR, Marsh H and S Robson (2007) *Vessel speed increases collision risk for the green turtle* Chelonia mydas. Endangered Species Research. 3: 105-113.

Heck KL, Hays G, and Orth RJ (2003) *Critical evaluation of the nursery role hypothesis for seagrass meadows.* Marine Ecology Progress Series 253: 123-136

Holmer M, Duarte CM, Marba N (2005) *Iron additions reduce sulphate reduction rates and improve seagrass growth on organic-enriched carbonate sediments*. Ecosystems.

Hu, M, Yan, H, Chung, W, Shial, J, and Hwang, P (2009) *Acoustically evoked potentials in two cephalopods inferred using the auditory brainstem response (ABR) approach*. Comparative Biochemistry and Physiology, Vol. 153 Part A, pp. 278–283.

Hutchings PA, Ward T, Waterhouse JH, and Walker L (1993) *Infauna of marine sediments and seagrass beds of upper Spencer Gulf near Port Pirie, South Australia.* Transactions of the Royal Society of South Australia.

Hutchins D.A. and K.W. Bruland (1998) *Iron-limited diatom growth and Si:N uptake ratios in a coastal upwelling regime.* Nature.

Hutchins DA, Witter AE, Butler A and Luther III GW (1999) Competition among marine phytoplankton for different chelated iron species

Hydro Survey (2012) Cape Hardy Marine Investigation Project Appendix A

International Union for Conservation of Nature (2013) The ICUN Red List of threatened species online database accessed at http://www.iucnredlist.org/ November 2013



Inshore Fish Group (2008) viewed 16 February 2015 at http://www.ifg.bioteck.org/Fish%20species/

IOSEA (2013) Indian Ocean – South-East Asia Marine Turtle Memorandum of Understanding: species database and assessments http://www.ioseaturtles.org/ accessed November 2013

Jackson EL, Rowden AA, Attrill MJ, Bossey SJ, and Jones MB (2001) *The importance of seagrass beds as a habitat for fisheries species.* Oceanography and Marine Biology: An Annual Review

Jacobs (2014a) Central Eyre Iron Project: Port Site Ecological Survey report number E-F-34-RPT-0015 prepared for Iron Road Ltd by Jacobs.

Jacobs (2014b) Central Eyre Iron Project: Infrastructure Air Quality Impact Assessment report number E-F-34-RPT-0040 prepared for Iron Road Ltd by Jacobs.

Jacobs (2014c) Central Eyre Iron Project: Environmental Noise and Vibration Assessment – Cape Hardy Port Facility report number E-F-34-RPT-0038 prepared for Iron Road Ltd by Jacobs.

Jacobs (2014d) Cape Hardy Coastal Modelling Report report number E-F-34-RPT-0030 prepared for Iron Road Ltd by Jacobs

Jacobs SWL and Les DH (2009) *New combinations in Zostera (Zosteraceae)*. Royal Botanic Gardens and Domain Trust. Telopea 12(3) 419-423

Kailola P, Williams M, Stewart P, Reichelt R, McNee A, Grieve C (1993) *Australian Fisheries Resources*, Bureau of Resource Sciences and the Fisheries Research and Development Corporation, Canberra.

Kersten M (1988) Geochemistry of Priority Pollutants in Anoxic Sludges: Cadmium, Arsenic, Methyl Mercury and Chlorinated Organics. In Chemistry and Biology of Solid Waste ed. Salomons and Forstner p189-195

Kemper CM (2005) *Records of humpback whales* Megaptera novaeangliae *in South Australia*, Transactions of the Royal Society of South Australia, Adelaide.

Kemper C, Coughran D, Warneke R, Pirzl R, Watson M, Gales R and Gibbs S (2008) *Southern right whale* (Eubalaena australis) *mortalities and human interactions in Australia, 1950-2006*. Journal of Cetacean Research and Management 10, 1-8.

Kemper, C. and Ling, J. (1991) Whale strandings in South Australia (1881 – 1989), Transactions of the Royal Society of South Australia, 115: 37-52.

Kite-Powell HK, Knowlton NA and M Brown (2007). *Modelling the effect of vessel speed on right whale ship strike risk*. Project report for NOAA/NMFS Project NAO4NMF47202394, April 2007.

Knight M and Tsolos A (2012) South Australian Wild Fisheries Information and Statistics Report 2010/2011. South Australian Research and Development Institute (Aquatic Sciences), Adelaide.

Kostoglou P and McCarthy J (1991) Whaling and Sealing Sites in South Australia, Australian Institute for Maritime Archaeology

Kuo J (2005) A revision of the genus Heterozostera (Zosteraceae). Aquatic Botany 81: 97–140.

Laist D W, Knowlton A R, Mead J G, Collet A S and Podesta M (2001) *Collisions between ships and whales*. Marine Mammal Science 17, 35-75.



Langston WJ (1984) Availability of arsenic to estuarine and marine organisms: a field and laboratory evaluation. Marine Biology 80: 143-154

Lennon GW, Bowers GD, Nunes RA, Scott BD, Ali M, Boyle J, Wenju C, Herzfeld M, Johansson G, Nield S, Petrusevics P, Stephenson P, Suskin AA, and Wijffels SEA (1987) *Gravity currents and the release of salt from an inverse estuary,* Nature.

Levings A, Mitchell BD, McGarvey R, Mathews J, Laurenson L, Austin C, Heeron T, Murphy N, Miller A, Rowsell M, Jones P (2001) *Fisheries biology of the giant crab,* Pseudocarcinus gigas, Final report to the fisheries research and development corporation for projects 93/220 & 97/132, School of Ecology and Environment, Deakin University, Geelong.

Lim P, Leaw C, Usup G, Kobiyama A, Koike K and Ogata T (2006) Effects of Light and Temperature on Growth, Nitrate Uptake, and Toxin Production of two Tropical Dinoflagellates: Alexandrium tamiyavanichii and Alexandrium minutum (Dinophyceae)

Linnane A, McGarvey R, Feenstra J and Hoare M (2012) *Northern Zone Rock Lobster* Jasus edwardsii *Fishery Status Report 2011/12. Status Report to PIRSA Fisheries and Aquaculture.* South Australian Research and Development Institute (Aquatic Sciences), Adelaide.

Llansó RJ (2002) *Methods for Calculating the Chesapeake Bay Benthic Index of Biotic Integrity.* Versar, Inc., 9200 Rumsey Road, Columbia, MD 21045, Revised 11 November 2002. Available at: http://www.baybenthos.versar.com

Long ER, Macdonald DD, Smith SL, and Calder FD (1995) *Incidence of adverse biological effects within ranges of chemical concentrations in marine and estuarine sediments.* 19: 81-97.

Luoma SN, Bryan GW, Langston WJ (1982) *Scavenging of Heavy Metals from Particulates by Brown Seaweed,* Marine Pollution Bulletin 13: 394-396

McConnaughey RA, Syrjala S (2008) *Appendix 2: Ecosystems Considerations Chapter*, Alaska Marine Ecosystems Considerations REEM ed. Heifetz J. Available at http://access.afsc.noaa.gov/reem/EcoWeb/EcoChaptMainFrame.cfm?ID=90

Madden-Hallett DM, Hammer M, Gursanky W, Donato DB (2011) *Lipson Island baseline flora and fauna report and assessment of risk.* For Golder Associates. Donato Environmental Services, Darwin

MARPOL, International Convention for the Prevention of Pollution from Ships http://www.imo.org/About/Conventions/ListOfConventions/Pages/International-Convention-for-the-Prevention-of-Pollution-from-Ships-(MARPOL).aspx

Martin JH, Coale KH, Johnson KS, Fitzwater SE, Gordon RM, Tanner SJ, Hunter CN, Elrod VA, Nowicki JL, Coley TL, Barber RT, Lindley S, Watson AJ, Van Scoy K, Law CS, Liddicoat MI, Ling R, Stanton T, Stockel J, Collins C, Anderson A, Bidigare R, Ondrusek M, Latasa M, Millero FJ, Lee K, Yao W, Zhang JZ, Friederich G, Sakamoto C, Chavez F, Buck K, Kolber Z, Greene R, Falkowski P, Chisholm SW, Hoge F, Swift R, Yungel J, Turner S, Nightingale P, Hatton A, Liss P, and Tindale NW (1994) *Testing the iron hypothesis in ecosystems of the equatorial Pacific Ocean*, Nature 371, 123 - 129

Masini RJ, Burt JS and Simpson CJ (1995) *Light and Posidonia seagrass meadows in the temperate coastal waters of Western Australia. III Minimum light requirements*. Technical Series No. 65, Perth, Western Australia: Department of Environmental Protection.



MES (2012) Central Eyre Iron Project – Geophysical Surveys, Cape Hardy, South Australia prepared by Marine & Earth Sciences for Sinclair Knight Merz Adelaide Office.

Middleton J, Doubell M, James C, Luick J and van Ruth P (2013) *PIRSA Initiative II: carrying capacity of Spencer Gulf: hydrodynamic and biogeochemical measurement modelling and performance monitoring*. Final Report for the Fisheries Research and Development Corporation. South Australian Research and Development Institute (Aquatic Sciences), Adeliade. SARDI Publication No. F2013/000311-1. SARDI Research Report Series No. 705. 97pp.

Möller L M, Bolgmann K, Charlton-Robb K and Beheregaray L (2008) *Multi-gene evidence for a new bottlenose dolphin species in southern Australia*. Molecular Phylogenetics and Evolution

Mooney TA, Hanlon R, Madsen PT, Christensen-Dalsgaard J, Ketten DR and Nachtigall PE (2012) *Potential for sound sensitivity in cephalopods*

Montgomery JC, Jeffs A, Simpson SD, Meekan M, Tindle C, (2006) *Sound as an Orientation Cue for the Pelagic Larvae of Reef Fishes and Decapod Crustaceans*, in: Alan J. Southward, and D.W.S. (Ed.), *Advances in Marine Biology*. Academic Press, pp. 143–196.

Myrberg AA (1990) *The effects of man-made noise on the behaviour of marine animals*, Environment International.

National Assessment Guidelines for Dredging (2009), Commonwealth of Australia, Canberra

National Oceanic and Atmospheric Administration (2007) *Southern Right Whale 5-Year Review: Summary and Evaluation* National Marine Fisheries Services, Office of Protected Resources

National Oceanic and Atmospheric Administration (2013) *Draft Guidance for Assessing the Effects of Anthropogenic Sound on Marine Mammals – Acoustic Threshold Levels for Onset of Permanent and Temporary Threshold Shifts*, U.S. Department of Commerce

Nature Maps http://www.naturemaps.sa.gov.au/ accessed in March 2014

NIMPIS (2014) The Australian Government, NIMPIS – the National Introduced Marine Pest Information System http://data.daff.gov.au/marinepests/ accessed March 2014

O'Brien A L, Ross D J and Keough M J (2006) Effects of Sabella spallanzanii physical structure on soft sediment macrofaunal assemblages. Marine Freshwater and Research

Organisation for Economic Co-operation and Development (1997) *The environmental effects of freight*, OECD, Paris.

Online Zoological Collections of Australian Museums (2012) Online Zoological Collections of Australian Museums at http://www.ozcam.org.au/

Page B, Peters KD, McIntosh RR, Hamer D, Baylis AMM, Summer MD, McGarvey R, and Goldsworthy SD (2009) Assessment of different home range estimates and spatial scales to describe the distribution of Australian Sea Lion foraging effort. In. Part B: Goldsworthy, SD, Page, B, Shaughnessy, PD, Hamer, D, Peters, KD, McIntosh, RR, Baylis, AMM, McKenzie, J. (2009) Innovative solutions for aquaculture planning and



management: addressing seal interactions in the finfish aquaculture industry. SARDI Aquatic Sciences Publication Number F2008/000222–1. SARDI Research Report Series Number 288

Peterson ML and Carpenter R (1986). Biogeochemical processes affecting total arsenic and arsenic species distribution in an intermittently anoxic fjord. *Marine Chemistry* 12: 295-321

PIRSA (2008) Primary Industries and Regions South Australia (PIRSA) Report supporting the Aquaculture (zones – Port Neill) Policy 2008. Available as a pdf from the PIRSA website http://www.pir.sa.gov.au/aquaculture/management policies/zonal policies

PIRSA (2014) Primary Industries and Regions South Australia (PIRSA) website accessed November 2014 http://www.pir.sa.gov.au/fisheries/recreational_fishing/protected_species

Primary Industries and Regions South Australia (2014) PIRSA, Aquaculture lease public register database accessed May 2014 http://www.pir.sa.gov.au/aquaculture/public register

Poore GCB (1995) *Biogeography and diversity of Australia's marine biota*. In 'The State of the Marine Environment Report for Australia. *Technical Annex 1: The Marine Environment* (Ed. L Zann) Department of Environment, Sport and Territories, Canberra, pp. 75-84.

Poore, GCB and O'Hara, TD (Eds) (2007) *Marine biogeography and biodiversity of Australia*. In 'Marine Ecology'. (Oxford University Press: Sydney).

Popper, A and Hastings, M (2009) REVIEW PAPER: *The effects of anthropogenic sources of sound on fishes*. Journal of Fish Biology, vol. 75, pp 455-489.

Port Bonython (2013) Spencer Gulf Port Link – Port Bonython Bulk Commodities Export Facility Environmental Impact Statement prepared by Spencer Gulf Port Link and Arup Pty Ltd.

Port Spencer (2012) *Port Spencer Stage 1 Public Environmental Report* prepared by Golder Associates Pty Ltd on behalf of Centrex Metals Ltd.

Primary Industries and Resources SA (2008) *Report Supporting the Aquqculture (Zones – Port Neill) Policy 2008*. Government of South Australia

Rabalais NN (2002) Nitrogen in Aquatic Ecosystems, Journal of the Human Environment 31(2): 102-112.

Ravenscroft P, McArthur JM, and Hoque BA (2001) *Geochemical and paleohydrological on pollution of groundwater by arsenic*. In W.R. Chappell *et al.* (eds), *Arsenic exposure and health effects IV*. Elsevier, Amsterdam.

Reynoldson TB (1987) *Interactions between sediment contaminants and benthic organisms*, Hydrobiologia 149: 53

Richardson WJ, Würsig B (1997) *Influences of man-made noise and other human actions on cetacean behaviour*. Marine and Freshwater Behaviour and Physiology 29, 183–209.

Richardson WJ, Würsig B, Greene CR (1990) *Reactions of bowhead whales, Balaena mysticetus, to drilling and dredging noise in the Canadian Beaufort Sea.* Marine Environmental Research 29, 135–160.



Richardson WJ, Würsig B (1986) *Reactions of bowhead whales, Balaenamysticetus, to seismic exploration in the Canadian Beaufort Sea.* The Journal of the Acoustical Society of America 79, 1117–1128.

Ruiz-Halpern S, Macko SA and JW Fourqurean (2008) *The effects of manipulation of sedimentary iron and organic matter on sediment biogeochemistry and seagrasses in a subtropical carbonate environment.*Biogeochemistry. 87:113-126.

Russell BD, Elsdon TS, Gillanders BM, and Connell SD (2005). *Nutrients increase epiphyte loads: broad-scale observations and an experimental assessment.* Marine Biology 147: 551-558.

Seddon S (2000) Causes and ecological consequences of the Spencer Gulf seagrass dieback. PhD dissertation, University of Adelaide. 171 pp.

Seddon S, Connolly RM, and Edyvane KS (2000) *Large-scale seagrass dieback in northern Spencer Gulf, South Australia*. Aquatic Botany 66:297-310.

Seraux C, Robinson-Laverick SM, Maho YL, Ropert-Coudert Y and Charadia A (2011) *Plasticity in foraging strategies of inshore birds: how little Penguins maintain body reserves while feeding offspring.* Ecology 92(10):1909-1916

Shanks S (2005) *Management Plan for the South Australian Pilchard Fishery* Fisheries Division, PIRSA, South Australian Government Paper No. 47.

Shaughnessy PD (1999) *The Action Plan for Australian Seals.* CSIRO Wildlife and Ecology produced for the Australian Government Natural Heritage Trust, Canberra.

Shaughnessy PD, Dennis TE, and Seager PG (2005) *Status of Australian sea lions*, Neophoca cinerea, *and New Zealand fur seals*, Arctocephalus forsteri, *on Eyre Peninsula and the Far West Coast of South Australia*. Wildlife Research 32:85-101

Shaughnessy PD, Goldsworthy SD, Hamer DJ, Page B, and McIntosh RR (2011) *Australian sea lions* Neophoca cinerea *at colonies in South Australia: distribution and abundance* , *2004 to 2008*. Endangered Species Research 13:87-98

Shepherd SA (1983) *Benthic communities of upper Spencer Gulf, South Australia*. Transactions of the Royal Society of South Australia 107(2):69-85.

Shepherd SA and Robertson EL (1989) *Regional studies: Seagrasses of South Australia, Western Vicoria and Bass Strait.* In *Biology of Seagrasses: A treatise on the biology of seagrasses with special reference to the Australian region* (Eds. AWD Larkham, AJ McComb and SA Shepherd) 211-229. (Elsevier, Amsterdam).

Shirihai H (2007) A Complete Guide to Antarctic Wildlife: the Birds and Marine Mammals of the Antarctic Continent and the Southern Ocean (2^{nd} edit.) A & C Black, London.

Silber G K, Slutsky J and Bettridge S (2010) *Hydrodynamics of a ship/whale collision*. Journal of Experimental Marine Biology and Ecology 391, 10-19.

Simpson K and Day N (1999) Field Guide to the Birds of Australia (6th Edition), Penguin Books Australia.

Simpson K and Day N (2004) Field Guide to the Birds of Australia (7th Edition), Penguin Books Australia.



Sinclair Knight Merz (2010) *Port Options Briefing Document, Central Eyre Iron Project*. Report prepared for Iron Road Limited by Sinclair Knight Merz, Adelaide and Melbourne Offices Australia.

Sinclair Knight Merz (2011) Lucky Bay common user export facility marine investigations. Report prepared for Iron Clad Mining Ltd, by Sinclair Knight Merz Adelaide Office, Australia

Sloan S (2002) A Report Prepared for Environment Australia on the Management of the South Australian Giant Crab (Pseudocarcinus gigas) Fishery. Report for PIRSA available online at: http://www.pir.sa.gov.au/ data/assets/pdf file/0006/12777/giant crab2002.pdf

Slabbekoorn H, Bouton N, van Opzeeland I, Coers A, ten Cate C, Popper AN (2010) A noisy spring: the impact of globally rising underwater sound levels on fish, Trends in Ecology and Evolution.

Sloan S and Crosthwaite K (2007) *Management plan for the South Australian northern zone rock lobster fishery* PIRSA Fisheries publication

Smedley PL and Kinniburgh DG (2002) *A review of the source, behaviour and distribution of arsenic in natural waters.* Applied Geochemistry 17: 517-568.

Sonus (2014) Central Eyre Iron Project; Marine Environmental Noise Assessment. Prepared for Jacobs Adelaide South Australia.

Southall B L (2005) *Shipping Noise and Marine Mammals: a Forum for Science, Management, and Technology.* Final Report of the National Oceanic and Atmospheric Administration (NOAA) International Symposium

SPRAT (2014) Species Profile and Threats Database http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl for EPBC listed species online fact sheets accessed November 2014

Steer MA, McGarvey R, Fowler AJ, Burch P, Feenstra JE, Jackson WB and Lloyd MT (2012) *Southern Garfish* (Hyporhamphus melanonchir) *Fishery. Report to PIRSA Fisheries and Aquaculture*. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/000720-3. SARDI Research Report Series No. 658. 76pp.

Street JH, Paytan A (2005) *Iron, phytoplankton growth, and the carbon cycle.* Metals Ions in Biological Systems. 43:153-193

Stobart, B. (2014). South Australian Giant Crab (Pseudocarcinus gigas) Fishery Status Report 2012/13. Fishery Status Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2011/000332-4. SARDI Research Report Series No. 763. 15pp.

Stobart B, Mayfield S, Dent J, Matthews DJ and Chick RC (2012) *Western Zone Abalone* (Haliotis rubra & H. laevigata) *Fishery (Region A) – Fishery Stock Assessment Report to PIRSA Fisheries and Aquaculture.* South Australian Research and Development Institute (Aquatic Sciences), Adelaide. Publication No. F2007/000561-4, SARDI Research Report Series No. 660. 118 pp.

Stoner AW (1980) *The Role of Seagrass Biomass in the Organization of Benthic Macrofaunal Assemblages.* Bulletin of Marine Science 30: 537–551.

Tanner JE (2003) *Patch shape and orientation influences on seagrass epifauna are mediated by dispersal abilities.* Oikos 100, 517–524.



Thompson PA, Butler ECV and RJ Watson (2008) A review of the role of iron, and the possible ramifications of its enhancement, in the phytoplankton ecology of lower Spencer Gulf, South Australia. CSIRO Marine and Atmospheric Research, 31 October 2008.

Ussher SJ, Achterberg EP, Worsfold PJ (2004) *Marine biogeochemistry of iron.* Environmental Chemistry. 1: 67-80

Usup G, Pin L C, Ahmad A and Teen L P (2002). *Alexandrium (Dinophyceae) species in Malaysian waters*. Harmful Algae 1

Valenzuela LO, Sironi M, Rowntree VJ and J Seger (2009) *Isotopic and genetic evidence for culturally inherited site fidelity to feeding grounds in southern right whales* (Eubalaena australis) Molecular Ecology 18: 782–791

van der Hoop J M, Moore M J, Barco S G, Cole T V N, Daoust P Y, Henry A G, McAlpine D F, McLellan W A, Wimmer T and Solow A R (2013) *Assessment of management to mitigate anthropogenic effects on large whales*. Conservation Biology 27, 121-133.

van der Hoop J M, Vanderlaan A S M and C T Taggart (2012) Absolute probability estimates of lethal vesselstrikes to North Atlantic right whales in Roseway Basin, Scotian Shelf. Ecological Applications 22(7):2012-2033

Vanderlaan, A. S. M., and C. T. Taggart. (2007). *Vessel collisions with whales: the probability of lethal injury based on vessel speed*. Marine Mammal Science 23:144–156.

Vila M, Grazia Giacobbeb M, Masoa M, Gangemib E, Pennac A, Sampedroa N, Azzarob F, Campa J and Galluzzic L (2005) *A comparative study on recurrent blooms of Alexandrium minutum in two Mediterranean coastal areas*. Harmful Algae 4

Walker DI and McComb AJ (1992) *Seagrass degradation in Australian coastal waters*. Marine Pollution Bulletin 25: 191-195.

Ward TJ (1987) *Temporal variation of metals in the seagrass* Posidonia australias *and its potential as a sentinel accumulator near a lead smelter.* Marine Biology 95: 315-321.

Ward TJ and Hutchings PA (1996) Effects of trace metals on infaunal species composition in polluted intertidal and subtidal marine sediments near a lead smelter, Spencer Gulf, South Australia. Marine Ecology Progress Series 135:123-135.

Ward, T.M. and Burch, P. (2012). Revised estimates of historical commercial fishery catches/effort in draft sanctuary and habitat protection zones in South Australia's Marine Parks. Report to PIRSA Fisheries and Aquaculture. South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2011/000307-6. SARDI Research Report Series No. 642. 120pp.

Ward TM, Burch P and Ivey AR (2012) *South Australia Sardine* (Sardinops sagax) *Fishery: Stock Assessment Report 2012. Report to PIRSA Fisheries and Aquaculture.* South Australian Research and Development Institute (Aquatic Sciences), Adelaide. SARDI Publication No. F2007/000765-4. SARDI Research Series No. 667. 101pp.

Wardle C, Carter T, Urquhart G, Johnstone AD, Ziolkowski A, Hampson G, Mackie D (2001) *Effects of seismic air guns on marine fish*. Continental Shelf Research 21, 1005–1027.

Waring J and Maher W (2005) *Arsenic bioaccumulation and species in marine Polychaeta (Review)*. Applied Organometallic Chemistry 19: 917-929



Westphalen G, Collings G, Wear R, Fernandes M, Bryars S and Cheshire A (2004) *A review of seagrass loss on the Adelaide metropolitan coastline*, ACWS Technical Report No. 2 prepared for the Adelaide Coastal Waters Study Steering Committee, South Australian Research and Development Institute

Wiltshire KH (2010) *Caulerpa taxifolia* – 2010 survey of current distribution and high risk areas and summary of distribution patterns 2003-2010. SARDI Aquatic Sciences, Adelaide Publication No. F2010/000612-1 SARDI Report Series No. 474

Wiltshire KH and Deveney MR (2011) Final monitoring report for the 2010-2011 marine pest survey of the Port Adelaide, South Australia. South Australian Research and Development Institute (Aquatic Sciences), Adelaide, SARDI Publication No. F2011/000386-1. SARDI Report Series No. 575

Wiltshire, K. Rowling, K. and Deveney, M. 2010. *Introduced marine species in South Australia: a review of records and distribution mapping*. SARDI

Yoshida M, Ogata T, Thuoc C V, Matsuoka K, Fukuyo Y, Hoi N C and Kodama M (2000) *The first finding of toxic dinoflagellate Alexandrium minutum in Vietnam*

ZooSA (2013) http://www.zoossa.com.au/conservation-ark/research/past-projects website accessed November 2013



Appendix A Bathymetry Survey Reports

AUSTRALIAN HYDROGRAPHIC SERVICE



SURVEY SUMMARY

INSTRUCTIONS FOR RENDERING

This Survey Summary form should be completed for all data rendered to the Australian Hydrographic Office which is not accompanied by a full written report, Method Statement, Plans or other reports which would normally include details such as those in this form.

This will provide the minimum information required to manage data within Australia's area of charting interest.

The preferred format of bathymetric data is:

- Processed,
- Ungridded for singlebeam and multibeam, shoal biased, true position,
- Provided as either GSF, ascii .xyz, .dxf, .dgn, or Hydrographic Transfer Format (.htf, available from the AHS website http://www.hydro.gov.au). If these formats are not available, full source data will be accepted.

Any ancillary data such as tides, benchmarks, linework and final levelling heights etc. is also of assistance. If supplying such data, please include positional data of deployed equipment.

Please forward survey data with the completed Survey Summary to:

Hydrographer of Australia C/- Manager Digital Information Australian Hydrographic Office Locked Bag 8801 Wollongong NSW 2500

Guidance on Confidence Levels and Error Ellipse scaling is contained in ICSM Publication Number 1 (SP-1), uncertainties from IHO publication S-44 or by contacting the Validation and Assessment Section at the Australian Hydrographic Office on 02 4223 6500.

SURVEY SUMMARY

General

| Survey Title and ID | Locality | | | |
|---|--|--|--|--|
| Cape Hardy Soundings | Spencer Gulf, South Australia | | | |
| Survey Authority | Survey Sponsor/Custodian | | | |
| HydroSurvey Australia | Flinders Ports South Australia Pty Ltd | | | |
| Surveyor in Charge and Qualification | Date this Survey Summary was completed | | | |
| Peter Woolfall (Level 1) | 23 rd March 2012 | | | |
| Start Date of Survey | End Date of Survey | | | |
| 18 th February 2012 | 23 rd February 2012 | | | |
| Survey Platform/Vessel Name | Scale (if applicable) | | | |
| S.V. Pathfinder | 5000 | | | |
| Purpose of the Survey | | | | |
| Cape Hardy Marine Investigation Project | | | | |
| List of Data Provided | | | | |
| 2 x csv files, 3 x dwg file | | | | |
| Do you require the media returned and to which address? | | | | |

Horizontal Control

| Soundings are on the following datum (WGS 84 preferred by not essential). | | | | | | |
|--|-------|-----|-------------|-------------|-------|----|
| Datum | GDA94 | | Spheroid | GRS80 | GRS80 | |
| Projection | MGA94 | | Zone | 53 South | | |
| Was the positioning system validated? | | Yes | Were laybac | ks applied? | • | No |
| Estimated horizontal accuracy of soundings at 2σ (95%) confidence level (Calculations can be included as an attachment) +/- 1 metres | | | | | | |

Vertical Control

| Tides applied | Soundings corrected using Observed Tides | | Sounding Datum | Indian S ₁ | prings Low Water |
|---|--|---|----------------|-----------------------|------------------|
| Tide Statio | n Details | Tumby Bay (34°22'39"S 136°06'21'E) Cape Hardy (34°11'56"S 136°18'00'E) | | | |
| and height | (BM) used difference M and Datum | CD(approx ISLW) is 3.885 BM PSMP = EL 2.658m A | | chmark PS | SMP 6129/1003. |
| Geoid detail GPS tides | | n/a | | | |
| Were sound | dings corrected | for draught? | | | Yes |
| Estimated vertical accuracy of soundings at 1.96σ (95 (Calculations can be included as an attachment) | | | %) confidenc | e level | +/- 0.39 metres |

Details of Survey Execution

| The following positioning systems were used | |
|---|--|
| Positioning System 1 Trimble DSM232 DGPS | |

| Positioning System 2 | POS MV Wavemaster V4 | | | | |
|---------------------------------|---|---------------|--|--|--|
| Base Station | OmniStar ITRF to GDA94 corrections (2012.5 epoch) | | | | |
| The following sounding | The following sounding systems were used | | | | |
| | Model / System Details Frequency | | | | |
| Echosounder 1 | GeoSwath Plus MBES | 250 kHz | | | |
| Echosounder 2 | Odom Echotrac Mk2 200 / 24 kHz | | | | |
| Motion Reference Unit | POS MV Wavemaster V4 RM IP68 | | | | |
| Towed Side Scan Sonar | n/a kHz | | | | |
| Hull Mounted Side Scan Sonar | n/a kHz | | | | |
| | Model / System Details | Spot density | | | |
| LIDAR | n/a | X by X metres | | | |
| Sweep System | n/a | | | | |

| Logging and Processing Systems used Version | | | | | |
|---|---|-------|-----------|--------------|--|
| Logging | GeoSwath Plus / Hyd | roPro | 3.50r / 2 | 2.40 | |
| Processing | GeoSwath Plus / HydroPro / Terramodel 3.50r / 2 | | | 2.40 / 10.61 | |
| Was the survey | Was the survey systematically controlled with planned survey lines or methods? Yes | | | | |
| Were soundings corrected for sound velocity? Yes | | | | | |
| Was full feature detection achieved as defined in IHO publication S-44? Yes | | | | | |
| Were all shoal depths systematically investigated and their least depths determined? Yes | | | | | |
| Has the data be | Has the data been thinned from that collected? Yes | | | | |
| If thinned, what thinning method and bin size was used? 2m bin out of GS+, thinned to 45m circle of influence in Terramodel (shoal bias) for plotting purposes (1 to 5000) | | | | | |
| If thinned, have least depths and their positions been retained? Yes | | | | | |

| Remarks (If applicable): | |
|--------------------------|--|
| | |

Shoals and Dangers

This section seeks comment on any features that may be dangerous to surface navigation.

Small Yellow Buoy located at 34°11'22.9"S 136°19'55.4E



Appendix B Sediment Analysis Results



B1. Particle Size Distribution Lab Results



Laser Diffraction particle sizing of 15 Marine Sediment Samples

Project Number 2012/005 **Ian Wark Research Institute ARC Special Research Centre**

Minerals and Materials Science -Nanotechnology - Interfaces

Director

Professor John Ralston AO FAA FTSE

Deputy Director Professor Hans Griesser

Mawson Lakes Campus Mawson Lakes Adelaide SA 5095 Australia

t +61 8 8302 3694 f +61 8 8302 3683

www.unisa.edu.au/iwri

Australian Research Council Special Research Centre for Particle and Material Interfaces

CRICOS Provider Number 00121B

Attention Dr Greg Barbara

Prepared by

Prepared for

Dr Scott Abbott

Sinclair Knight Merz

Adelaide SA 5000

L5, 33 King William St

Date of issue

18 January 2012

Distribution Sinclair Knight Merz

Ian Wark Research Institute:

Scientific Services

Important Notice

This report applies only to the subject of the project.

This report is confidential and was prepared exclusively for the client named above. It is not intended for, nor does the University of South Australia accept any responsibility for its use by any third party. This report consists of one cover page and thirty four (total 35) pages of text and figures plus an attached spreadsheet of data.

<u>Introduction</u>

15 marine sediment samples were received at the Ian Wark Research Institute (commonly referred to as The WarkTM) on Friday 13th January 2012 for laser particle size distribution (PSD) analysis. A purchase order complete with number was received on the same day. The supplied samples were labelled as shown in Table 1.

Experimental

Before analysis, the samples were wet sieved using a 2,000µm mesh sieve. Significant observations made during the sieving process are also summarised in Table 1. Particle size distribution of the -2,000µm fraction was then measured using a Malvern Mastersizer 2000 Static Light Scattering/Laser diffraction instrument covering the measurement range of 0.02 to 2,000µm. Three measurements of each sample were conducted for 20 seconds and these 3 results for each sample were averaged. The exception was sample C6. The majority of the particulate in sample C6 was very coarse (>1mm) and as such tended to 'drop-out' and settle in the Mastersizer feed tube. As such, only one measurement could be recorded before this settling was observed.

Results

Samples consisted of a variety of forms from medium sand through to coarse shell fragments, with a wide range of larger material, including some small rocks.

The measured volume % PSD plots (Figure 1 and Figure 2) are overlaid on pages 4 and 5 of this report for ease of comparison purposes. You will note that the overlays presented in Figures 1 and 2 are different. Also attached to this report are the individual insize and undersize Mastersizer result analysis reports for your perusal.

| Sample | Significant Observations |
|--------|--|
| A1 | Sediment was not too coarse. Packed well in the vial as was |
| | difficult to remove for analysis. |
| B1 | As above |
| C1 | As above |
| D1 | As above |
| A4 | Sediment was coarser than those above and as such was |
| | removed from the supplied vial much easier. |
| B4 | Appeared of similar consistency to A—D1, maybe slightly coarser. |
| C4 | As for B4 |
| D4 | As for B4. Very large stones removed from this sample on the |
| | sieve. Stones up to 4cm in size (4 of them). |
| A5 | Large shell and stones again removed from this sample on the |
| | sieve. |
| B5 | - |
| C5 | - |
| D5 | - |
| B6 | The majority of this sample was >2mm in size and as such most |
| | of the sample remained on the sieve. |
| C6 | <2mm sediment was difficult to size due to the sample being |
| | coarse and 'dropping out' in the Mastersizer feed tuning. Only one |
| | measurement made. No averaging performed here. |
| D6 | - |
| I | I . |

Table 1: Supplied marine sediment samples and observations made during the sizing analysis.




Figure 1: Overlay of the measured PSD plots for ease of comparison purposes.




Figure 2: Overlay of the measured PSD plots for ease of comparison purposes.





Sample Name: A1 - Average

SOP Name:

Tuesday, 17 January 2012 10:53:59 AM

Sample Source & type:

Measured by:

Analysed:

abbottsj

Tuesday, 17 January 2012 10:54:01 AM

Sample bulk lot ref:

Result Source: Averaged

Particle Name:

Accessory Name: Hydro 2000G (A)

Sensitivity:

Silica 1.45 Particle RI:

Absorption:

Normal Obscuration:

1.450 **Dispersant Name:**

to 2000.000 um

Dispersant RI:

0.020 Weighted Residual: 13.88 % **Result Emulation:**

Water

1.330

5.323

Size range:

Analysis model:

General purpose

Concentration:

Span:

Uniformity:

Result units:

0.3346 %Vol 1.729


Volume

Specific Surface Area: 0.037

Surface Weighted Mean D[3,2]: 162.231 um

Vol. Weighted Mean D[4,3]: 647.587 um

d(0.1): 111.734 um d(0.5): 627.177 um d(0.9): 1196.139 um



Operator notes:





Sample Name:

SOP Name:

B1 - Average

Tuesday, 17 January 2012 11:13:59 AM

Sample Source & type:

Measured by:

Analysed:

abbottsj

Sample bulk lot ref:

Result Source:

Averaged

Tuesday, 17 January 2012 11:14:01 AM

Accessory Name: Hydro 2000G (A)

Sensitivity:

Silica 1.45 Particle RI:

Normal

1.450

Absorption:

Obscuration:

Dispersant Name:

Particle Name:

0.1

to 2000.000 um 15.33 %

d(0.9):

Dispersant RI:

Water

Weighted Residual:

Analysis model:

General purpose

Result Emulation:

1.330

1.416

Size range:

Off

Concentration:

Span:

Uniformity:

0.020

Result units:

0.0505 %Vol 4.300

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]:

Volume

Specific Surface Area: 0.37

16.237 um

225.161

d(0.1): 8.772


um

d(0.5):

132.553

um

578.695 um



Operator notes:





Sample Name:

SOP Name:

C1 - Average

Measured by:

Tuesday, 17 January 2012 11:35:01 AM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Averaged

Tuesday, 17 January 2012 11:35:03 AM

Particle Name:

Accessory Name: Silica 1.45 Hydro 2000G (A)

um

Analysis model: Sensitivity: General purpose Normal

Particle RI: 1.450

Absorption:

Size range: Obscuration:

Dispersant Name:

0.1

0.020 to 2000.000 um 19.99 %

Dispersant RI:

Weighted Residual:

Result Emulation:

Water

1.330

1.804

Off

d(0.9):

Concentration:

Span:

Uniformity:

Result units: Volume

0.0595 %Vol 2.871

Surface Weighted Mean D[3,2]:

228.844

Vol. Weighted Mean D[4,3]:

284.412

Specific Surface Area: 0.439

6.900


d(0.1):

13.679 um

d(0.5):

um

663.849 um



C1 - Average, Tuesday, 17 January 2012 11:35:01 AM

| Size (µm) | volume in % | Size (µm) | volume in % | |
|-----------|-------------|-----------|-------------|--|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | | 0.120 | | |
| 0.013 | 0.00 | 0.138 | 0.00 | |
| 0.015 | 0.00 | 0.158 | 0.00 | |
| 0.017 | 0.00 | 0.182 | 0.00 | |
| 0.020 | 0.00 | 0.209 | 0.00 | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| | 0.00 | | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.01 | |
| 0.030 | 0.00 | 0.316 | 0.08 | |
| 0.035 | 0.00 | 0.363 | 0.16 | |
| 0.040 | 0.00 | 0.417 | 0.23 | |
| 0.046 | 0.00 | 0.479 | 0.23 | |
| 0.052 | | 0.550 | | |
| 0.060 | 0.00 | 0.631 | 0.30 | |
| 0.069 | 0.00 | 0.724 | 0.30 | |
| 0.079 | 0.00 | 0.832 | 0.29 | |
| 0.079 | 0.00 | 0.955 | 0.25 | |
| | 0.00 | | 0.22 | |
| 0.105 | | 1.096 | | |

| Size (µm) | Volume In % |
|-----------|-------------|
| 1.096 | 0.19 |
| 1.259 | |
| 1.445 | 0.18 |
| 1.660 | 0.20 |
| 1.905 | 0.25 |
| 2.188 | 0.32 |
| 2.512 | 0.41 |
| - | 0.52 |
| 2.884 | 0.64 |
| 3.311 | 0.76 |
| 3.802 | 0.87 |
| 4.365 | 0.97 |
| 5.012 | 1.06 |
| 5.754 | 1.13 |
| 6.607 | _ |
| 7.586 | 1.19 |
| 8.710 | 1.22 |
| 10.000 | 1.24 |
| | 1.23 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | |
| 13.183 | 1.21 |
| 15.136 | 1.18 |
| 17.378 | 1.13 |
| 19.953 | 1.08 |
| 22.909 | 1.04 |
| 26.303 | 1.02 |
| 30.200 | 1.03 |
| 34.674 | 1.07 |
| 39.811 | 1.14 |
| 45.709 | 1.26 |
| 52.481 | 1.40 |
| 60.256 | 1.55 |
| 69.183 | 1.70 |
| 79.433 | 1.83 |
| | 1.93 |
| 91.201 | 2.01 |
| 104.713 | 2.08 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 120.226 | 2.16 |
| 138.038 | 2.10 |
| 158.489 | |
| 181.970 | 2.49 |
| 208.930 | 2.81 |
| 239.883 | 3.25 |
| 275.423 | 3.81 |
| 316.228 | 4.46 |
| 363.078 | 5.13 |
| 416.869 | 5.70 |
| 478,630 | 6.06 |
| 549.541 | 6.09 |
| 630.957 | 5.68 |

724.436

831.764

954.993

1096.478

1258.925

4 84

3.64

2.22

0.92

0.29

| Size (µm) | Volume In % |
|-----------|-----------------|
| 1258.925 | VOIGITIO III 70 |
| 1445.440 | 0.00 |
| | 0.00 |
| 1659.587 | 0.00 |
| 1905.461 | 0.00 |
| 2187.762 | 0.00 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311.311 | 0.00 |
| 3801.894 | |
| 4365.158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

D1 - Average

Tuesday, 17 January 2012 11:54:27 AM

Sample Source & type:

Measured by: abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Averaged

Tuesday, 17 January 2012 11:54:29 AM

Particle Name: Silica 1.45

Accessory Name:

Analysis model: General purpose

Sensitivity: Normal

Particle RI:

Hydro 2000G (A)

Size range:

Obscuration:

Absorption:

to 2000.000 um

17.49 %

1.450 **Dispersant Name:** 0.1

Weighted Residual:

Result Emulation:

Water

Dispersant RI: 1.330

1.059

Off

Span:

Uniformity:

0.020

Result units:

Concentration: 0.0440 %Vol

6.518

Volume In %

0.00 0.00 0.00 0.00 0.00 0.00 0.01 0.08 0.18 0.26 0.31 0.34 0.35 0.32

0.28

0.23

1.096

Volume

Specific Surface Area: 0.502

Surface Weighted Mean D[3,2]: 11.943 um

Vol. Weighted Mean D[4,3]: 202.146

d(0.1): 6.517


um

d(0.5):

89.308

um

d(0.9): 588.636 um



| Size (µm) | Volume In % | Size (µm) | |
|-----------|-------------|-----------|---|
| 0.010 | 0.00 | 0.105 | |
| 0.011 | 0.00 | 0.120 | |
| 0.013 | 0.00 | 0.138 | |
| 0.015 | 0.00 | 0.158 | |
| 0.017 | 0.00 | 0.182 | |
| 0.020 | | 0.209 | |
| 0.023 | 0.00 | 0.240 | |
| 0.026 | 0.00 | 0.275 | |
| 0.030 | 0.00 | 0.316 | |
| 0.035 | 0.00 | 0.363 | |
| 0.040 | 0.00 | 0.417 | |
| 0.046 | 0.00 | 0.479 | |
| 0.052 | 0.00 | 0.550 | |
| 0.060 | 0.00 | 0.631 | |
| 0.069 | 0.00 | 0.724 | |
| 0.079 | 0.00 | 0.832 | |
| 0.091 | 0.00 | 0.955 | |
| | 0.00 | | П |

| Size (µm) | Volume In % |
|-----------|-------------|
| 1.096 | 0.19 |
| 1.259 | 0.13 |
| 1.445 | 0.17 |
| 1.660 | 0.16 |
| 1.905 | |
| 2.188 | 0.28 |
| 2.512 | 0.38 |
| 2.884 | 0.49 |
| 3.311 | 0.61 |
| 3.802 | 0.75 |
| 4.365 | 0.90 |
| 5.012 | 1.05 |
| 5.754 | 1.20 |
| 6.607 | 1.35 |
| 7.586 | 1.49 |
| 8.710 | 1.61 |
| | 1.71 |
| 10.000 | 1.78 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 1.82 |
| 13.183 | |
| 15.136 | 1.83 |
| 17.378 | 1.83 |
| 19.953 | 1.82 |
| 22.909 | 1.81 |
| 26.303 | 1.84 |
| 30.200 | 1.90 |
| 34.674 | 2.01 |
| | 2.16 |
| 39.811 | 2.35 |
| 45.709 | 2.55 |
| 52.481 | 2.75 |
| 60.256 | 2.92 |
| 69.183 | 3.04 |
| 79.433 | |
| 91.201 | 3.12 |
| 104.713 | 3.15 |
| 120.226 | 3.15 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 120.226 | 3.13 |
| 138.038 | 3.10 |
| 158.489 | 3.10 |
| 181.970 | 3.03 |
| 208.930 | 2.98 |
| 239.883 | |
| 275.423 | 2.92 |
| 316.228 | 2.87 |
| 363.078 | 2.81 |
| 416.869 | 2.75 |
| 478.630 | 2.70 |
| 549.541 | 2.62 |
| 630.957 | 2.51 |
| 724.436 | 2.32 |
| 831.764 | 2.06 |
| | 1 71 |

954.993

1096.478

1258.925

1.29

0.88

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 0.48 |
| 1445.440 | |
| 1659.587 | 0.02 |
| 1905.461 | 0.00 |
| 2187.762 | 0.00 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311 311 | 0.00 |
| 3801.894 | 0.00 |
| 4365.158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

Measured:

A4 - Average

Tuesday, 17 January 2012 12:18:14 PM

Sample Source & type:

Measured by: abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Averaged

Tuesday, 17 January 2012 12:18:16 PM

Particle Name:

Accessory Name: Hydro 2000G (A)

Analysis model: Sensitivity: General purpose Normal

Silica 1.45

Particle RI:

Absorption:

Size range:

Obscuration:

1.450

0.1

0.020 to 2000.000 um

12.04 %

Dispersant Name:

d(0.1):

Dispersant RI:

Weighted Residual:

Result Emulation:

Water

1.330

4.757 %

Off

Concentration:

Span:

Uniformity:

Result units:

0.4640 %Vol

1.623

0..00

Volume

Specific Surface Area: 0.0239 m²/g

Surface Weighted Mean D[3,2]: 250.903 um

Vol. Weighted Mean D[4,3]: 636.603 um

185.391


um

d(0.5):

593.174

um

d(0.9): 1148.302 um



| Size (µm) | Volume In % | Size (µ | m) Volume In % | Size (µm) | Volume In % |
|-----------|-------------|-----------|-------------|-----------|-------------|-----------|--------------|---------|----------------|-----------|-------------|
| 0.010 | 0.00 | 0.105 | 0.00 | 1.096 | 0.00 | 11.482 | 0.14 | 120.2 | 26 | 1258.925 | 3.86 |
| 0.011 | 0.00 | 0.120 | 0.00 | 1.259 | 0.00 | 13.183 | 0.15 | 138.0 | 38 1.48 | 1445.440 | 2.30 |
| 0.013 | 0.00 | 0.138 | 0.00 | 1.445 | 0.00 | 15.136 | 0.15 | 158.4 | 1.73 | 1659.587 | 0.33 |
| 0.015 | 0.00 | 0.158 | 0.00 | 1.660 | 0.00 | 17.378 | 0.16 | 181.9 | 70 2.02 | 1905.461 | 0.01 |
| 0.017 | 0.00 | 0.182 | 0.00 | 1.905 | 0.00 | 19.953 | 0.16 | 208.9 | 2.40 | 2187.762 | 0.00 |
| 0.020 | 0.00 | 0.209 | 0.00 | 2.188 | 0.00 | 22.909 | 0.15 | 239.8 | 33 2.91 | 2511.886 | 0.00 |
| 0.023 | 0.00 | 0.240 | 0.00 | 2.512 | 0.00 | 26.303 | 0.13 | 275.4 | 3.59 | 2884.032 | 0.00 |
| 0.026 | 0.00 | 0.275 | 0.00 | 2.884 | 0.00 | 30.200 | 0.14 | 316.2 | 28 4.48 | 3311.311 | 0.00 |
| 0.030 | 0.00 | 0.316 | 0.00 | 3.311 | 0.00 | 34.674 | 0.12 | 363.0 | 78 5.55 | 3801.894 | 0.00 |
| 0.035 | 0.00 | 0.363 | 0.00 | 3.802 | 0.00 | 39.811 | 0.10 | 416.8 | 6.72 | 4365.158 | 0.00 |
| 0.040 | 0.00 | 0.417 | 0.00 | 4.365 | 0.02 | 45.709 | 0.10 | 478.6 | 7.84 | 5011.872 | 0.00 |
| 0.046 | 0.00 | 0.479 | 0.00 | 5.012 | 0.06 | 52.481 | 0.12 | 549.5 | 8.74 | 5754.399 | 0.00 |
| 0.052 | 0.00 | 0.550 | 0.00 | 5.754 | 0.09 | 60.256 | 0.17 | 630.9 | 9.22 | 6606.934 | 0.00 |
| 0.060 | 0.00 | 0.631 | 0.00 | 6.607 | 0.09 | 69.183 | 0.25 | 724.4 | 9.14 | 7585.776 | 0.00 |
| 0.069 | 0.00 | 0.724 | 0.00 | 7.586 | 0.10 | 79.433 | | 831.7 | 8.43 | 8709.636 | 0.00 |
| 0.079 | 0.00 | 0.832 | 0.00 | 8.710 | 0.11 | 91.201 | 0.56 0.76 | 954.9 | 7.17 | 10000.000 | 0.00 |
| 0.091 | 0.00 | 0.955 | 0.00 | 10.000 | 0.12 | 104.713 | 1.00 | 1096.4 | 78 5.55 | | |
| 0.105 | 0.00 | 1.096 | 0.00 | 11.482 | 0.13 | 120.226 | 1.00 | 1258.9 | 25 | | |

Operator notes:





Sample Name:

SOP Name:

B4 - Average

Measured by:

Tuesday, 17 January 2012 12:31:36 PM

abbottsj

Analysed:

Sample bulk lot ref:

Sample Source & type:

Result Source:

Averaged

Tuesday, 17 January 2012 12:31:37 PM

Particle Name:

Accessory Name: Hydro 2000G (A)

Sensitivity: Normal

Silica 1.45 Particle RI: 1.450

Absorption:

Dispersant Name:

Size range: 0.020

Obscuration:

Dispersant RI:

to 2000.000 um 19.63 % **Result Emulation:**

Water 1.330

um

2.114

Analysis model:

General purpose

Weighted Residual:

Off

Concentration:

Span:

0.1

Uniformity:

Result units:

0.0778 %Vol 2.844

Volume

Specific Surface Area: 0.328

Surface Weighted Mean D[3,2]: 18.298

Vol. Weighted Mean D[4,3]: 370.929

10.615


d(0.1):

d(0.5): 297.105 um

d(0.9):

855.563

um



| 0.010 0.011 0.013 0.015 0.017 0.020 0.023 0.026 0.030 0.035 | Volume In % 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0 | 0.105 0.120 0.138 0.158 0.182 0.209 0.240 0.275 0.316 0.363 0.417 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | 1.096 1.259 1.445 1.660 1.905 2.188 2.512 2.884 3.311 3.802 4.365 | 0.13 0.12 0.13 0.16 0.21 0.28 | 11.482 13.183 15.136 17.378 19.953 22.909 26.303 30.200 34.674 39.811 45.709 | 0.95 0.93 0.89 0.85 0.82 0.79 0.78 0.81 | 120.22 138.03 158.48 181.97 208.93 239.88 275.42 316.22 363.07 416.86 | 2.30 2.46 2.66 2.91 3.22 3.3 3.61 4.07 8.4.57 6.9 5.48 5.75 | 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3801.894 4365.158 5011.872 | 1.24 0.09 0.00 0.00 0.00 0.00 0.00 0.00 0.0 |
|--|---|---|--|---|--|--|--|--|--|--|---|
| 0.020 0.023 0.026 0.030 0.035 | 0.00 0.00 0.00 0.00 0.00 | 0.209 0.240 0.275 0.316 0.363 | 0.00 0.00 0.00 0.07 0.13 0.17 0.21 0.23 0.23 0.21 0.18 | 2.188 2.512 2.884 3.311 3.802 | 0.28 0.36 0.44 0.53 0.61 0.69 0.77 0.83 0.88 0.92 0.94 | 22.909 26.303 30.200 34.674 39.811 | 0.82 0.79 0.78 0.81 0.87 0.97 1.11 1.28 1.46 1.65 1.83 1.99 | 239.88 275.42 316.22 363.07 416.86 | 3.22 3.61 3.61 4.07 5.06 5.48 5.75 5.79 7 5.54 4.99 4.19 3.321 3.21 | 2511.886 2884.032 3311.311 3801.894 4365.158 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 |

Operator notes:





Sample Name:

SOP Name:

C4 - Average

Measured by:

Tuesday, 17 January 2012 1:40:57 PM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 1:40:58 PM

Averaged

Particle Name: Silica 1.45

Accessory Name:

Sensitivity:

Particle RI:

Hydro 2000G (A)

Normal

1.450

Absorption:

Obscuration:

Dispersant Name:

Concentration:

0.1

um

Size range: to 2000.000 um

22.36 %

Dispersant RI:

Weighted Residual:

Analysis model:

General purpose

Result Emulation:

Water

2.813

Off

1.330

Uniformity:

0.020

0.0926

0.327

d(0.1):

Span:

Result units: Volume

Specific Surface Area:

%Vol

11.695

um

2.725

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]:


457.101

18.372

d(0.5): 382.678 um

d(0.9):

1054.336 um



| Size (µm) | Volume In % | Size (µm) | Volume In % | |
|-----------|-------------|-----------|-------------|--|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | |
| 0.013 | 0.00 | 0.138 | 0.00 | |
| 0.015 | | 0.158 | 0.00 | |
| 0.017 | 0.00 | 0.182 | | |
| 0.020 | 0.00 | 0.209 | 0.00 | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.00 | |
| 0.030 | 0.00 | 0.316 | 0.01 | |
| 0.035 | 0.00 | 0.363 | 0.07 | |
| 0.040 | 0.00 | 0.417 | 0.13 | |
| 0.046 | 0.00 | 0.479 | 0.18 | |
| 0.052 | 0.00 | 0.550 | 0.21 | |
| 0.060 | 0.00 | 0.631 | 0.23 | |
| 0.069 | 0.00 | 0.724 | 0.23 | |
| 0.009 | 0.00 | 0.724 | 0.22 | |
| 0.079 | 0.00 | 0.052 | 0.19 | |
| 0.091 | 0.00 | 1.006 | 0.16 | |

| Size (µm) | Volume In % | |
|-----------|-------------|--|
| 1.096 | 0.14 | |
| 1.259 | 0.14 | |
| 1.445 | 0.12 | |
| 1.660 | 0.13 | |
| 1.905 | | |
| 2.188 | 0.20 | |
| 2.512 | 0.26 | |
| 2.884 | 0.33 | |
| 3.311 | 0.40 | |
| 3.802 | 0.48 | |
| 4.365 | 0.55 | |
| 5.012 | 0.62 | |
| 5.754 | 0.69 | |
| 6.607 | 0.75 | |
| 7.586 | 0.80 | |
| 8.710 | 0.84 | |
| 10.000 | 0.87 | |
| 11 400 | 0.89 | |

| Size (µm) | Volume In % | |
|-----------|-------------|---|
| 11.482 | 0.90 | |
| 13.183 | 0.89 | |
| 15.136 | | |
| 17.378 | 0.88 | |
| 19.953 | 0.86 | |
| 22.909 | 0.85 | |
| 26.303 | 0.85 | |
| 30.200 | 0.89 | |
| 34.674 | 0.95 | |
| 39.811 | 1.06 | |
| 45.709 | 1.21 | |
| 52.481 | 1.37 | |
| | 1.55 | |
| 60.256 | 1.72 | |
| 69.183 | 1.85 | |
| 79.433 | 1.95 | |
| 91.201 | 1.99 | |
| 104.713 | 1.99 | |
| 120 226 | 1.00 | 1 |

| ĺ | Size (µm) | Volume In % |
|---|-----------|--------------|
| | 120.226 | 1.96 |
| | 138.038 | 1.90 |
| | 158.489 | 1.91 |
| | 181.970 | 1.86 |
| | 208.930 | 1.00 |
| | 239.883 | |
| | 275.423 | 2.12 2.45 |
| | 316.228 | |
| | 363.078 | 2.96 3.63 |
| | 416.869 | 5.55 |
| | 478.630 | 4.41 |
| | 549.541 | 5.20 |
| | 630.957 | 5.87 |
| | 724.436 | 6.28 |
| | 831.764 | 6.30 |
| | | 5 88 |

1096.478

1258.925

5.06

3.96

| _ | | |
|---|---|----------------------|
| | Size (µm) | Volume In % |
| | 1258.925 | 2.77 |
| | 1445.440 | 168 |
| | 1659.587 | |
| | 1905.461 | 0.25 |
| | 2187.762 | 0.01 |
| | 2511.886 | 0.00 |
| | 2884.032 | 0.00 |
| | 3311.311 | 0.00 |
| | 3801.894 | 0.00 |
| | 4365.158 | 0.00 |
| | | 0.00 |
| | | 0.00 |
| | | 0.00 |
| | | 0.00 |
| | | 0.00 |
| | 8709.636 | 0.00 |
| | 10000.000 | 0.00 |
| | | |
| | | |
| | 5011.872 5754.399 6606.934 7585.776 8709.636 10000.000 | 0.00 0.00 0.00 |

Operator notes:

Malvern Instruments Ltd. Tel := +[44] (0) 1684-892456 Fax +[44] (0) 1684-892789

Mastersizer 2000 Ver. 5.60 Serial Number: MAL1006765 File name: SKM 170112.mea Record Number: 32 18/01/2012 10:39:20 4





Sample Name:

SOP Name:

D4 - Average Sample Source & type: Tuesday, 17 January 2012 2:10:11 PM

Measured by: abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Averaged

Tuesday, 17 January 2012 2:10:13 PM

Particle Name:

Accessory Name: Hydro 2000G (A)

Sensitivity:

Silica 1.45 Particle RI:

Normal

1.450

Absorption:

Dispersant Name:

0.1

Size range: to 2000.000 um Obscuration: 16.69 % **Result Emulation:**

Water

Dispersant RI:

Weighted Residual: 2.435

Analysis model:

General purpose

Off

Concentration:

Span:

1.330

Uniformity:

0.020

Result units:

0.0526 %Vol 3.413

Volume

Specific Surface Area: 0.487

Surface Weighted Mean D[3,2]: 12.323 um

Vol. Weighted Mean D[4,3]:

367.026

d(0.1):

6.026


um

d(0.5):

267.250

um

d(0.9): 918.145 um



| Size (µm) 0.010 0.011 0.013 0.015 0.017 0.020 0.023 0.026 0.030 0.035 0.040 0.046 0.052 0.060 0.069 | Volume In % 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0 | Size (µm) 0.105 0.120 0.138 0.158 0.182 0.209 0.240 0.275 0.316 0.363 0.417 0.479 0.550 0.631 0.724 | Volume In % | Size (µm) 1.096 1.259 1.445 1.660 1.905 2.188 2.512 2.884 3.311 3.802 4.365 5.012 5.754 6.607 7.586 | 0.23 0.22 0.23 0.28 0.36 0.46 0.58 0.71 0.84 0.98 1.11 1.23 | Size (µm) 11.482 13.183 15.136 17.378 19.953 22.909 26.303 30.200 34.674 39.811 45.709 52.481 60.256 69.183 79.433 | 1.44 1.40 1.35 1.31 1.29 1.32 1.38 1.47 1.56 1.65 1.71 1.73 | Size (µm) 120.226 138.038 158.489 181.970 208.930 239.883 275.423 316.228 363.078 416.869 478.630 549.541 630.957 724.436 831.764 | Volume In % 1.24 1.19 1.23 1.38 1.66 2.12 2.72 3.42 4.14 4.80 5.27 5.50 5.42 5.03 | Size (µm) 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3801.894 4365.158 5011.872 5754.399 6606.934 67585.776 8709.636 | Volume In % 1.75 0.99 0.07 0.00 0.00 0.00 0.00 0.00 0.00 | t |
|--|---|---|-------------|---|--|--|--|---|--|---|--|---|
| 0.052 0.060 | 0.00 | 0.550 0.631 | 0.34 | 5.754 6.607 | 1.32 | 60.256 69.183 | 1.73 | 630.957 724.436 | 5.42 | 6606.934 7585.776 | 0.00 | |

Operator notes:

Mastersizer 2000 Ver. 5.60 Serial Number: MAL1006765





Sample Name:

SOP Name:

A5 - Average

Tuesday, 17 January 2012 2:25:54 PM

Sample Source & type:

Measured by: abbottsj

Analysed:

Sample bulk lot ref:

Tuesday, 17 January 2012 2:25:56 PM

Result Source:

Averaged

Particle Name: Silica 1.45

Accessory Name: Hydro 2000G (A)

Analysis model: General purpose

Sensitivity: Normal

Particle RI:

Absorption:

Size range:

1.450

to 2000.000 um

Obscuration:

Dispersant Name:

0.1

0.020

13.48 %

1.0423

Dispersant RI:

Weighted Residual:

Result Emulation:

Water

1.330

5.186

Off

Concentration:

Span:

Uniformity:

Result units: Volume

Specific Surface Area:

%Vol

1.103


Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]: 593.948

0.0118

506.982 um

um



| Size (µm) | Volume In % | |
|-----------|-------------|--|
| 0.010 | 0.00 | |
| 0.011 | 0.00 | |
| 0.013 | 0.00 | |
| 0.015 | | |
| 0.017 | 0.00 | |
| 0.020 | 0.00 | |
| 0.023 | 0.00 | |
| 0.026 | 0.00 | |
| 0.030 | 0.00 | |
| 0.035 | 0.00 | |
| 0.040 | 0.00 | |
| 0.046 | 0.00 | |
| 0.052 | 0.00 | |
| 0.060 | 0.00 | |
| 0.069 | 0.00 | |
| | 0.00 | |
| 0.079 | 0.00 | |
| 0.091 | 0.00 | |
| | | |

| iay, i | / Januar | <u>y</u> 2 |
|----------|-------------|------------|
| ize (µm) | Volume In % | |
| 0.105 | 0.00 | |
| 0.120 | 0.00 | |
| 0.138 | | |
| 0.158 | 0.00 | |
| 0.182 | 0.00 | |
| 0.209 | 0.00 | |
| 0.240 | 0.00 | |
| 0.275 | 0.00 | |
| 0.316 | 0.00 | |
| 0.363 | 0.00 | |
| 0.303 | 0.00 | |
| - | 0.00 | |
| 0.479 | 0.00 | |
| 0.550 | 0.00 | |
| 0.631 | 0.00 | |
| 0.724 | 0.00 | |
| 0.832 | 0.00 | |
| 0.955 | 0.00 | |
| 4 000 | 0.00 | |

| Size (µm) | Volume In % | |
|-----------|-------------|--|
| 1.096 | 0.00 | |
| 1.259 | | |
| 1.445 | 0.00 | |
| 1.660 | 0.00 | |
| 1.905 | 0.00 | |
| 2.188 | 0.00 | |
| 2.512 | 0.00 | |
| 2.884 | 0.00 | |
| 3.311 | 0.00 | |
| | 0.00 | |
| 3.802 | 0.00 | |
| 4.365 | 0.00 | |
| 5.012 | 0.00 | |
| 5.754 | 0.00 | |
| 6.607 | 0.00 | |
| 7.586 | | |
| 8.710 | 0.00 | |
| 10.000 | 0.00 | |
| 11.482 | 0.00 | |
| .1102 | | |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 0.00 |
| 13.183 | |
| 15.136 | 0.00 |
| 17.378 | 0.00 |
| 19.953 | 0.00 |
| 22,909 | 0.00 |
| 26.303 | 0.00 |
| 30.200 | 0.00 |
| 34.674 | 0.00 |
| 39.811 | 0.00 |
| | 0.00 |
| 45.709 | 0.00 |
| 52.481 | 0.00 |
| 60.256 | 0.00 |
| 69.183 | 0.00 |
| 79.433 | 0.00 |
| 91.201 | 0.00 |
| 104.713 | 5.55 |
| 400.000 | 0.00 |

| 1 | O: () | 1 |
|---|-----------|-------------|
| | Size (µm) | Volume In % |
| | 120.226 | 0.00 |
| | 138.038 | 0.00 |
| | 158.489 | 5.55 |
| | 181.970 | 0.02 |
| | 208,930 | 0.23 |
| | 239.883 | 1.32 |
| | 275.423 | 2.89 |
| | 316.228 | 4.89 |
| | 0.00 | 7.17 |
| | 363.078 | 9.36 |
| | 416.869 | 11.17 |
| | 478.630 | 12.26 |
| | 549.541 | 12.43 |
| | 630.957 | 11.65 |
| | 724.436 | |
| | 831.764 | 10.02 |
| | 954,993 | 7.81 |
| | 4000 470 | 5.39 |

1096.478

1258.925

2.98

Operator notes:

Malvern Instruments Ltd. Tel := +[44] (0) 1684-892456 Fax +[44] (0) 1684-892789

Mastersizer 2000 Ver. 5.60 Serial Number: MAL1006765 File name: SKM 170112.mea Record Number: 40 18/01/2012 10:39:22





Sample Name:

SOP Name:

B5 - Average

Tuesday, 17 January 2012 2:56:04 PM

Sample Source & type:

Measured by: Analysed: abbottsj

Sample bulk lot ref:

Result Source:

Averaged

Tuesday, 17 January 2012 2:56:06 PM

Particle Name:

Accessory Name: Hydro 2000G (A)

Sensitivity:

Silica 1.45 Particle RI:

Normal

1.450

Absorption: 0.1

Obscuration:

Dispersant Name:

Dispersant RI:

to 2000.000 um 15.04 %

Water

Weighted Residual:

Result Emulation:

1.330

3.362

0.020

Off

Concentration: 0.1782 %Vol Span:

Uniformity:

Result units:

Specific Surface Area:

2.039

Vol. Weighted Mean D[4,3]:

Volume

0.0748

Surface Weighted Mean D[3,2]: 80.228 um

500.129

Analysis model:

General purpose

Size range:

d(0.1): 59.017 um


d(0.5):

458.886

um

d(0.9): 994.662

um



| 0.010 0.011 0.013 0.015 0.017 0.020 0.023 0.026 0.030 0.035 0.040 0.052 0.060 0.069 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | Size (µm) 0.105 0.120 0.138 0.158 0.182 0.209 0.240 0.275 0.316 0.363 0.417 0.479 0.550 0.631 0.724 0.832 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | Size (µm) 1.096 1.259 1.445 1.660 1.905 2.188 2.512 2.884 3.311 3.802 4.365 5.012 5.754 6.607 7.586 8.710 | Volume In % 0.00 0.00 0.00 0.00 0.07 0.10 0.13 0.18 0.23 0.27 0.32 0.36 0.40 0.43 0.45 0.45 | 11.482 13.183 15.136 17.378 19.953 22.909 26.303 30.200 34.674 39.811 45.709 52.481 60.256 69.183 79.433 91.201 | Volume In % 0.49 0.48 0.46 0.45 0.44 0.45 0.57 0.66 0.78 0.91 1.06 1.21 1.37 | 120.226 138.038 158.489 181.970 208.930 239.883 275.423 316.228 363.078 416.869 478.630 549.541 630.957 724.436 831.764 954.993 | Volume In % 1.72 1.92 2.16 2.47 2.88 3.41 4.08 4.88 5.75 6.60 7.30 7.72 7.72 7.24 6.30 5.03 | Size (µm) 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3801.894 4365.158 5011.872 5754.399 6606.934 7585.776 8709.636 10000.000 | Volume In % 2.22 0.74 0.03 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | |
|--|--|---|--|---|--|--|---|--|---|--|--|--|
| 0.091 0.105 | 0.00 | 0.955 1.096 | 0.00 | 10.000 11.482 | 0.48 | 104.713 120.226 | 1.54 | 1096.478 1258.925 | 3.61 | | | |

Operator notes:





Sample Name:

SOP Name:

C5 - Average

Measured by:

Tuesday, 17 January 2012 3:07:35 PM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 3:07:36 PM

Averaged

Particle Name:

Accessory Name: Hydro 2000G (A) Silica 1.45

Particle RI:

Absorption: 0.1

Dispersant Name:

Dispersant RI:

Water

0.0508

1.450

1.330

Analysis model: General purpose

Size range:

to 2000.000 um

Weighted Residual:

3.315

Sensitivity: Normal

Obscuration:

14.77 %

Result Emulation:

Off

Concentration: 0.2673

%Vol

Specific Surface Area:

Span: 1.859

Surface Weighted Mean D[3,2]:

118.227 um **Uniformity:**

0.020

Result units: Volume

Vol. Weighted Mean D[4,3]:

493.829

d(0.1):

109.265

um

d(0.5):


448.521

um

d(0.9):

943.215

um



| Size (µm) | Volume In % | Size (µm) | Volume In % | Size (µm) | Volume In % | |
|-----------|-------------|-----------|-------------|-----------|-------------|--|
| 0.010 | 0.00 | 0.105 | 0.00 | 1.096 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | 1.259 | 0.00 | |
| 0.013 | 0.00 | 0.138 | 0.00 | 1.445 | 0.00 | |
| 0.015 | 0.00 | 0.158 | 0.00 | 1.660 | 0.00 | |
| 0.017 | 0.00 | 0.182 | 0.00 | 1.905 | 0.00 | |
| 0.020 | 0.00 | 0.209 | 0.00 | 2.188 | 0.02 | |
| 0.023 | 0.00 | 0.240 | 0.00 | 2.512 | 0.02 | |
| 0.026 | 0.00 | 0.275 | 0.00 | 2.884 | 0.10 | |
| 0.030 | 0.00 | 0.316 | 0.00 | 3.311 | 0.10 | |
| 0.035 | 0.00 | 0.363 | 0.00 | 3.802 | 0.16 | |
| 0.040 | 0.00 | 0.417 | 0.00 | 4.365 | 0.10 | |
| 0.046 | 0.00 | 0.479 | 0.00 | 5.012 | 0.13 | |
| 0.052 | 0.00 | 0.550 | 0.00 | 5.754 | 0.24 | |
| 0.060 | 0.00 | 0.631 | 0.00 | 6.607 | 0.27 | |
| 0.069 | 0.00 | 0.724 | 0.00 | 7.586 | 0.29 | |
| 0.079 | 0.00 | 0.832 | 0.00 | 8.710 | 0.30 | |
| 0.091 | 0.00 | 0.955 | 0.00 | 10.000 | 0.32 | |
| 0.105 | 0.00 | 1 096 | 0.00 | 11 482 | 0.02 | |

| Size (µm) | Volume In % | Size (µm) | Volume In % | Size (µm) | Volume In % |
|------------------|-------------|--------------------|-------------|----------------------|-------------|
| 11.482 | 0.33 | 120.226 | 1.56 | 1258.925 | 1.74 |
| 13.183 | 0.33 | 138.038 | 1.94 | 1445.440 | 0.15 |
| 15.136 | 0.33 | 158.489 | 2.41 | 1659.587 | 0.00 |
| 17.378 | 0.33 | 181.970 | 2.98 | 1905.461 | 0.00 |
| 19.953 | 0.32 | 208.930 | 3.65 | 2187.762 | 0.00 |
| 22.909 | 0.32 | 239.883 | 4.41 | 2511.886 | 0.00 |
| 26.303 | 0.32 | 275.423 | 5.23 | 2884.032 | 0.00 |
| 30.200 | 0.33 | 316.228 | 6.09 | 3311.311 | 0.00 |
| 34.674 | 0.34 | 363.078 | 6.89 | 3801.894 | 0.00 |
| 39.811 | 0.37 | 416.869 | 7.57 | 4365.158 | 0.00 |
| 45.709 | 0.42 | 478.630 | 8.01 | 5011.872 | 0.00 |
| 52.481 60.256 | 0.48 | 549.541 630.957 | 8.10 | 5754.399 6606.934 | 0.00 |
| 69.183 | 0.57 | 724.436 | 7.79 | 7585.776 | 0.00 |
| 79.433 | 0.68 | 831.764 | 7.04 | 8709.636 | 0.00 |
| 91,201 | 0.83 | 954.993 | 5.91 | 10000.000 | 0.00 |
| 104.713 | 1.02 | 1096.478 | 4.53 | 10000.000 | |
| 120.226 | 1.26 | 1258.925 | 3.10 | | |
| | | 50.020 | | | |

Operator notes:

Malvern Instruments Ltd. Tel := +[44] (0) 1684-892456 Fax +[44] (0) 1684-892789

Mastersizer 2000 Ver. 5.60 Serial Number: MAL1006765 File name: SKM 170112.mea Record Number: 48 18/01/2012 10:39:24





Sample Name:

SOP Name:

D5 - Average

Analysed:

Sample Source & type:

Measured by: abbottsj

Tuesday, 17 January 2012 3:19:03 PM

Tuesday, 17 January 2012 3:19:01 PM

Sample bulk lot ref:

Result Source:

Averaged

Particle Name: Silica 1.45

Accessory Name: Hydro 2000G (A)

General purpose

Sensitivity: Normal

Particle RI:

Absorption:

Size range:

Analysis model:

Obscuration:

1.450

0.1

0.020 to 2000.000 um 12.49 %

Dispersant Name: Water

Dispersant RI:

Weighted Residual:

Result Emulation:

1.330

3.547

Off

Concentration:

Span:

Uniformity:

Result units:

0.1342 %Vol 1.944

Volume

Specific Surface Area:

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]: 511.165

0.0813

73.819 um

d(0.9): 990.660 um

d(0.1): 64.419

um

d(0.5): 476.553 um

Particle Size Distribution 10 9 8 7 6 5 4 3 2 1 0.01 0.1 100 1000 3000 Particle Size (µm)

Size (µm) Volume In % 0.00 0.011 0.00 0.013 0.00 0.015 0.00 0.017 0.00 0.020 0.00 0.023 0.00 0.026 0.00 0.030 0.00 0.035 0.00 0.040 0.00 0.046 0.00 0.052 0.00 0.060 0.00 0.069 0.00 0.079 0.00 0.091 0.00 0.105

D5 - Average, Tuesday, 17 January 2012 3:19:01 PM Size (µm) Volume In % 0.120 0.00 0.138 0.00 0.158 0.00 0.182 0.00 0.209 0.00 0.240 0.00 0.275 0.00 0.316 0.00 0.363 0.00 0.417 0.00 0.479 0.00 0.550 0.00 0.631 0.00 0.724 0.00 0.832 0.00 0.955

0.00

1.096

Size (µm) Volume In % 0.00 1.259 0.00 1.445 0.01 1.660 0.06 1.905 0.09 2.188 0.13 2.512 0.18 2.884 0.23 3.311 0.28 3.802 0.33 4.365 0.38 5.012 5.754 0 44 6.607 0.46 7.586 0.47 8.710 0.47 10.000 0.46 11.482

Size (µm) Volume In % 13,183 0.42 15.136 0.40 17.378 0.37 19.953 0.36 22.909 0.35 26,303 0.36 30.200 0.38 34.674 0.43 39.811 0.49 45.709 0.58 52.481 0.67 60.256 0.78 69.183 0.90 79,433 1.02 91.201 1.13 104.713 1.26 120.226

| Size (µm) | Volume In % |
|-----------|-------------|
| 120.226 | 1.41 |
| 138.038 | 1.60 |
| 158.489 | 1.86 |
| 181.970 | 2.22 |
| 208.930 | |
| 239.883 | 2.71 |
| 275.423 | 3.36 |
| 316.228 | 4.15 |
| 363.078 | 5.09 |
| 416.869 | 6.08 |
| 478,630 | 7.02 |
| 549.541 | 7.77 |
| 630.957 | 8.19 |
| 724.436 | 8.15 |
| 724.400 | 7.50 |

831.764

954.993

1096.478

1258.925

6.55

5.16

3.65

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 2.16 |
| 1445.440 | 0.54 |
| 1659.587 | 0.00 |
| 1905.461 | |
| 2187.762 | 0.00 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311.311 | 0.00 |
| 3801.894 | 0.00 |
| 4365,158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | 0.00 |
| 10000.000 | |
| | |

Operator notes:





Sample Name:

SOP Name:

B6 - Average

Tuesday, 17 January 2012 3:31:12 PM

Sample Source & type:

Measured by:

Analysed:

abbottsj

Tuesday, 17 January 2012 3:31:14 PM

Sample bulk lot ref:

Particle Name:

Result Source:

Averaged

Analysis model:

Sensitivity:

Silica 1.45

Accessory Name: Hydro 2000G (A)

General purpose

Normal

Particle RI:

Absorption:

Size range:

Obscuration:

1.450

0.1

0.020 to 2000.000 um 11.59 %

Dispersant Name:

Dispersant RI:

Weighted Residual:

Result Emulation:

Water

1.330

6.876

Off

d(0.9):

d(0.1):

Span:

Uniformity:

Result units: Volume

Concentration: 0.2626 %Vol

1.699

Surface Weighted Mean D[3,2]:

705.996

Vol. Weighted Mean D[4,3]:

Specific Surface Area: 0.139

87.251

um

43.246

um


d(0.5):

718.950

um

1286.731

um



| Size (µm) Volume In % | | Size (µm) | Volume In % | Size (µm) | Volume In % | (1 / | Volume In % | | Volume In % |
|-----------------------|--------|-----------|-------------|-----------|-------------|----------|-------------|-----------|-------------|
| 0.010 0.00 | 0.00 | 1.096 | 0.06 | 11.482 | 0.42 | 120.226 | 0.81 | 1258.925 | 5.64 |
| 0.011 0.00 | 0.00 | 1.259 | 0.05 | 13.183 | 0.43 | 138.038 | 0.82 | 1445.440 | 3.66 |
| 0.013 0.1 | 8 | 1.445 | | 15.136 | | 158.489 | | 1659.587 | |
| 0.015 | 0.00 | 1.660 | 0.05 | 17.378 | 0.44 | 181.970 | 0.81 | 1905.461 | 1.51 |
| 0.00 | 0.00 | 1.905 | 0.05 | 19.953 | 0.44 | 208.930 | 0.78 | 2187.762 | 0.22 |
| 0.020 0.00 | 0.00 | 2.188 | 0.06 | 22.909 | 0.44 | 239.883 | 0.81 | 2511.886 | 0.00 |
| 0.023 | 0.00 | 2.512 | 0.08 | 26.303 | 0.42 | 275.423 | 0.97 | 2884.032 | 0.00 |
| 0.026 0.00 0.2 | 0.00 | 2.884 | 0.10 | 30.200 | 0.41 | 316.228 | 1.37 | 3311.311 | 0.00 |
| 0.00 | 0.00 | | 0.12 | | 0.39 | | 2.11 | | 0.00 |
| 0.030 0.00 | 0.00 | 3.311 | 0.14 | 34.674 | 0.38 | 363.078 | 3.22 | 3801.894 | 0.00 |
| 0.035 | 0.07 | 3.802 | 0.17 | 39.811 | 0.37 | 416.869 | 4.69 | 4365.158 | 0.00 |
| 0.040 0.00 0.4 | 7 0.09 | 4.365 | 0.20 | 45.709 | 0.37 | 478.630 | 6.36 | 5011.872 | 0.00 |
| 0.046 0.00 0.4 | 9 0.10 | 5.012 | 0.25 | 52.481 | 0.39 | 549.541 | 8.04 | 5754.399 | 0.00 |
| 0.052 0.00 | 0.10 | 5.754 | 0.28 | 60.256 | 0.43 | 630.957 | 9.41 | 6606.934 | 0.00 |
| 0.060 0.6 | 1 | 6.607 | | 69.183 | | 724.436 | | 7585.776 | |
| 0.069 | 0.11 | 7.586 | 0.31 | 79.433 | 0.50 | 831.764 | 10.17 | 8709.636 | 0.00 |
| 0.079 0.00 0.8 | 0.11 | 8.710 | 0.34 | 91.201 | 0.58 | 954,993 | 10.13 | 10000.000 | 0.00 |
| 0.00 | 5 0.09 | 10.000 | 0.37 | 104.713 | 0.67 | 1096,478 | 9.22 | | |
| 0.105 | 0.08 | 11.482 | 0.39 | 120,226 | 0.76 | 1258.925 | 7.62 | | |

Operator notes:





Sample Name:

SOP Name:

Tuesday, 17 January 2012 4:55:57 PM

Sample Source & type:

Measured by:

Analysed:

abbottsj

Tuesday, 17 January 2012 4:55:59 PM

Sample bulk lot ref:

Result Source: Measurement

Particle Name:

Accessory Name: Hydro 2000G (A)

Sensitivity:

Silica 1.45 Particle RI:

Normal

1.450

Absorption:

Obscuration:

Dispersant Name:

0.1

to 2000.000 um 5.58

Dispersant RI:

Weighted Residual:

%

Water

1.330

6.472

Size range:

Analysis model:

General purpose

Result Emulation:

Concentration:

Span:

Uniformity:

0.020

Result units:

0.1396 %Vol 1.563

Volume

Specific Surface Area: 0.0337


Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]: 770.974

177.992 um

695.114 um

1393.505 um



C6, Tuesday, 17 January 2012 4:55:57 PM

| Size (µm) | Volume In % | Size (µr |
|-----------|-------------|----------|
| 0.010 | | 0.10 |
| 0.011 | 0.00 | 0.12 |
| 0.013 | 0.00 | 0.13 |
| 0.015 | 0.00 | 0.15 |
| 0.017 | 0.00 | 0.18 |
| 0.020 | 0.00 | 0.20 |
| 0.023 | 0.00 | 0.24 |
| 0.026 | 0.00 | 0.27 |
| 0.030 | 0.00 | 0.3 |
| 0.035 | 0.00 | 0.36 |
| 0.040 | 0.00 | 0.4 |
| 0.046 | 0.00 | 0.47 |
| 0.052 | 0.00 | 0.55 |
| 0.060 | 0.00 | 0.63 |
| 0.069 | 0.00 | 0.72 |
| 0.079 | 0.00 | 0.83 |
| 0.091 | 0.00 | 0.95 |
| 0.105 | 0.00 | 1.09 |
| | | |

| ary 2012 4.33.37 PW | | | | | |
|--|--|--|--|--|--|
| e (µm) | Volume In % | | Size (µm) | Volume In % | |
| 0.105 0.120 0.138 0.158 0.182 | 0.00 0.00 0.00 0.00 | | 1.096 1.259 1.445 1.660 1.905 | 0.00 0.00 0.00 0.00 | |
| 0.209 0.240 0.275 0.316 0.363 0.417 0.479 0.550 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 | | 2.188 2.512 2.884 3.311 3.802 4.365 5.012 5.754 | 0.00 0.00 0.05 0.08 0.10 0.12 0.14 | |
| 0.631 0.724 0.832 0.955 1.096 | 0.00 0.00 0.00 0.00 0.00 | | 6.607 7.586 8.710 10.000 11.482 | 0.19 0.20 0.22 0.23 0.24 | |

| - | | |
|---|-----------|-------------|
| | Size (µm) | Volume In % |
| | 11.482 | 0.24 |
| | 13.183 | 0.24 |
| | 15.136 | |
| | 17.378 | 0.23 |
| | 19.953 | 0.21 |
| | 22,909 | 0.18 |
| | 26.303 | 0.16 |
| | 30.200 | 0.14 |
| | | 0.13 |
| | 34.674 | 0.14 |
| | 39.811 | 0.16 |
| | 45.709 | 0.20 |
| | 52.481 | 0.23 |
| | 60.256 | 0.26 |
| | 69.183 | |
| | 79.433 | 0.25 |
| | 91,201 | 0.21 |
| | 104.713 | 0.05 |
| | 400.000 | 0.00 |

| - 1 | o: / \ | 1.1.1.00 |
|-----|-----------|-------------|
| | Size (µm) | Volume In % |
| | 120.226 | 0.00 |
| | 138.038 | 0.00 |
| | 158.489 | 0.00 |
| | 181.970 | 0.01 |
| | 208,930 | 0.29 |
| | 239.883 | 0.90 |
| | 275.423 | 1.81 |
| | 316.228 | 2.96 |
| | 0.00 | 4.29 |
| | 363.078 | 5.63 |
| | 416.869 | 6.86 |
| | 478.630 | 7.83 |
| | 549.541 | 8.48 |
| | 630.957 | 0.10 |
| | 724.436 | 8.79 |
| | 831,764 | 8.79 |
| | 054.003 | 8.52 |

1096.478

1258.925

8.03

7.30

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 6.27 |
| 1445.440 | 4.89 |
| 1659.587 | 2.87 |
| 1905.461 | 0.70 |
| 2187.762 | 00 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311.311 | 0.00 |
| 3801.894 | 0.00 |
| 4365.158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | 0.00 |
| 10000.000 | |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

D6 - Average

Measured by:

Tuesday, 17 January 2012 5:16:05 PM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 5:16:07 PM

Averaged

Particle Name: Silica 1.45

Accessory Name: Hydro 2000G (A)

Sensitivity:

Particle RI:

Normal

1.450

Absorption:

Obscuration:

Dispersant Name:

0.1

Dispersant RI:

to 2000.000 um Weighted Residual:

12.39 % **Result Emulation:**

Water

0.0688

1.330

2.998

Size range:

0.020

Analysis model:

General purpose

Off

Concentration:

Span:

Uniformity:

Result units:

Specific Surface Area:

%Vol

3.000

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]:

Volume

0.297

20.195 um 545.749

d(0.1):

9.451

um


d(0.5):

429.527

um

d(0.9):

1298.067 um



| Size (µm) | Volume In % | Size (µm) | Volume In % | |
|-----------|-------------|-----------|-------------|--|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | | 0.120 | | |
| 0.013 | 0.00 | 0.138 | 0.00 | |
| 0.015 | 0.00 | 0.158 | 0.00 | |
| 0.017 | 0.00 | 0.182 | 0.00 | |
| 0.020 | 0.00 | 0.209 | 0.00 | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| | 0.00 | | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.00 | |
| 0.030 | 0.00 | 0.316 | 0.01 | |
| 0.035 | 0.00 | 0.363 | 0.08 | |
| 0.040 | 0.00 | 0.417 | 0.13 | |
| 0.046 | 0.00 | 0.479 | 0.16 | |
| 0.052 | | 0.550 | | |
| 0.060 | 0.00 | 0.631 | 0.18 | |
| 0.069 | 0.00 | 0.724 | 0.19 | |
| 0.079 | 0.00 | 0.832 | 0.18 | |
| 0.091 | 0.00 | 0.955 | 0.16 | |
| 0.031 | 0.00 | 1.006 | 0.14 | |

| | |
|--------------|-------------|
| Size (µm) | Volume In % |
| 1.096 | 0.12 |
| 1.259 | |
| 1.445 | 0.11 |
| 1.660 | 0.12 |
| 1.905 | 0.17 |
| 2.188 | 0.22 |
| 2.512 | 0.28 |
| 2.884 | 0.37 |
| 3.311 | 0.46 |
| 3.802 | 0.57 |
| | 0.68 |
| 4.365 | 0.79 |
| 5.012 | 0.90 |
| 5.754 | 1.00 |
| 6.607 | 1.09 |
| 7.586 | 1.17 |
| 8.710 | 1.22 |
| 10.000 | 1.26 |
| 11 //82 | 1.20 |

| - | | |
|---|-----------|-------------|
| | Size (µm) | Volume In % |
| | 11.482 | 1.27 |
| | 13.183 | |
| | 15.136 | 1.26 |
| | 17.378 | 1.23 |
| | 19.953 | 1.19 |
| | 22.909 | 1.14 |
| | 26.303 | 1.09 |
| | 30.200 | 1.03 |
| | 34.674 | 0.97 |
| | 39.811 | 0.91 |
| | | 0.84 |
| | 45.709 | 0.77 |
| | 52.481 | 0.69 |
| | 60.256 | 0.61 |
| | 69.183 | 0.55 |
| | 79.433 | 0.52 |
| | 91.201 | 0.52 |
| | 104.713 | |
| | 100 000 | 0.64 |

| Size (µm) | Volume In % |
|-----------|--------------|
| 120.226 | 0.83 |
| 138.038 | 1.11 |
| 158.489 | 1.48 |
| 181.970 | 1.40 |
| 208.930 | 2.42 |
| 239.883 | 2.42 |
| 275.423 | 3.39 |
| 316.228 | 3.80 |
| 363.078 | 3.60 4.14 |
| 416.869 | 4.14 |
| 478.630 | 4.40 4.62 |
| 549.541 | 4.62 |
| 630.957 | |
| 724,436 | 4.99 |

831.764

954.993

1096.478

1258.925

5.33

5.36

5.17

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 4.63 |
| 1445.440 | |
| 1659.587 | 3.71 |
| 1905.461 | 2.21 |
| 2187.762 | 0.54 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311 311 | 0.00 |
| 00111011 | 0.00 |
| 3801.894 | 0.00 |
| 4365.158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| 6606.934 | 5.55 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | 0.00 |
| 10000.000 | |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

A1 - Average

Measured by:

Tuesday, 17 January 2012 10:53:59 AM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 10:54:01 AM

Averaged

Particle Name:

Accessory Name: Silica 1.45 Hydro 2000G (A)

Sensitivity: Normal

Particle RI:

Absorption:

1.450

Obscuration:

Dispersant Name:

0.1

to 2000.000 um 13.88 %

d(0.1):

Dispersant RI:

Water

1.330

Weighted Residual: 5.323

Analysis model:

General purpose

Size range:

Result Emulation: Off

Concentration:

Span:

Uniformity:

0.020

Result units:

0.3346 %Vol 1.729

Volume

Specific Surface Area:

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]: 647.587

0.037

162.231 um

> 627.177 um

d(0.9):

1196.139 um

111.734

um

d(0.5):

Particle Size Distribution 100 90 80 70 60 50 40 30 20 10 0.01 10 100 1000 3000 Particle Size (µm) -A1 - Average, Tuesday, 17 January 2012 10:53:59 AM

| Size (µm) | Volume In % | Size (µm) | Volume In % | |
|-----------|-------------|-----------|-------------|--|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | |
| 0.013 | 0.00 | 0.138 | 0.00 | |
| 0.015 | | 0.158 | | |
| 0.017 | 0.00 | 0.182 | 0.00 | |
| 0.020 | 0.00 | 0.209 | 0.00 | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.00 | |
| 0.030 | 0.00 | 0.316 | 0.00 | |
| 0.035 | 0.00 | 0.363 | 0.00 | |
| 0.040 | 0.00 | 0.417 | 0.00 | |
| 0.046 | 0.00 | 0.417 | 0.00 | |
| | 0.00 | 1 1 | 0.00 | |
| 0.052 | 0.00 | 0.550 | 0.00 | |
| 0.060 | 0.00 | 0.631 | 0.00 | |
| 0.069 | 0.00 | 0.724 | 0.00 | |
| 0.079 | 0.00 | 0.832 | 0.00 | |
| 0.091 | 0.00 | 0.955 | 0.00 | |
| 0.105 | 0.00 | 1 006 | 0.00 | |

| Size (µm) | Volume In % | |
|-----------|-------------|--|
| 1.096 | 0.00 | |
| 1.259 | | |
| 1.445 | 0.00 | |
| 1.660 | 0.00 | |
| 1.905 | 0.00 | |
| 2.188 | 0.00 | |
| 2.512 | 0.00 | |
| 2.884 | 0.00 | |
| 3.311 | 0.05 | |
| 3.802 | 0.09 | |
| | 0.11 | |
| 4.365 | 0.13 | |
| 5.012 | 0.15 | |
| 5.754 | 0.17 | |
| 6.607 | 0.18 | |
| 7.586 | 0.20 | |
| 8.710 | 0.21 | |
| 10.000 | | |
| 11.482 | 0.22 | |

| Size (µm) | Volume In % |
|-----------|---|
| 11.482 | 0.24 |
| 13.183 | 0.24 |
| 15.136 | 0.24 |
| 17.378 | 0.24 |
| 19.953 | 0.23 |
| 22.909 | 0.21 |
| 26.303 | 0.16 |
| 30.200 | 0.13 |
| 34.674 | 0.13 |
| 39.811 | 0.13 |
| 45.709 | 0.21 |
| 52.481 | 0.52 |
| 60.256 | 0.52 |
| 69.183 | • |
| 79.433 | 1.06 |
| 91.201 | 1.36 |
| 104.713 | 1.62 |
| 120.226 | 1.81 |

| 1 | Size (µm) | Volume In % |
|---|-----------|-------------|
| | 120.226 | 1.90 |
| | 138.038 | 1.89 |
| | 158.489 | |
| | 181.970 | 1.80 |
| | 208.930 | 1.69 |
| | 239.883 | 1.64 |
| | 275.423 | 1.75 |
| | 316.228 | 2.11 |
| | 363.078 | 2.82 |
| | 416.869 | 3.86 |
| | 478.630 | 5.21 |
| | 549.541 | 6.68 |
| | 630.957 | 8.05 |
| | 724.436 | 9.05 |
| | 831.764 | 9.43 |
| | | 9.05 |
| | 954.993 | 7.93 |

1096.478

1258.925

6.29

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 4.46 |
| 1445.440 | |
| 1659.587 | 2.65 |
| 1905.461 | 0.77 |
| 2187.762 | 0.00 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311 311 | 0.00 |
| 3801.894 | 0.00 |
| 4365.158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

B1 - Average

Measured by:

Tuesday, 17 January 2012 11:13:59 AM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Analysis model:

General purpose

Averaged

Tuesday, 17 January 2012 11:14:01 AM

Particle Name:

Accessory Name: Hydro 2000G (A) Silica 1.45

Sensitivity: Normal

Particle RI: 1.450

Absorption:

Dispersant Name:

Size range: 0.020

Obscuration:

Dispersant RI:

to 2000.000 um 15.33 % Weighted Residual:

Water 1.330

um

1.416

Result Emulation: Off

Concentration:

d(0.1):

Span:

0.1

Uniformity:

Result units:

578.695

0.0505 %Vol 4.300

Volume

Specific Surface Area:

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]:

225.161

0.37


8.772

16.237 um

> d(0.5): 132.553 um

d(0.9):

um



| Size (µm) | Volume In % | Size (µm) | Volume In % | 97 |
|-----------|-------------|-----------|-------------|----|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | |
| 0.013 | 0.00 | 0.138 | 0.00 | |
| 0.015 | | 0.158 | | |
| 0.017 | 0.00 | 0.182 | 0.00 | |
| 0.020 | 0.00 | 0.209 | 0.00 | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.00 | |
| | 0.00 | | 0.00 | |
| 0.030 | 0.00 | 0.316 | 0.01 | |
| 0.035 | 0.00 | 0.363 | 0.13 | |
| 0.040 | 0.00 | 0.417 | 0.19 | |
| 0.046 | 0.00 | 0.479 | 0.23 | |
| 0.052 | | 0.550 | | |
| 0.060 | 0.00 | 0.631 | 0.25 | |
| 0.069 | 0.00 | 0.724 | 0.25 | |
| 0.079 | 0.00 | 0.832 | 0.24 | |
| 0.091 | 0.00 | 0.955 | 0.21 | |
| 0.091 | 0.00 | 1.006 | 0.17 | |

| Size (µm) | Volume In % | |
|-----------|-------------|--|
| 1.096 | 0.14 | |
| 1.259 | 0.13 | |
| 1.445 | 0.13 | |
| 1.660 | 0.14 | |
| 1.905 | 0.17 | |
| 2.188 | 0.23 | |
| 2.512 | | |
| 2.884 | 0.39 | |
| 3.311 | 0.50 | |
| 3.802 | 0.60 | |
| 4.365 | 0.71 | |
| 5.012 | 0.81 | |
| 5.754 | 0.91 | |
| 6.607 | 1.00 | |
| 7.586 | 1.08 | |
| 8,710 | 1.16 | |
| 10.000 | 1.21 | |
| 11.482 | 1.25 | |
| | | |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 1.27 |
| 13.183 | 1.27 |
| 15.136 | 1.24 |
| 17.378 | 1.24 |
| 19.953 | 1.17 |
| 22.909 | |
| 26.303 | 1.16 |
| 30.200 | 1.19 |
| 34.674 | 1.29 |
| 39.811 | 1.46 |
| 45.709 | 1.73 |
| 52,481 | 2.07 |
| 60.256 | 2.46 |
| 69.183 | 2.87 |
| 79.433 | 3.25 |
| 91.201 | 3.56 |
| 104.713 | 3.78 |
| | 3.88 |
| 120.226 | |

| Size (µm) | Volume In % |
|-----------|-------------|
| 120.226 | |
| 138.038 | 3.88 |
| 158.489 | 3.80 |
| 181.970 | 3.68 |
| 208,930 | 3.57 |
| 239.883 | 3.50 |
| 275,423 | 3.51 |
| 316.228 | 3.61 |
| 363.078 | 3.78 |
| 416.869 | 3.96 |
| 478.630 | 4.06 |
| 549.541 | 3.99 |
| 0.1010.11 | 3.65 |
| 630.957 | 3.03 |
| 724.436 | 2.11 |
| 831.764 | 1.15 |
| | |

1096.478

1258.925

0.75

0.49

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 0.24 |
| 1445.440 | 0.24 |
| 1659.587 | 0.00 |
| 1905.461 | 0.00 |
| 2187.762 | 5.55 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311.311 | 0.00 |
| 3801.894 | 0.00 |
| 4365,158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| | 0.00 |
| 10000.000 | |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

C1 - Average

Measured by:

Tuesday, 17 January 2012 11:35:01 AM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 11:35:03 AM

Averaged

Particle Name:

Accessory Name: Silica 1.45 Hydro 2000G (A)

Sensitivity: Normal

Particle RI: 1.450

Absorption:

Obscuration:

Dispersant Name:

0.1

to 2000.000 um

19.99 %

0.0595

d(0.1):

Dispersant RI:

Weighted Residual:

Analysis model:

General purpose

Result Emulation:

Water

1.330

1.804

Size range:

0.020

Off

Concentration:

Span:

um

Uniformity:

Result units:

%Vol

6.900

2.871

Vol. Weighted Mean D[4,3]:

Volume

Specific Surface Area:

Surface Weighted Mean D[3,2]:

d(0.5):

284.412

0.439


13.679 um

> 228.844 um

d(0.9):

663.849

um



| Size (µm) | Volume In % | Size (µm) | Volume In % | Size (µm) | Vo |
|-----------|-------------|-----------|-------------|-----------|----|
| 0.010 | 0.00 | 0.105 | 0.00 | 1.096 | |
| 0.011 | 0.00 | 0.120 | 0.00 | 1.259 | |
| 0.013 | 0.00 | 0.138 | 0.00 | 1.445 | |
| 0.015 | | 0.158 | | 1.660 | |
| 0.017 | 0.00 | 0.182 | 0.00 | 1.905 | |
| 0.020 | 0.00 | 0.209 | 0.00 | 2.188 | |
| 0.023 | 0.00 | 0.240 | 0.00 | 2.512 | |
| 0.026 | 0.00 | 0.275 | 0.00 | 2.884 | |
| 0.030 | 0.00 | 0.316 | 0.01 | 3.311 | |
| 0.035 | 0.00 | 0.363 | 0.08 | 3.802 | |
| 0.040 | 0.00 | 0.417 | 0.16 | 4.365 | |
| 0.046 | 0.00 | 0.479 | 0.23 | 5.012 | |
| 0.052 | 0.00 | 0.550 | 0.27 | 5.754 | |
| 0.060 | 0.00 | 0.631 | 0.30 | 6.607 | |
| 0.069 | 0.00 | 0.031 | 0.30 | 7.586 | |
| | 0.00 | - | 0.29 | | |
| 0.079 | 0.00 | 0.832 | 0.25 | 8.710 | |
| 0.091 | 0.00 | 0.955 | 0.22 | 10.000 | |
| | | | | | |

| 1 | Volume In % | Ì | Cine (um) | Volume In % |
|---|-------------|---|-----------|-------------|
| | volume in % | | | volume in % |
| | 0.19 | | 11.482 | 1.21 |
| | 0.18 | | 13.183 | 1.18 |
| | 0.20 | | 15.136 | 1.13 |
| | 0.25 | | 17.378 | 1.08 |
| | 0.32 | | 19.953 | 1.04 |
| | 0.41 | | 22.909 | 1.02 |
| | 0.52 | | 26.303 | 1.03 |
| | 0.64 | | 30.200 | 1.07 |
| | 0.76 | | 34.674 | 1.14 |
| | 0.87 | | 39.811 | 1.26 |
| | 0.97 | | 45.709 | 1.40 |
| | 1.06 | | 52.481 | 1.55 |
| | | | 60.256 | |
| | 1.13 | | 69.183 | 1.70 |
| | 1.19 | | 79.433 | 1.83 |
| | 1.22 | | 91.201 | 1.93 |
| | 1.24 | | 104.713 | 2.01 |
| | 1.23 | | 120.226 | 2.08 |
| | | | | |

| Size (µm) | Volume In % | Size (µm) | Volume In % |
|--|---|--|---|
| Size (µm) 120,226 138,038 158,489 181,970 208,930 239,883 275,423 316,228 363,078 416,869 478,630 549,541 630,957 724,436 831,764 9954,993 1096,478 1258,925 | Volume In % 2.16 2.29 2.49 2.81 3.25 3.81 4.46 5.13 5.70 6.06 6.09 5.68 4.84 3.64 2.22 0.92 0.29 | Size (µm) 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3801.894 4365.158 5011.872 5754.399 6606.934 7585.776 8709.636 10000.000 | Volume In % 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0 |
| 13/020 | | | |

Operator notes:





Sample Name:

SOP Name:

D1 - Average

Measured by:

Tuesday, 17 January 2012 11:54:27 AM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 11:54:29 AM

Averaged

Particle Name: Silica 1.45

Accessory Name:

Hydro 2000G (A)

Analysis model: General purpose

Sensitivity: Normal

Particle RI:

Absorption:

Size range:

Obscuration:

1.450

0.1

0.020 to 2000.000 um 17.49 %

Dispersant Name:

Dispersant RI:

Weighted Residual:

Result Emulation:

Water

1.059

Off

0.0440

1.330

Uniformity:

Result units:

Concentration: %Vol Span: 6.518

Volume

588.636

Specific Surface Area: 0.502

Surface Weighted Mean D[3,2]: 11.943

Vol. Weighted Mean D[4,3]: 202.146

d(0.1):

6.517

um

d(0.5):


um

89.308

um

d(0.9):

um



| Size (µm) | Volume In % | Size (µm) | Volume In % | 0) |
|-----------|-------------|-----------|-------------|----|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | |
| 0.013 | | 0.138 | | |
| 0.015 | 0.00 | 0.158 | 0.00 | |
| 0.017 | 0.00 | 0.182 | 0.00 | |
| 0.020 | 0.00 | 0.209 | 0.00 | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| | 0.00 | | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.01 | |
| 0.030 | 0.00 | 0.316 | 0.08 | |
| 0.035 | 0.00 | 0.363 | 0.18 | |
| 0.040 | 0.00 | 0.417 | 0.16 | |
| 0.046 | | 0.479 | | |
| 0.052 | 0.00 | 0.550 | 0.31 | |
| 0.060 | 0.00 | 0.631 | 0.34 | |
| 0.069 | 0.00 | 0.724 | 0.35 | |
| 0.009 | 0.00 | - | 0.32 | |
| | 0.00 | 0.832 | 0.28 | |
| 0.091 | 0.00 | 0.955 | 0.23 | |
| 0.405 | | | | |

| e (µm) | Volume In % | Size (µm) | Volume In % |
|--------|-------------|-----------|-------------|
| 1.096 | 0.19 | 11.482 | 1.82 |
| 1.259 | | 13.183 | |
| 1.445 | 0.17 | 15.136 | 1.83 |
| 1.660 | 0.18 | 17.378 | 1.83 |
| 1.905 | 0.22 | 19.953 | 1.82 |
| 2.188 | 0.28 | 22.909 | 1.81 |
| 2.512 | 0.38 | 26.303 | 1.84 |
| 2.884 | 0.49 | 30.200 | 1.90 |
| | 0.61 | | 2.01 |
| 3.311 | 0.75 | 34.674 | 2.16 |
| 3.802 | 0.90 | 39.811 | 2.35 |
| 4.365 | 1.05 | 45.709 | 2.55 |
| 5.012 | 1.20 | 52.481 | 2.75 |
| 5.754 | 1.35 | 60.256 | 2.92 |
| 6.607 | 1.49 | 69.183 | 3.04 |
| 7.586 | | 79.433 | |
| 8.710 | 1.61 | 91.201 | 3.12 |
| 10.000 | 1.71 | 104.713 | 3.15 |
| 11.482 | 1.78 | 120.226 | 3.15 |
| | | | |

| Size (µm) | Volume In % |
|-----------|--|
| 1258.925 | 0.48 |
| 1445.440 | 0.40 |
| 1659.587 | 0.02 |
| 1905.461 | 5.55 |
| 2187.762 | 0.00 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| | 0.00 |
| | 0.00 |
| | 0.00 |
| | 0.00 |
| | 0.00 |
| | 0.00 |
| | 0.00 |
| | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | 0.00 |
| | |
| | |
| | 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3310.391 4365.158 5011.872 5754.399 6606.934 7585.776 8709.636 |

Operator notes:

Malvern Instruments Ltd. Tel := +[44] (0) 1684-892456 Fax +[44] (0) 1684-892789

Mastersizer 2000 Ver. 5.60 Serial Number: MAL1006765 File name: SKM 170112.mea Record Number: 20 18/01/2012 10:53:28 4





Sample Name:

SOP Name:

A4 - Average

Measured by:

Tuesday, 17 January 2012 12:18:14 PM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 12:18:16 PM

Averaged

Particle Name: Silica 1.45

Accessory Name: Hydro 2000G (A)

Analysis model: General purpose

Sensitivity: Normal

Particle RI:

Absorption:

Size range:

Obscuration:

1.450

0.1

0.020 to 2000.000 um 12.04 %

Dispersant Name:

Concentration:

Dispersant RI:

Weighted Residual:

Result Emulation:

Water

4.757

Off

d(0.9):

1.330

Uniformity:

Result units:

0.4640

d(0.1):

Span: %Vol 1.623

Volume

Specific Surface Area:

185.391

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]: 636.603

250.903 0.0239


um

um

d(0.5):

um

1148.302 um



593.174

| 0.010 0.011 0.013 0.015 0.017 0.020 0.023 0.026 0.030 | Volume In % 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0 | 0.105 0.120 0.138 0.158 0.182 0.209 0.240 0.275 0.316 0.363 | Volume In % 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0 | 1.096 1.259 1.445 1.660 1.905 2.188 2.512 2.884 3.311 3.802 | Volume In % 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0 | Size (μm) 11.482 13.183 15.136 17.378 19.953 22.909 26.303 30.200 34.674 39.811 | Volume In % 0.14 0.15 0.15 0.16 0.16 0.15 0.14 0.12 0.11 0.10 | 120.226 138.038 158.489 181.970 208.930 239.883 275.423 316.228 363.078 416.869 | 1.23 1.48 1.73 2.02 2.40 2.91 3.59 4.48 5.55 6.72 | 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3801.894 4365.158 | Volume In % 3.86 2.30 0.33 0.01 0.00 0.00 0.00 0.00 0.00 0 | |
|---|---|--|---|--|---|---|--|--|--|--|---|--|
| 0.023 | 0.00 0.00 0.00 | 0.240 0.275 0.316 | 0.00 0.00 0.00 | 2.512 2.884 3.311 | 0.00 0.00 0.00 | 26.303 30.200 34.674 | 0.14 0.12 0.11 | 275.423 316.228 363.078 | 3.59 4.48 5.55 | 2884.032 3311.311 3801.894 | 0.00 0.00 0.00 | |

Operator notes:





Sample Name: B4 - Average

SOP Name:

Tuesday, 17 January 2012 12:31:36 PM

Sample Source & type:

Measured by:

Analysed:

abbottsj

Tuesday, 17 January 2012 12:31:37 PM

Sample bulk lot ref:

Result Source: Averaged

Particle Name:

Accessory Name:

Sensitivity:

Silica 1.45

Hydro 2000G (A)

Normal

Particle RI:

Absorption:

General purpose

1.450

Obscuration: 19.63 %

Dispersant Name:

Concentration:

d(0.1):

0.1 **Dispersant RI:** to 2000.000 um

d(0.9):

Water

Weighted Residual: 2.114

Analysis model:

Size range:

Result Emulation:

1.330

0.020

Off

Span:

Uniformity:

Result units: Volume

855.563

0.0778 %Vol 2.844

Surface Weighted Mean D[3,2]:

d(0.5):

Vol. Weighted Mean D[4,3]: 370.929

Specific Surface Area: 0.328


10.615

um

18.298

297.105 um

um



| Size (µm) | Volume In % | Size (µm) | Volume In % | |
|-----------|-------------|-----------|-------------|---|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | l |
| 0.013 | | 0.138 | | l |
| 0.015 | 0.00 | 0.158 | 0.00 | ì |
| 0.017 | 0.00 | 0.182 | 0.00 | l |
| 0.020 | 0.00 | 0.209 | 0.00 | l |
| 0.023 | 0.00 | 0.240 | 0.00 | ì |
| 0.026 | 0.00 | 0.275 | 0.00 | ì |
| 0.020 | 0.00 | 0.275 | 0.00 | l |
| | 0.00 | | 0.07 | ì |
| 0.035 | 0.00 | 0.363 | 0.13 | ì |
| 0.040 | 0.00 | 0.417 | 0.17 | Ì |
| 0.046 | 0.00 | 0.479 | 0.21 | Ì |
| 0.052 | 0.00 | 0.550 | 0.23 | l |
| 0.060 | 0.00 | 0.631 | 0.23 | l |
| 0.069 | | 0.724 | 0.23 | l |
| 0.079 | 0.00 | 0.832 | | l |
| 0.091 | 0.00 | 0.955 | 0.18 | l |
| | 0.00 | | 0.15 | ì |

| Size (µm) | Volume In % |
|-----------|-------------|
| 1.096 | 0.13 |
| 1.259 | |
| 1.445 | 0.12 |
| 1.660 | 0.13 |
| 1.905 | 0.16 |
| 2,188 | 0.21 |
| 2.512 | 0.28 |
| 2.884 | 0.36 |
| 3.311 | 0.44 |
| 3.802 | 0.53 |
| 4.365 | 0.61 |
| | 0.69 |
| 5.012 | 0.77 |
| 5.754 | 0.83 |
| 6.607 | 0.88 |
| 7.586 | 0.92 |
| 8.710 | 0.94 |
| 10.000 | 0.95 |
| 11 //82 | 0.33 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 0.95 |
| 13.183 | |
| 15.136 | 0.93 |
| 17.378 | 0.89 |
| 19.953 | 0.85 |
| 22.909 | 0.82 |
| 26,303 | 0.79 |
| 30.200 | 0.78 |
| 34.674 | 0.81 |
| 39.811 | 0.87 |
| 45.709 | 0.97 |
| 52.481 | 1.11 |
| | 1.28 |
| 60.256 | 1.46 |
| 69.183 | 1.65 |
| 79.433 | 1.83 |
| 91.201 | 1.99 |
| 104.713 | 2.15 |
| 120,226 | 2.10 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 120.226 | 2.30 |
| 138.038 | 2.46 |
| 158.489 | 2.40 |
| 181.970 | 2.91 |
| 208.930 | 3.22 |
| 239.883 | 3.61 |
| 275.423 | 4.07 |
| 316.228 | 4.57 |
| 363.078 | 5.06 |
| 416.869 | 5.48 |
| 478.630 | 5.75 |
| 549.541 | 5.79 |
| 630.957 | 5.54 |
| 724.436 | 4.99 |
| 831.764 | 4.19 |
| 954.993 | 3 21 |

1096.478

1258.925

3.21

2.20

| S | Size (µm) | Volume In % |
|---|-----------|-------------|
| | 1258.925 | 1.24 |
| | 1445.440 | 0.09 |
| | 1659.587 | 0.09 |
| | 1905.461 | 0.00 |
| | 2187.762 | 0.00 |
| | 2511.886 | 0.00 |
| | 2884.032 | 0.00 |
| | 3311.311 | 0.00 |
| | 3801.894 | 0.00 |
| | 4365.158 | 0.00 |
| | 5011.872 | 0.00 |
| | 5754.399 | 0.00 |
| | 6606.934 | 0.00 |
| | 7585.776 | 0.00 |
| | 8709.636 | 0.00 |
| | | 0.00 |
| 1 | 0000.000 | |
| | | |
| | | |

Operator notes:





Sample Name:

SOP Name:

C4 - Average

Measured by:

Tuesday, 17 January 2012 1:40:57 PM

Sample Source & type:

abbottsj

Averaged

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 1:40:58 PM

Particle Name: Accessory Name: Hydro 2000G (A) Silica 1.45

Sensitivity: Normal

Particle RI: 1.450

0.0926

Absorption:

Dispersant Name:

Size range: 0.020

Obscuration: 22.36 %

Dispersant RI:

to 2000.000 um Weighted Residual:

Water 1.330 2.813

Analysis model:

General purpose

Result Emulation: Off

Concentration:

Span:

0.1

Uniformity:

Result units:

2.725

Vol. Weighted Mean D[4,3]: Surface Weighted Mean D[3,2]:

Volume

Specific Surface Area: 0.327

%Vol

18.372 um

457.101


d(0.1): 11.695 um

d(0.5):

382.678

um

d(0.9): 1054.336 um



| Size (µm) | Volume In % | | Size (µm) | Volume In % | |
|-----------|-------------|---|-----------|-------------|---|
| 0.010 | 0.00 | | 0.105 | 0.00 | |
| 0.011 | 0.00 | | 0.120 | 0.00 | |
| 0.013 | | | 0.138 | | |
| 0.015 | 0.00 | | 0.158 | 0.00 | |
| 0.017 | 0.00 | | 0.182 | 0.00 | |
| 0.020 | 0.00 | | 0.209 | 0.00 | |
| 0.023 | 0.00 | | 0.240 | 0.00 | |
| 0.026 | 0.00 | | 0.275 | 0.00 | |
| 0.030 | 0.00 | | 0.316 | 0.01 | |
| 0.035 | 0.00 | | 0.363 | 0.07 | |
| 0.040 | 0.00 | | 0.417 | 0.13 | |
| 0.040 | 0.00 | | 0.417 | 0.18 | |
| | 0.00 | | | 0.21 | |
| 0.052 | 0.00 | | 0.550 | 0.23 | |
| 0.060 | 0.00 | | 0.631 | 0.23 | |
| 0.069 | 0.00 | | 0.724 | 0.22 | |
| 0.079 | 0.00 | | 0.832 | 0.19 | |
| 0.091 | 0.00 | | 0.955 | 0.16 | |
| 0.405 | 0.00 | 1 | 4 000 | 0.10 | l |

| Size (µm) | Volume In % |
|-----------|-------------|
| 1.096 | 0.14 |
| 1.259 | • |
| 1.445 | 0.12 |
| 1.660 | 0.13 |
| 1.905 | 0.16 |
| 2.188 | 0.20 |
| | 0.26 |
| 2.512 | 0.33 |
| 2.884 | 0.40 |
| 3.311 | 0.48 |
| 3.802 | 0.55 |
| 4.365 | 5.55 |
| 5.012 | 0.62 |
| 5.754 | 0.69 |
| 6.607 | 0.75 |
| 7.586 | 0.80 |
| | 0.84 |
| 8.710 | 0.87 |
| 10.000 | 0.89 |
| 11.482 | 0.03 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 0.90 |
| 13.183 | |
| 15.136 | 0.89 |
| 17.378 | 0.88 |
| 19.953 | 0.86 |
| 22,909 | 0.85 |
| 26.303 | 0.85 |
| 30.200 | 0.89 |
| 34.674 | 0.95 |
| 39.811 | 1.06 |
| 45.709 | 1.21 |
| | 1.37 |
| 52.481 | 1.55 |
| 60.256 | 1.72 |
| 69.183 | 1.85 |
| 79.433 | 1.95 |
| 91.201 | 1.99 |
| 104.713 | 1.99 |
| 120,226 | 1.99 |

| Size (um) | Volume In % |
|-----------|-------------|
| 120.226 | |
| 138.038 | 1.96 |
| 158,489 | 1.91 |
| 181.970 | 1.87 |
| 208.930 | 1.86 |
| 239.883 | 1.93 |
| 275.423 | 2.12 |
| 316.228 | 2.45 |
| 363.078 | 2.96 |
| 416.869 | 3.63 |
| 478.630 | 4.41 |
| | 5.20 |
| 549.541 | 5.87 |
| 630.957 | 6.28 |
| 724.436 | 6.30 |

831.764

954.993

1096.478

1258.925

5.88

5.06

3.96

| Size (µm) | Volume In % |
|--|---|
| 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3801.894 4365.158 5011.872 5754.399 | Volume In % 2.77 1.68 0.25 0.01 0.00 0.00 0.00 0.00 0.00 0.00 0.0 |
| 5754.399 6606.934 | 0.00 |
| 6606.934 7585.776 8709.636 | 0.00 |
| 10000.000 | 0.00 |

Operator notes:





Sample Name:

SOP Name:

D4 - Average

Measured by:

Tuesday, 17 January 2012 2:10:11 PM

Sample Source & type:

abbottsj

Averaged

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 2:10:13 PM

Particle Name: Accessory Name: Hydro 2000G (A) Silica 1.45

General purpose

Sensitivity: Normal

Particle RI:

Absorption:

Size range:

Obscuration:

1.450

0.1

0.020 to 2000.000 um Weighted Residual:

16.69 %

Dispersant Name: Water

Dispersant RI:

Analysis model:

Result Emulation:

0.0526

1.330

2.435

Off

Concentration:

Span:

Uniformity:

Result units: Volume

Specific Surface Area:

%Vol

3.413

Vol. Weighted Mean D[4,3]:

0.487

Surface Weighted Mean D[3,2]: 12.323 um

367.026

d(0.1): 6.026


um

d(0.5):

267.250

um

d(0.9): 918.145 um



| Size (µm) 0.010 0.011 0.013 0.015 0.017 0.020 0.023 0.026 0.030 0.035 0.040 0.046 0.052 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | Size (µm) 0.105 0.120 0.138 0.158 0.182 0.209 0.240 0.275 0.316 0.363 0.417 0.479 0.550 0.631 | 0.00 0.00 0.00 0.00 0.00 0.00 0.01 0.07 0.16 0.24 0.30 | Size (µm) 1.096 1.259 1.445 1.660 1.905 2.188 2.512 2.884 3.311 3.802 4.365 5.012 5.754 6.607 | 0.23 0.22 0.23 0.28 0.36 0.46 0.58 0.71 0.84 0.98 1.11 | Size (µm) 11.482 13.183 15.136 17.378 19.953 22.909 26.303 30.200 34.674 39.811 45.709 52.481 60.256 69.183 | 1.44 1.40 1.35 1.31 1.29 1.32 1.38 1.47 1.56 1.65 | 120,226 138,038 158,489 181,970 208,930 239,883 275,423 316,228 363,078 416,869 478,630 549,541 630,957 | Volume In % 1.24 1.19 1.23 1.38 1.66 2.12 2.72 3.42 4.14 4.80 5.27 5.50 5.42 | 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3801.894 4365.158 5011.872 5754.399 6606.934 | Volume in % 1.75 0.99 0.07 0.00 0.00 0.00 0.00 0.00 0.00 |
|---|---|---|--|---|--|---|--|---|---|--|---|
| | 0.00 | | 0.30 | | 1.23 1.32 1.40 1.45 | | 1.71 1.73 1.69 1.60 1.47 | | 5.50 | | 0.00 |

Operator notes:





Sample Name:

SOP Name:

A5 - Average

Measured by:

Tuesday, 17 January 2012 2:25:54 PM

Sample Source & type:

abbottsj

Averaged

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 2:25:56 PM

Particle Name: Accessory Name: Hydro 2000G (A) Silica 1.45

Analysis model: General purpose

Sensitivity: Normal

Particle RI:

Absorption:

Size range:

Obscuration:

1.450 **Dispersant Name:** 0.1

0.020 to 2000.000 um 13.48 %

Water

1.0423

d(0.1):

Dispersant RI:

Weighted Residual:

Result Emulation:

5.186

1.330

Off

Concentration:

%Vol

320.863

um

Span:

Uniformity:

Result units: Volume

Specific Surface Area:

1.103

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]: 593.948

0.0118

506.982 um


d(0.5):

553.821

um

d(0.9): 931.814

um



| Size (µIII) | VOIGITIE III 70 | Size (µIII) | VOIGITIE III 70 |
|-------------|-----------------|-------------|-----------------|
| 0.010 | 0.00 | 0.105 | 0.00 |
| 0.011 | 0.00 | 0.120 | 0.00 |
| 0.013 | | 0.138 | |
| 0.015 | 0.00 | 0.158 | 0.00 |
| 0.017 | 0.00 | 0.182 | 0.00 |
| 0.020 | 0.00 | 0.209 | 0.00 |
| 0.023 | 0.00 | 0.240 | 0.00 |
| 0.026 | 0.00 | 0.275 | 0.00 |
| 0.020 | 0.00 | 0.275 | 0.00 |
| 0.035 | 0.00 | 0.363 | 0.00 |
| | 0.00 | | 0.00 |
| 0.040 | 0.00 | 0.417 | 0.00 |
| 0.046 | 0.00 | 0.479 | 0.00 |
| 0.052 | 0.00 | 0.550 | 0.00 |
| 0.060 | 0.00 | 0.631 | 0.00 |
| 0.069 | 0.00 | 0.724 | 0.00 |
| 0.079 | 0.00 | 0.832 | 0.00 |
| 0.091 | | 0.955 | |
| 0.105 | 0.00 | 1.096 | 0.00 |

| Size (µm) | Volume In % | |
|-----------|-------------|--|
| 1.096 | 0.00 | |
| 1.259 | 0.00 | |
| 1.445 | 0.00 | |
| 1.660 | 0.00 | |
| 1.905 | 0.00 | |
| 2.188 | | |
| 2.512 | 0.00 | |
| 2.884 | 0.00 | |
| 3.311 | 0.00 | |
| 3.802 | 0.00 | |
| 4.365 | 0.00 | |
| 5.012 | 0.00 | |
| 5.754 | 0.00 | |
| 6.607 | 0.00 | |
| 7.586 | 0.00 | |
| 8.710 | 0.00 | |
| 10.000 | 0.00 | |
| | 0.00 | |
| 11.482 | | |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 0.00 |
| 13.183 | 0.00 |
| 15.136 | |
| 17.378 | 0.00 |
| 19.953 | 0.00 |
| 22,909 | 0.00 |
| 26.303 | 0.00 |
| 30.200 | 0.00 |
| 34.674 | 0.00 |
| 39.811 | 0.00 |
| 45.709 | 0.00 |
| | 0.00 |
| 52.481 | 0.00 |
| 60.256 | 0.00 |
| 69.183 | 0.00 |
| 79.433 | 0.00 |
| 91.201 | 0.00 |
| 104.713 | 0.00 |
| 400 000 | 0.00 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 120.226 | 0.00 |
| 138.038 | |
| 158.489 | 0.00 |
| 181.970 | 0.02 |
| 208.930 | 0.23 |
| 239.883 | 1.32 |
| 275,423 | 2.89 |
| 316.228 | 4.89 |
| 363.078 | 7.17 |
| 416.869 | 9.36 |
| 478.630 | 11.17 |
| | 12.26 |
| 549.541 | 12.43 |
| 630.957 | 11.65 |
| 724.436 | 10.02 |
| 831.764 | 7.81 |
| 95/1 993 | |

1096.478

1258.925

5.39

2.98

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 0.42 |
| 1445.440 | 0.00 |
| 1659.587 | 0.00 |
| 1905.461 | 5.55 |
| 2187.762 | 0.00 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311.311 | 0.00 |
| 3801.894 | 0.00 |
| 4365.158 | 0.00 |
| 5011.872 | 0.00 |
| | 0.00 |
| 5754.399 | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | 0.00 |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

B5 - Average

Measured by:

Tuesday, 17 January 2012 2:56:04 PM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Averaged

Tuesday, 17 January 2012 2:56:06 PM

Particle Name: Accessory Name: Hydro 2000G (A) Silica 1.45

Sensitivity: Normal

Particle RI: 1.450

Absorption:

Dispersant Name:

Obscuration: to 2000.000 um

0.1

0.020

15.04 %

Water

Dispersant RI:

Weighted Residual:

Analysis model:

General purpose

Size range:

Result Emulation:

0.1782

0.0748

d(0.1):

1.330

3.362

Off

Concentration:

Span:

Uniformity:

Result units: Volume

Specific Surface Area:

%Vol

2.039

Surface Weighted Mean D[3,2]: 80.228

um

Vol. Weighted Mean D[4,3]:


500.129

59.017 um d(0.5):

458.886

um

d(0.9): 994.662 um



| 0.01 0.01 0.01 0.01 0.02 0.02 0.02 0.03 0.03 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | 0.105 0.120 0.138 0.158 0.182 0.209 0.240 0.275 0.316 0.363 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | 1.096 1.259 1.445 1.660 1.905 2.188 2.512 2.884 3.311 3.802 4.365 | 0.00 0.00 0.00 0.00 0.07 0.10 0.13 0.18 0.23 0.27 | 11.482 13.183 15.136 17.378 19.953 22.909 26.303 30.200 34.674 39.811 45.709 | Volume In % 0.49 0.48 0.48 0.46 0.45 0.44 0.45 0.47 0.51 0.57 | 120.226 138.038 158.489 181.970 208.930 239.883 275.423 316.228 363.078 416.869 478.630 | 1.72 1.92 2.16 2.47 2.88 3.41 4.08 4.88 5.75 6.60 7.30 | Size (µm) 1258.925 1445.440 1659.587 1905.461 2187.762 2511.886 2884.032 3311.311 3801.894 4365.158 5011.872 | ' |
|--|---|--|---|---|--|--|--|---|--|--|---|
| 0.02 0.03 0.03 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | 0.275 0.316 0.363 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | 2.884 3.311 3.802 | 0.13 0.18 0.23 0.27 0.32 0.36 0.40 0.43 0.45 | 30.200 34.674 39.811 | 0.45 0.47 0.51 0.57 | 316.228 363.078 416.869 | 4.08 4.88 5.75 6.60 | 3311.311 3801.894 4365.158 | |

Operator notes:

2.22 0.74 0.03 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00





Sample Name:

SOP Name:

C5 - Average

Tuesday, 17 January 2012 3:07:35 PM

Sample Source & type:

Measured by:

Analysed:

abbottsj

Tuesday, 17 January 2012 3:07:36 PM

Sample bulk lot ref:

Particle Name:

Result Source:

Averaged

Analysis model:

Sensitivity:

Silica 1.45

Accessory Name: Hydro 2000G (A)

General purpose Normal

Particle RI: 1.450

Absorption:

Size range: to 2000.000 um

Obscuration: 14.77 %

Dispersant Name:

Concentration:

0.1

Weighted Residual:

Water

Dispersant RI:

Result Emulation:

1.330

3.315

Off

0.2673

Span:

Uniformity:

0.020

Result units: Volume

Specific Surface Area:

%Vol

1.859

Vol. Weighted Mean D[4,3]:

0.0508

118.227 um 493.829

d(0.1): 109.265 um

d(0.5):

Surface Weighted Mean D[3,2]:


448.521

um

d(0.9):

943.215

um



| Size (µm) | Volume In % | Size (µn |) Volume In % | Size (µm) | Volume In % | Size (µm) | Volume In % | Size (|
|-----------|-------------|----------|---------------|-----------|-------------|-----------|-------------|--------|
| 0.010 | 0.00 | 0.10 | 0.00 | 1.096 | 0.00 | 11.482 | 0.33 | 120. |
| 0.011 | | 0.12 | 0 | 1.259 | | 13.183 | | 138. |
| 0.013 | 0.00 | 0.13 | 0.00 | 1.445 | 0.00 | 15.136 | 0.33 | 158. |
| 0.015 | 0.00 | 0.15 | 0.00 | 1.660 | 0.00 | 17.378 | 0.33 | 181. |
| 0.017 | 0.00 | 0.18 | 0.00 | 1.905 | 0.00 | 19.953 | 0.33 | 208. |
| 0.020 | 0.00 | 0.20 | 0.00 | 2.188 | 0.00 | 22.909 | 0.32 | 239. |
| 0.023 | 0.00 | 0.24 | 0.00 | 2.512 | 0.02 | 26.303 | 0.32 | 275. |
| | 0.00 | | 0.00 | | 0.07 | | 0.32 | |
| 0.026 | 0.00 | 0.27 | 0.00 | 2.884 | 0.10 | 30.200 | 0.33 | 316. |
| 0.030 | 0.00 | 0.31 | 0.00 | 3.311 | 0.13 | 34.674 | 0.34 | 363. |
| 0.035 | 0.00 | 0.36 | 0.00 | 3.802 | 0.16 | 39.811 | 0.37 | 416. |
| 0.040 | 0.00 | 0.41 | 7 0.00 | 4.365 | 0.19 | 45.709 | 0.42 | 478. |
| 0.046 | | 0.47 | 9 | 5.012 | | 52.481 | | 549. |
| 0.052 | 0.00 | 0.55 | 0.00 | 5.754 | 0.22 | 60.256 | 0.48 | 630. |
| 0.060 | 0.00 | 0.63 | 0.00 | 6.607 | 0.24 | 69.183 | 0.57 | 724. |
| 0.069 | 0.00 | 0.72 | 0.00 | 7.586 | 0.27 | 79.433 | 0.68 | 831. |
| 0.079 | 0.00 | 0.83 | 0.00 | 8.710 | 0.29 | 91.201 | 0.83 | 954. |
| | 0.00 | | 0.00 | | 0.30 | | 1.02 | |
| 0.091 | 0.00 | 0.95 | 0.00 | 10.000 | 0.32 | 104.713 | 1.26 | 1096. |
| 0.105 | | 1.09 | Ö | 11.482 | | 120.226 | | 1258. |

(µm) Volume In % Size (µm) Volume In % 1.74 1445,440 3.038 1.94 0.15 3.489 1659.587 2.41 0.00 1.970 1905.461 2.98 0.00 3.930 2187.762 3.65 0.00 2511.886 9.883 4.41 0.00 5.423 2884.032 5.23 0.00 5.228 3311.311 6.09 0.00 3801.894 3.078 6.89 0.00 6.869 4365.158 7.57 0.00 3.630 5011.872 8.01 0.00 5754.399 9.541 8.10 0.00 0.957 6606.934 0.00 7 79 1.436 7.04 0.00 .764 8709.636 5.91 0.00 1.993 10000.000 4.53 6.478 3.925

Operator notes:





Sample Name: D5 - Average

SOP Name:

Tuesday, 17 January 2012 3:19:01 PM

Sample Source & type:

Measured by: Analysed:

Sample bulk lot ref:

abbottsj **Result Source:**

Tuesday, 17 January 2012 3:19:03 PM

Averaged

Particle Name: Silica 1.45

Accessory Name:

Analysis model: General purpose

Sensitivity: Normal

Particle RI:

Hydro 2000G (A) Absorption:

Size range:

Obscuration:

1.450

0.1

0.020 to 2000.000 um 12.49 %

Dispersant Name:

Dispersant RI:

Weighted Residual:

Result Emulation:

Water

1.330

3.547

Off

Concentration:

Span:

Uniformity:

Result units: Volume

0.1342 %Vol 1.944


Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]: 511.165

Specific Surface Area: 0.0813

73.819 um

um



D5 - Average, Tuesday, 17 January 2012 3:19:01 PM

| Size (µm) | Volume In % | Size (µm) | Volume In % | |
|-----------|-------------|-----------|-------------|--|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | |
| 0.013 | 0.00 | 0.138 | 0.00 | |
| 0.015 | 0.00 | 0.158 | 0.00 | |
| 0.017 | 0.00 | 0.182 | 0.00 | |
| 0.020 | 0.00 | 0.209 | 0.00 | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.00 | |
| 0.030 | 0.00 | 0.316 | 0.00 | |
| 0.035 | 0.00 | 0.363 | 0.00 | |
| 0.040 | 0.00 | 0.417 | 0.00 | |
| 0.046 | 0.00 | 0.479 | 0.00 | |
| 0.052 | 0.00 | 0.550 | 0.00 | |
| 0.060 | 0.00 | 0.631 | 0.00 | |
| 0.069 | 0.00 | 0.724 | 0.00 | |
| 0.079 | 0.00 | 0.832 | 0.00 | |
| 0.091 | 0.00 | 0.955 | 0.00 | |
| 0 105 | | 1 096 | | |

| 012 3. | 17.01110 |
|-----------|-------------|
| Size (µm) | Volume In % |
| 1.096 | 0.00 |
| 1.259 | 0.00 |
| 1.445 | 0.00 |
| 1.660 | 0.06 |
| 1.905 | 0.09 |
| 2.188 | 0.03 |
| 2.512 | 0.13 |
| 2.884 | 0.16 |
| 3.311 | 0.23 |
| 3.802 | 0.28 |
| 4.365 | |
| 5.012 | 0.38 |
| 5.754 | 0.41 |
| 6.607 | 0.44 |
| 7.586 | 0.46 |
| 8.710 | 0.47 |
| 10.000 | 0.47 |
| 11.482 | 0.46 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 0.44 |
| 13.183 | |
| 15.136 | 0.42 |
| 17.378 | 0.40 |
| 19.953 | 0.37 |
| 22,909 | 0.36 |
| 26.303 | 0.35 |
| 30.200 | 0.36 |
| 34.674 | 0.38 |
| 39.811 | 0.43 |
| | 0.49 |
| 45.709 | 0.58 |
| 52.481 | 0.67 |
| 60.256 | 0.78 |
| 69.183 | 0.90 |
| 79.433 | 1.02 |
| 91.201 | 1.13 |
| 104.713 | |
| 400 000 | 1.26 |

| Size (µm) | Volume In % |
|-----------|--------------|
| 120.226 | 1.41 |
| 138.038 | 1.60 |
| 158.489 | 1.86 |
| 181.970 | 2.22 |
| 208.930 | 2.71 |
| 239.883 | 3.36 |
| 275.423 | 4.15 |
| 316.228 | 5.09 |
| 363.078 | 6.08 |
| 416.869 | 7.02 |
| 478.630 | 7.77 |
| 549.541 | 8.19 |
| 630.957 | 8.15 |
| 724.436 | 7.59 |
| 831.764 | 7.55 6.55 |

954.993

1096.478

1258.925

5.16

3.65

| Sizo (um) | Volume In % |
|-----------|-----------------|
| 1258,925 | VOIUITIE III 76 |
| | 2.16 |
| 1445.440 | 0.54 |
| 1659.587 | 0.00 |
| 1905.461 | 0.00 |
| 2187.762 | 0.00 |
| 2511.886 | 5.55 |
| 2884.032 | 0.00 |
| 3311.311 | 0.00 |
| 3801.894 | 0.00 |
| 4365.158 | 0.00 |
| | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | |
| 10000.000 | 0.00 |
| | |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

B6 - Average

Measured by:

Tuesday, 17 January 2012 3:31:12 PM

Sample Source & type:

abbottsj

Analysed:

Sample bulk lot ref:

Result Source:

Tuesday, 17 January 2012 3:31:14 PM

Averaged

Particle Name: Silica 1.45

Accessory Name:

Analysis model: Sensitivity: Normal

Particle RI:

Hydro 2000G (A) Absorption:

General purpose

1.450

0.020 to 2000.000 um Obscuration: 11.59 %

Dispersant Name:

0.1

Dispersant RI:

Weighted Residual:

Result Emulation:

Water

1.330

6.876

Size range:

Off

Concentration:

Span:

Uniformity:

Result units:

0.2626 %Vol 1.699

Surface Weighted Mean D[3,2]:

Vol. Weighted Mean D[4,3]:

Volume

Specific Surface Area: 0.139

43.246 um

718.950

d(0.1): 87.251


um

d(0.5):

705.996

um

d(0.9): 1286.731 um



| Size (µm) | Volume In % | Size (µm) | Volume In % | |
|-----------|-------------|-----------|-------------|--|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | |
| 0.013 | 0.00 | 0.138 | 0.00 | |
| 0.015 | 0.00 | 0.158 | 0.00 | |
| 0.017 | 0.00 | 0.182 | 0.00 | |
| 0.020 | | 0.209 | | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.00 | |
| 0.030 | 0.00 | 0.316 | 0.00 | |
| 0.035 | 0.00 | 0.363 | 0.00 | |
| 0.040 | 0.00 | 0.417 | 0.07 | |
| 0.046 | 0.00 | 0.479 | 0.09 | |
| 0.052 | 0.00 | 0.550 | 0.10 | |
| 0.060 | 0.00 | 0.631 | 0.11 | |
| 0.069 | 0.00 | 0.724 | 0.11 | |
| 0.079 | 0.00 | 0.832 | 0.11 | |
| 0.079 | 0.00 | 0.955 | 0.09 | |
| 0.091 | 0.00 | 1.006 | 0.08 | |

| , . _ | , . |
|------------------|-------------|
| Size (µm) | Volume In % |
| 1.096 | 0.06 |
| 1.259 | |
| 1.445 | 0.05 |
| 1.660 | 0.05 |
| 1.905 | 0.05 |
| 2.188 | 0.06 |
| 2.512 | 0.08 |
| 2.884 | 0.10 |
| 3.311 | 0.12 |
| 3.802 | 0.14 |
| 4.365 | 0.17 |
| | 0.20 |
| 5.012 | 0.25 |
| 5.754 | 0.28 |
| 6.607 | 0.31 |
| 7.586 | 0.34 |
| 8.710 | 0.37 |
| 10.000 | 0.39 |
| 11 //82 | 0.39 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 0.42 |
| 13.183 | 0.42 |
| 15.136 | 51.15 |
| 17.378 | 0.44 |
| 19.953 | 0.44 |
| 22,909 | 0.44 |
| 26,303 | 0.42 |
| 30.200 | 0.41 |
| 34.674 | 0.39 |
| | 0.38 |
| 39.811 | 0.37 |
| 45.709 | 0.37 |
| 52.481 | 0.39 |
| 60.256 | 0.43 |
| 69.183 | 0.50 |
| 79.433 | 0.58 |
| 91.201 | 0.50 |
| 104.713 | |
| 120.226 | 0.76 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 120.226 | 0.8 |
| 138.038 | 0.8 |
| 158.489 | 0.8 |
| 181.970 | 0.8 |
| 208.930 | 0.76 |
| 239.883 | 0.8 |
| 275.423 | 1.3 |
| 316.228 | 2.1 |
| 363.078 | 3.2 |
| 416.869 | 4.69 |
| 478.630 | 6.36 |
| 549.541 | 8.04 |
| 630.957 | 9.4 |
| 724.436 | 10.17 |
| 831.764 | 10.1 |
| 954.993 | 10.1 |
| | |

1096.478

1258.925

9.22

7.62

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 5.64 |
| 1445.440 | 3.66 |
| 1659.587 | 1.51 |
| 1905.461 | 0.22 |
| 2187.762 | 0 |
| 2511.886 | 0.00 |
| 2884.032 | 0.00 |
| 3311.311 | 0.00 |
| 3801.894 | 0.00 |
| 4365.158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | |
| | |
| | |

Operator notes:





Sample Name:

SOP Name:

Tuesday, 17 January 2012 4:55:57 PM

Sample Source & type:

Measured by:

Analysed:

abbottsj

Tuesday, 17 January 2012 4:55:59 PM

Sample bulk lot ref:

Result Source: Measurement

Particle Name:

Accessory Name: Hydro 2000G (A)

Sensitivity: Normal

Silica 1.45 Particle RI:

Absorption:

General purpose Size range:

1.450

0.1

Obscuration:

Dispersant Name:

to 2000.000 um

5.58 %

Water

Dispersant RI:

Weighted Residual:

Analysis model:

Result Emulation:

1.330

6.472

0.020

Off

Concentration:

Span:

Uniformity:

Result units:

0.1396 %Vol 1.563

Volume

Specific Surface Area: 0.0337

Surface Weighted Mean D[3,2]: 177.992 um

Vol. Weighted Mean D[4,3]:

770.974


d(0.1): 306.801 um

d(0.5):

695.114

um

d(0.9): 1393.505 um



| Size (µm) 0.010 0.011 0.013 0.015 0.017 0.020 0.023 0.026 0.030 0.035 0.040 0.046 0.052 0.060 0.069 0.079 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | Size (µm) 0.105 0.120 0.138 0.158 0.182 0.209 0.240 0.275 0.316 0.363 0.417 0.479 0.550 0.631 0.724 0.832 0.955 | 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.0 | Size (µm) 1.096 1.259 1.445 1.660 1.905 2.188 2.512 2.884 3.311 3.802 4.365 5.012 5.754 6.607 7.586 8.710 10.000 | 0.00 0.00 0.00 0.00 0.00 0.00 0.05 0.08 0.10 0.12 0.14 0.17 0.19 0.20 0.22 | Size (µm) 11.482 13.183 15.136 17.378 19.953 22.909 26.303 30.200 34.674 39.811 45.709 52.481 60.256 69.183 79.433 91.201 104.713 | 0.24 0.23 0.21 0.18 0.16 0.14 0.13 0.14 0.16 0.20 0.23 0.26 0.25 0.21 | Size (µm) 120.226 138.038 158.489 181.970 208.930 239.883 275.423 316.228 363.078 416.869 478.630 549.541 630.957 724.436 831.764 954.993 | 0.29 0.90 1.81 2.96 4.29 5.63 6.86 7.83 8.48 8.79 8.79 8.52 8.03 | 12 14 16 15 22 28 33 38 43 50 56 66 75 87 | ize (µn 1258.922 1445.44 1659.58 1905.46 1905.46 12187.76 2511.88 2884.03 33311.31 33311.31 33311.31 336606.93 7585.77 |
|---|--|---|---|--|--|---|--|---|--|--|---|
| 0.105 | 0.00 | 1.096 | 0.00 | 11.482 | 0.24 | 120.226 | 0.00 | 1258.925 | 7.30 | | |

Operator notes:

4.89 2.87 0.70 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00





Sample Name:

SOP Name:

D6 - Average

Tuesday, 17 January 2012 5:16:05 PM

Sample Source & type:

Measured by: Analysed:

abbottsj Tuesday, 17 January 2012 5:16:07 PM

Sample bulk lot ref:

Result Source:

Averaged

Particle Name: Silica 1.45

Accessory Name: Hydro 2000G (A)

Sensitivity: Normal

Particle RI:

Absorption:

General purpose

1.450

Obscuration:

Dispersant Name:

0.1

to 2000.000 um 12.39 %

Water

0.0688

0.297

Dispersant RI:

Weighted Residual:

Analysis model:

Result Emulation:

1.330

2.998

Size range:

Off

Concentration:

Span:

Uniformity:

0.020

Result units: Volume

Specific Surface Area:

%Vol

3.000

Surface Weighted Mean D[3,2]: 20.195

um

Vol. Weighted Mean D[4,3]:

545.749

d(0.1): 9.451


um

d(0.5):

429.527

um

d(0.9): 1298.067 um



| Size (µm) | Volume In % | Size (µm) | Volume In % | |
|-----------|-------------|-----------|-------------|--|
| 0.010 | 0.00 | 0.105 | 0.00 | |
| 0.011 | 0.00 | 0.120 | 0.00 | |
| 0.013 | | 0.138 | | |
| 0.015 | 0.00 | 0.158 | 0.00 | |
| 0.017 | 0.00 | 0.182 | 0.00 | |
| 0.020 | 0.00 | 0.209 | 0.00 | |
| 0.023 | 0.00 | 0.240 | 0.00 | |
| 0.026 | 0.00 | 0.275 | 0.00 | |
| 0.030 | 0.00 | 0.316 | 0.00 | |
| 0.035 | 0.00 | 0.363 | 0.01 | |
| 0.040 | 0.00 | 0.417 | 0.08 | |
| 0.046 | 0.00 | 0.479 | 0.13 | |
| 0.040 | 0.00 | 0.479 | 0.16 | |
| | 0.00 | | 0.18 | |
| 0.060 | 0.00 | 0.631 | 0.19 | |
| 0.069 | 0.00 | 0.724 | 0.18 | |
| 0.079 | 0.00 | 0.832 | 0.16 | |
| 0.091 | 0.00 | 0.955 | 0.14 | |
| 0.405 | | 4 000 | | |

| o: () | |
|-----------|-------------|
| Size (µm) | Volume In % |
| 1.096 | 0.12 |
| 1.259 | 0.12 |
| 1.445 | |
| 1.660 | 0.12 |
| 1.905 | 0.17 |
| 2,188 | 0.22 |
| 2,512 | 0.28 |
| 2.884 | 0.37 |
| 3.311 | 0.46 |
| 3.802 | 0.57 |
| | 0.68 |
| 4.365 | 0.79 |
| 5.012 | 0.90 |
| 5.754 | 1.00 |
| 6.607 | 1.09 |
| 7.586 | 1.17 |
| 8.710 | 1.17 |
| 10.000 | |
| 11.482 | 1.26 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 11.482 | 1.27 |
| 13.183 | 1.26 |
| 15.136 | 1.23 |
| 17.378 | 1.19 |
| 19.953 | |
| 22.909 | 1.14 |
| 26.303 | 1.09 |
| 30,200 | 1.03 |
| 34.674 | 0.97 |
| 39.811 | 0.91 |
| 45.709 | 0.84 |
| 52.481 | 0.77 |
| 60.256 | 0.69 |
| 69.183 | 0.61 |
| | 0.55 |
| 79.433 | 0.52 |
| 91.201 | 0.54 |
| 104.713 | 0.64 |
| 120.226 | 0.01 |

| Size (µm) | Volume In % |
|-----------|-------------|
| 120.226 | 0.00 |
| 138.038 | 0.83 |
| 158.489 | 1.11 |
| 181.970 | 1.48 |
| 208.930 | 1.93 |
| 239,883 | 2.42 |
| 275,423 | 2.92 |
| 316.228 | 3.39 |
| 363.078 | 3.80 |
| 416.869 | 4.14 |
| 478.630 | 4.40 |
| 549.541 | 4.62 |
| | 4.80 |
| 630.957 | 4.99 |
| 724.436 | 5.18 |
| 831.764 | 5.33 |
| | |

1096.478

1258.925

5.36

5.17

| Size (µm) | Volume In % |
|-----------|-------------|
| 1258.925 | 4.63 |
| 1445.440 | 3.71 |
| 1659.587 | 2.21 |
| 1905.461 | 0.54 |
| 2187.762 | 0.00 |
| 2511.886 | 0.00 |
| 2884.032 | 5.55 |
| 3311.311 | 0.00 |
| 3801.894 | 0.00 |
| 4365.158 | 0.00 |
| 5011.872 | 0.00 |
| 5754.399 | 0.00 |
| 6606.934 | 0.00 |
| 7585.776 | 0.00 |
| 8709.636 | 0.00 |
| 10000.000 | 0.00 |
| 10000.000 | |
| | |
| | |

Operator notes:



B2. Chemical Analysis of Sediments Laboratory Results

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

INTERPRETIVE QUALITY CONTROL REPORT

Work Order : **EM1114568** Page : 1 of 8

Amendment : 1

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : DR GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

Telephone : +61 08 8424 3800 Telephone : +61-3-8549 9608 Facsimile : +61 08 8424 3810 Facsimile : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

Site : ----

C-O-C number : ---- Date Samples Received : 22-DEC-2011
Sampler : GB Issue Date : 28-FEB-2012

Order number :----

Quote number : FN/003/10 No. of samples received : 14

Quote number : FN/003/10 No. of samples analysed : 14

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for release.

This Interpretive Quality Control Report contains the following information:

- Analysis Holding Time Compliance
- Quality Control Parameter Frequency Compliance
- Brief Method Summaries
- Summary of Outliers

Environmental Division Melbourne
Part of the ALS Laboratory Group

4 Westall Rd Springvale VIC Australia 3171

Tel. +61-3-8549 9600 Fax. +61-3-8549 9601 www.alsglobal.com

A Campbell Brothers Limited Company

Page : 2 of 8

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Analysis Holding Time Compliance

The following report summarises extraction / preparation and analysis times and compares with recommended holding times. Dates reported represent first date of extraction or analysis and precludes subsequent dilutions and reruns. Information is also provided re the sample container (preservative) from which the analysis aliquot was taken. Elapsed period to analysis represents number of days from sampling where no extraction / digestion is involved or period from extraction / digestion where this is present. For composite samples, sampling date is assumed to be that of the oldest sample contributing to the composite. Sample date for laboratory produced leachates is assumed as the completion date of the leaching process. Outliers for holding time are based on USEPA SW 846, APHA, AS and NEPM (1999). A listing of breaches is provided in the Summary of Outliers.

Holding times for leachate methods (excluding elutriates) vary according to the analytes being determined on the resulting solution. For non-volatile analytes, the holding time compliance assessment compares the leach date with the shortest analyte holding time for the equivalent soil method. These soil holding times are: Organics (14 days); Mercury (28 days) & other metals (180 days). A recorded breach therefore does not guarantee a breach for all non-volatile parameters.

| Matrix: SOIL | | | | | Evaluation | : x = Holding time | breach ; ✓ = Withir | n holding time |
|---------------------------------|---------------|-------------|----------------|------------------------|------------|--------------------|---------------------|----------------|
| Method | | Sample Date | Ex | traction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EA055: Moisture Content | | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | | | | 31-DEC-2011 | 02-JAN-2012 | ✓ |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | B6, | | | | | | | |
| C6, | D6 | | | | | | | |
| EG005T: Total Metals by ICP-AES | | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 05-JAN-2012 | 16-JUN-2012 | ✓ | 06-JAN-2012 | 16-JUN-2012 | ✓ |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | B6, | | | | | | | |
| C6, | D6 | | | | | | | |
| EG020-SD: Total Metals in Sedim | ents by ICPMS | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 05-JAN-2012 | 16-JUN-2012 | ✓ | 05-JAN-2012 | 16-JUN-2012 | ✓ |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | В6, | | | | | | | |
| C6, | D6 | | | | | | | |
| EG035T: Total Recoverable Merc | cury by FIMS | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 05-JAN-2012 | 16-JAN-2012 | ✓ | 05-JAN-2012 | 16-JAN-2012 | 1 |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | В6, | | | | | | | |
| C6, | D6 | | | | | | | |

Page : 3 of 8

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Matrix: SOIL | | | | | Evaluation: | × = Holding time | breach ; ✓ = Withir | holding time. |
|---------------------------------|--------------------------------|-------------|--------------------------|--------------------|-------------|------------------|---------------------|---------------|
| Method | | Sample Date | Extraction / Preparation | | | Analysis | | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EP068A: Organochlorine Pesti | icides (OC) | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 29-DEC-2011 | 02-JAN-2012 | ✓ | 30-DEC-2011 | 07-FEB-2012 | ✓ |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | B6, | | | | | | | |
| C6, | D6 | | | | | | | |
| EP068B: Organophosphorus P | | | | | | | I | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 29-DEC-2011 | 02-JAN-2012 | ✓ | 30-DEC-2011 | 07-FEB-2012 | ✓ |
| C1, | D1, | | | | | | | |
| A4, C4, | B4, D4, | | | | | | | |
| A5, | D4, C5, | | | | | | | |
| D5, | B6, | | | | | | | |
| C6, | D6 | | | | | | | |
| | Hydrocarbons - NEPM 2010 Draft | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | I | | |
| A1, | B1, | 19-DEC-2011 | 29-DEC-2011 | 02-JAN-2012 | 1 | 31-DEC-2011 | 07-FEB-2012 | ✓ |
| C1, | D1, | 19-020-2011 | 29-DEC-2011 | 02-3AN-2012 | • | 31-DEC-2011 | 07-1 LD-2012 | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | B6, | | | | | | | |
| C6, | D6 | | | | | | | |
| EP080-SD / EP071-SD: Total Pe | etroleum Hydrocarbons | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 29-DEC-2011 | 02-JAN-2012 | ✓ | 30-DEC-2011 | 02-JAN-2012 | ✓ |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | B6, | | | | | | | |
| C6, | D6 | | | | | | | |
| Soil Glass Jar - Unpreserved | B1, | 10 850 0011 | 00 050 0044 | 00 IAN 0040 | | 04 850 0044 | 07 FED 2042 | |
| A1, C1, | ы, D1, | 19-DEC-2011 | 29-DEC-2011 | 02-JAN-2012 | ✓ | 31-DEC-2011 | 07-FEB-2012 | ✓ |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | B6, | | | | | | | |
| C6, | D6 | | | | | | | |
| <u> </u> | - | | | | | | | |

Page : 4 of 8

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Matrix: SOIL | | | | | Evaluation | × = Holding time | breach ; ✓ = Withir | n holding time |
|----------------------------------|----------------------|-------------|----------------|------------------------|------------|------------------|---------------------|----------------|
| Method | | | Ex | traction / Preparation | | Analysis | | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EP080-SD / EP071-SD: Total Recov | verable Hydrocarbons | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 29-DEC-2011 | 02-JAN-2012 | ✓ | 30-DEC-2011 | 02-JAN-2012 | 1 |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | B6, | | | | | | | |
| C6, | D6 | | | | | | | |
| EP080-SD: BTEXN | | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 29-DEC-2011 | 02-JAN-2012 | 1 | 30-DEC-2011 | 02-JAN-2012 | 1 |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | В6, | | | | | | | |
| C6, | D6 | | | | | | | |
| EP090: Organotin Compounds | | | | | | | | |
| Soil Glass Jar - Unpreserved | | | | | | | | |
| A1, | B1, | 19-DEC-2011 | 28-DEC-2011 | 02-JAN-2012 | ✓ | 03-JAN-2012 | 06-FEB-2012 | ✓ |
| C1, | D1, | | | | | | | |
| A4, | B4, | | | | | | | |
| C4, | D4, | | | | | | | |
| A5, | C5, | | | | | | | |
| D5, | В6, | | | | | | | |
| C6, | D6 | | | | | | | |

Page : 5 of 8

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Quality Control Parameter Frequency Compliance

The following report summarises the frequency of laboratory QC samples analysed within the analytical lot(s) in which the submitted sample(s) was(where) processed. Actual rate should be greater than or equal to the expected rate. A listing of breaches is provided in the Summary of Outliers.

Matrix: **SOIL** Evaluation: **×** = Quality Control frequency not within specification; ✓ = Quality Control frequency within specification.

| Quality Control Sample Type | | | ount | | Rate (%) | | Quality Control Specification |
|------------------------------------|-----------|---|---------|--------|----------|------------|--|
| Analytical Methods Method | | | Regular | Actual | Expected | Evaluation | |
| Laboratory Duplicates (DUP) | | | | | | | |
| Moisture Content | EA055-103 | 2 | 20 | 10.0 | 10.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| Organotin Analysis | EP090 | 2 | 9 | 22.2 | 10.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| Pesticides by GCMS | EP068 | 2 | 17 | 11.8 | 10.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| otal Mercury by FIMS (Low Level) | EG035T-LL | 2 | 14 | 14.3 | 10.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| Total Metals by ICP-AES | EG005T | 2 | 14 | 14.3 | 10.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| Total Metals in Sediments by ICPMS | EG020-SD | 2 | 14 | 14.3 | 10.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| FPH - Semivolatile Fraction | EP071-SD | 2 | 14 | 14.3 | 10.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| PH Volatiles/BTEX in Sediments | EP080-SD | 2 | 14 | 14.3 | 10.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| aboratory Control Samples (LCS) | | | | | | | |
| Organotin Analysis | EP090 | 1 | 9 | 11.1 | 5.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| Pesticides by GCMS | EP068 | 1 | 17 | 5.9 | 5.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| otal Mercury by FIMS (Low Level) | EG035T-LL | 1 | 14 | 7.1 | 5.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| otal Metals by ICP-AES | EG005T | 1 | 14 | 7.1 | 5.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| otal Metals in Sediments by ICPMS | EG020-SD | 1 | 14 | 7.1 | 5.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| PH - Semivolatile Fraction | EP071-SD | 1 | 14 | 7.1 | 5.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| PH Volatiles/BTEX in Sediments | EP080-SD | 1 | 14 | 7.1 | 5.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| Method Blanks (MB) | | | | | | | |
| Organotin Analysis | EP090 | 1 | 9 | 11.1 | 5.0 | ✓ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| Pesticides by GCMS | EP068 | 1 | 17 | 5.9 | 5.0 | √ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| otal Mercury by FIMS (Low Level) | EG035T-LL | 1 | 14 | 7.1 | 5.0 | 1 | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| otal Metals by ICP-AES | EG005T | 1 | 14 | 7.1 | 5.0 | 1 | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| otal Metals in Sediments by ICPMS | EG020-SD | 1 | 14 | 7.1 | 5.0 | √ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| PH - Semivolatile Fraction | EP071-SD | 1 | 14 | 7.1 | 5.0 | √ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| PH Volatiles/BTEX in Sediments | EP080-SD | 1 | 14 | 7.1 | 5.0 | √ | NEPM 1999 Schedule B(3) and ALS QCS3 requirement |
| Matrix Spikes (MS) | | | | | | | |
| Organotin Analysis | EP090 | 1 | 9 | 11.1 | 5.0 | 1 | ALS QCS3 requirement |
| Pesticides by GCMS | EP068 | 1 | 17 | 5.9 | 5.0 | √ | ALS QCS3 requirement |
| otal Mercury by FIMS (Low Level) | EG035T-LL | 1 | 14 | 7.1 | 5.0 | √ | ALS QCS3 requirement |
| otal Metals by ICP-AES | EG005T | 1 | 14 | 7.1 | 5.0 | √ | ALS QCS3 requirement |
| otal Metals in Sediments by ICPMS | EG020-SD | 1 | 14 | 7.1 | 5.0 | 1 | ALS QCS3 requirement |
| PH - Semivolatile Fraction | EP071-SD | 1 | 14 | 7.1 | 5.0 | √ | ALS QCS3 requirement |
| PH Volatiles/BTEX in Sediments | EP080-SD | 1 | 14 | 7.1 | 5.0 | 1 | ALS QCS3 requirement |

Page : 6 of 8

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Brief Method Summaries

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the US EPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request. The following report provides brief descriptions of the analytical procedures employed for results reported in the Certificate of Analysis. Sources from which ALS methods have been developed are provided within the Method Descriptions.

| Analytical Methods | Method | Matrix | Method Descriptions |
|--|-----------|--------|--|
| Moisture Content | EA055-103 | SOIL | A gravimetric procedure based on weight loss over a 12 hour drying period at 103-105 degrees C. This method is |
| | | | compliant with NEPM (2010 Draft) Schedule B(3) Section 7.1 and Table 1 (14 day holding time). |
| Total Metals by ICP-AES | EG005T | SOIL | (APHA 21st ed., 3120; USEPA SW 846 - 6010) (ICPAES) Metals are determined following an appropriate acid |
| | | | digestion of the soil. The ICPAES technique ionises samples in a plasma, emitting a characteristic spectrum |
| | | | based on metals present. Intensities at selected wavelengths are compared against those of matrix matched |
| | | | standards. This method is compliant with NEPM (1999) Schedule B(3) |
| Total Metals in Sediments by ICPMS | EG020-SD | SOIL | (APHA 21st ed., 3125; USEPA SW846 - 6020, ALS QWI-EN/EG020): The ICPMS technique utilizes a highly |
| | | | efficient argon plasma to ionize selected elements. Ions are then passed into a high vacuum mass spectrometer, |
| | | | which separates the analytes based on their distinct mass to charge ratios prior to their measurement by a discrete |
| Total Marsum, by FIMC (Lavel aval) | EG035T-LL | COII | dynode ion detector. Analyte list and LORs per NODG. |
| Total Mercury by FIMS (Low Level) | EG0331-LL | SOIL | AS 3550, APHA 21st ed., 3112 Hg - B (Flow-injection (SnCl2)(Cold Vapour generation) AAS) FIM-AAS is an automated flameless atomic absorption technique. Mercury in solids are determined following an appropriate acid |
| | | | digestion. Ionic mercury is reduced online to atomic mercury vapour by SnCl2 which is then purged into a heated |
| | | | quartz cell. Quantification is by comparing absorbance against a calibration curve. This method is compliant with |
| | | | NEPM (1999) Schedule B(3) |
| Pesticides by GCMS | EP068 | SOIL | (USEPA SW 846 - 8270B) Extracts are analysed by Capillary GC/MS and quantification is by comparison against |
| | | | an established 5 point calibration curve. This technique is compliant with NEPM (1999) Schedule B(3) (Method |
| | | | 504,505) |
| TPH - Semivolatile Fraction | EP071-SD | SOIL | (USEPA SW 846 - 8270B) Extracts are analysed by Capillary GC/MS and quantification is by comparison against |
| | | | an established 5 point calibration curve. This method is compliant with NEPM (1999) Schedule B(3) (Method 504) |
| TPH Volatiles/BTEX in Sediments | EP080-SD | SOIL | (USEPA SW 846 - 8260B) Extracts are analysed by Purge and Trap, Capillary GC/MS. Quantification is by |
| | | | comparison against an established 5 point calibration curve. This method is compliant with NEPM (1999) |
| | | | Schedule B(3) (Method 501) |
| Organotin Analysis | EP090 | SOIL | (USEPA SW 846 - 8270D) Prepared sample extracts are analysed by GC/MS coupled with high volume injection, |
| | | | and quanitified against an established calibration curve. |
| Preparation Methods | Method | Matrix | Method Descriptions |
| Hot Block Digest for metals in soils | EN69 | SOIL | USEPA 200.2 Mod. Hot Block Acid Digestion 1.0g of sample is heated with Nitric and Hydrochloric acids, then |
| sediments and sludges | | | cooled. Peroxide is added and samples heated and cooled again before being filtered and bulked to volume for |
| | | | analysis. Digest is appropriate for determination of selected metals in sludge, sediments, and soils. This method is |
| | | | compliant with NEPM (1999) Schedule B(3) (Method 202) |
| Methanolic Extraction of Soils for Purge | * ORG16 | SOIL | (USEPA SW 846 - 5030A) 5g of solid is shaken with surrogate and 10mL methanol prior to analysis by Purge and |
| and Trap | 000474 | 0011 | Trap - GC/MS. |
| Tumbler Extraction of Solids (Option A - | ORG17A | SOIL | In-house, Mechanical agitation (tumbler). 20g of sample, Na2SO4 and surrogate are extracted with 150mL 1:1 |
| Concentrating) | | | DCM/Acetone by end over end tumble. The solvent is decanted, dehydrated and concentrated (by KD) to the |
| Tumbler Extraction of Solids for LVI | ORG17D | SOIL | desired volume for analysis. In house: 10g of sample, Na2SO4 and surrogate are extracted with 50mL 1:1 DCM/Acetone by end over end |
| (Non-concentrating) | ONGITU | SUIL | tumbling. An aliquot is concentrated by nitrogen blowdown to a reduced volume for analysis if required. |
| (14011 0011001titatility) | | | taribing. 7 at anquet to concentrated by fine open blowdown to a reduced volume for analysis it required. |

Page : 7 of 8

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Preparation Methods | Method | Matrix | Method Descriptions |
|------------------------------|--------|--------|--|
| Organotin Sample Preparation | ORG35 | SOIL | In house. 20g sample is spiked with surrogate and leached in a methanol:acetic acid:UHP water mix and vacuum filtered. Reagents and solvents are added to the sample and the mixture tumbled. The butyltin compounds are simultaneously derivatised and extracted. The extract is further extracted with petroleum ether. The resultant extracts are combined and concentrated for analysis. |

Page : 8 of 8

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Summary of Outliers

Outliers: Quality Control Samples

The following report highlights outliers flagged in the Quality Control (QC) Report. Surrogate recovery limits are static and based on USEPA SW846 or ALS-QWI/EN/38 (in the absence of specific USEPA limits). This report displays QC Outliers (breaches) only.

Duplicates, Method Blanks, Laboratory Control Samples and Matrix Spikes

Matrix: SOIL

| Compound Group Name | Laboratory Sample ID | Client Sample ID | Analyte | CAS Number | Data | Limits | Comment |
|---|----------------------|------------------|------------------|------------|-------|-----------|---|
| Laboratory Control Spike (LCS) Recoveries | | | | | | | |
| EP068B: Organophosphorus Pesticides (OP) | 2492376-002 | | Prothiofos | 34643-46-4 | 112 % | 65.4-111% | Recovery greater than upper control limit |
| Matrix Spike (MS) Recoveries | | | | | | | |
| EP080-SD / EP071-SD: Total Petroleum Hydrocarbons | EM1114568-001 | A1 | C6 - C9 Fraction | | 130 % | 70-130% | Recovery greater than upper data quality |
| | | | | | | | objective |

- For all matrices, no Method Blank value outliers occur.
- For all matrices, no Duplicate outliers occur.

Regular Sample Surrogates

Sub-Matrix: SEDIMENT

| Compound Group Name | Laboratory Sample ID | Client Sample ID | Analyte | CAS Number | Data | Limits | Comment |
|-----------------------------|----------------------|------------------|--------------|------------|-------|----------|--|
| Samples Submitted | | | | | | | |
| EP090S: Organotin Surrogate | EM1114568-013 | C6 | Tripropyltin | | 146 % | 35-130 % | Recovery greater than upper data quality |
| | | | | | | | objective |

Outliers: Analysis Holding Time Compliance

This report displays Holding Time breaches only. Only the respective Extraction / Preparation and/or Analysis component is/are displayed.

No Analysis Holding Time Outliers exist.

Outliers : Frequency of Quality Control Samples

The following report highlights breaches in the Frequency of Quality Control Samples.

• No Quality Control Sample Frequency Outliers exist.

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

QUALITY CONTROL REPORT

Work Order : **EM1114568** Page : 1 of 11

Amendment : 1

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : DR GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

 Telephone
 : +61 08 8424 3800
 Telephone
 : +61-3-8549 9608

 Facsimile
 : +61 08 8424 3810
 Facsimile
 : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

Site : ----

 C-O-C number
 : -- Date Samples Received
 : 22-DEC-2011

 Sampler
 : GB
 Issue Date
 : 28-FEB-2012

No. of samples received : 14

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for

No. of samples analysed

This Quality Control Report contains the following information:

: ----

: EN/003/10

- Laboratory Duplicate (DUP) Report; Relative Percentage Difference (RPD) and Acceptance Limits
- Method Blank (MB) and Laboratory Control Spike (LCS) Report; Recovery and Acceptance Limits
- Matrix Spike (MS) Report; Recovery and Acceptance Limits



Order number

Quote number

release.

NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

: 14

| Signatories | Position | Accreditation Category |
|------------------------------|---|--------------------------------------|
| Di-An Dao | Soniar Organia Chamiet | Sydney Inorganics |
| Matt Frost Phalak Inthaksone | Senior Organic Chemist Laboratory Manager - Organics | Brisbane Organics Sydney Organics |
| Wisam Marassa | Inorganics Coordinator | Sydney Inorganics |

Page : 2 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

Key: Anonymous = Refers to samples which are not specifically part of this work order but formed part of the QC process lot

CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

RPD = Relative Percentage Difference

= Indicates failed QC

Page : 3 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Laboratory Duplicate (DUP) Report

The quality control term Laboratory Duplicate refers to a randomly selected intralaboratory split. Laboratory duplicates provide information regarding method precision and sample heterogeneity. The permitted ranges for the Relative Percent Deviation (RPD) of Laboratory Duplicates are specified in ALS Method QWI-EN/38 and are dependent on the magnitude of results in comparison to the level of reporting: Result < 10 times LOR:-No Limit; Result between 10 and 20 times LOR:-0% - 50%; Result > 20 times LOR:-0% - 20%.

| Laboratory sample ID | | | | Laboratory Duplicate (DUP) Report | | | | | | | |
|----------------------|-----------------------------|---|------------|-----------------------------------|-------|-----------------|------------------|---------|---------------------|--|--|
| Laboratory Sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) | | |
| EA055: Moisture Cor | ntent (QC Lot: 2111940) | | | | | | | | | | |
| EM1114568-003 | C1 | EA055-103: Moisture Content (dried @ 103°C) | | 1.0 | % | 35.6 | 35.6 | 0.0 | 0% - 20% | | |
| EM1114568-014 | D6 | EA055-103: Moisture Content (dried @ 103°C) | | 1.0 | % | 31.3 | 31.8 | 1.7 | 0% - 20% | | |
| EG005T: Total Metals | s by ICP-AES (QC Lot: 211 | 5272) | | | | | | | | | |
| EM1114568-001 | A1 | EG005T: Beryllium | 7440-41-7 | 1 | mg/kg | <1 | <1 | 0.0 | No Limit | | |
| | | EG005T: Barium | 7440-39-3 | 10 | mg/kg | <10 | <10 | 0.0 | No Limit | | |
| | | EG005T: Iron | 7439-89-6 | 50 | mg/kg | 2460 | 2520 | 2.4 | 0% - 20% | | |
| EM1114568-011 | D5 | EG005T: Beryllium | 7440-41-7 | 1 | mg/kg | <1 | <1 | 0.0 | No Limit | | |
| | | EG005T: Barium | 7440-39-3 | 10 | mg/kg | <10 | <10 | 0.0 | No Limit | | |
| | | EG005T: Iron | 7439-89-6 | 50 | mg/kg | 6540 | 6630 | 1.3 | 0% - 20% | | |
| EG020-SD: Total Met | tals in Sediments by ICPMS | (QC Lot: 2115274) | | | | | | | | | |
| EM1114568-001 | A1 | EG020-SD: Cadmium | 7440-43-9 | 0.1 | mg/kg | <0.1 | <0.1 | 0.0 | No Limit | | |
| | | EG020-SD: Cobalt | 7440-48-4 | 0.5 | mg/kg | <0.5 | 0.5 | 0.0 | No Limit | | |
| | | EG020-SD: Chromium | 7440-47-3 | 1.0 | mg/kg | 5.5 | 6.8 | 21.4 | No Limit | | |
| | | EG020-SD: Copper | 7440-50-8 | 1.0 | mg/kg | 1.3 | 1.3 | 0.0 | No Limit | | |
| | | EG020-SD: Lead | 7439-92-1 | 1.0 | mg/kg | 1.1 | 1.3 | 16.2 | No Limit | | |
| | | EG020-SD: Nickel | 7440-02-0 | 1.0 | mg/kg | <1.0 | <1.0 | 0.0 | No Limit | | |
| | | EG020-SD: Zinc | 7440-66-6 | 1.0 | mg/kg | 1.4 | 1.6 | 10.4 | No Limit | | |
| | | EG020-SD: Arsenic | 7440-38-2 | 1.00 | mg/kg | 2.54 | 2.25 | 11.9 | No Limit | | |
| | | EG020-SD: Manganese | 7439-96-5 | 10 | mg/kg | 10 | 11 | 0.0 | No Limit | | |
| | | EG020-SD: Vanadium | 7440-62-2 | 2.0 | mg/kg | <2.0 | <2.0 | 0.0 | No Limit | | |
| EM1114568-011 | D5 | EG020-SD: Cadmium | 7440-43-9 | 0.1 | mg/kg | <0.1 | <0.1 | 0.0 | No Limit | | |
| | | EG020-SD: Cobalt | 7440-48-4 | 0.5 | mg/kg | 0.5 | <0.5 | 0.0 | No Limit | | |
| | | EG020-SD: Chromium | 7440-47-3 | 1.0 | mg/kg | 13.5 | 12.4 | 8.6 | 0% - 50% | | |
| | | EG020-SD: Copper | 7440-50-8 | 1.0 | mg/kg | <1.0 | <1.0 | 0.0 | No Limit | | |
| | | EG020-SD: Lead | 7439-92-1 | 1.0 | mg/kg | 1.8 | 1.7 | 9.1 | No Limit | | |
| | | EG020-SD: Nickel | 7440-02-0 | 1.0 | mg/kg | 1.3 | 1.1 | 18.0 | No Limit | | |
| | | EG020-SD: Zinc | 7440-66-6 | 1.0 | mg/kg | 2.1 | 1.9 | 12.7 | No Limit | | |
| | | EG020-SD: Arsenic | 7440-38-2 | 1.00 | mg/kg | 8.99 | 8.31 | 7.9 | No Limit | | |
| | | EG020-SD: Manganese | 7439-96-5 | 10 | mg/kg | 14 | 12 | 13.0 | No Limit | | |
| | | EG020-SD: Vanadium | 7440-62-2 | 2.0 | mg/kg | 19.1 | 17.6 | 8.3 | No Limit | | |
| EG035T: Total Reco | verable Mercury by FIMS (| QC Lot: 2115273) | | | | | | | | | |
| EM1114568-001 | A1 | EG035T-LL: Mercury | 7439-97-6 | 0.01 | mg/kg | <0.01 | <0.01 | 0.0 | No Limit | | |
| EM1114568-011 | D5 | EG035T-LL: Mercury | 7439-97-6 | 0.01 | mg/kg | <0.01 | <0.01 | 0.0 | No Limit | | |
| EP068A: Organochio | orine Pesticides (OC) (QC L | ot: 2106215) | | | | | | | • | | |
| EM1114568-001 | A1 | EP068: alpha-BHC | 319-84-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |

Page : 4 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: SOIL | | | | | | Laboratory Duplicate (DUP) Report | | | | | |
|----------------------|-----------------------------|--------------------------------|------------|------|-------|-----------------------------------|------------------|---------|---------------------|--|--|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) | | |
| EP068A: Organochio | orine Pesticides (OC) (QC I | Lot: 2106215) - continued | | | | | | | | | |
| EM1114568-001 | A1 | EP068: Hexachlorobenzene (HCB) | 118-74-1 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: beta-BHC | 319-85-7 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: gamma-BHC | 58-89-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: delta-BHC | 319-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Heptachlor | 76-44-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Aldrin | 309-00-2 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Heptachlor epoxide | 1024-57-3 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: trans-Chlordane | 5103-74-2 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: alpha-Endosulfan | 959-98-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: cis-Chlordane | 5103-71-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Dieldrin | 60-57-1 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: 4.4`-DDE | 72-55-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Endrin | 72-20-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: beta-Endosulfan | 33213-65-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: 4.4`-DDD | 72-54-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Endrin aldehyde | 7421-93-4 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Endosulfan sulfate | 1031-07-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Endrin ketone | 53494-70-5 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: 4.4`-DDT | 50-29-3 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | | |
| | | EP068: Methoxychlor | 72-43-5 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | | |
| EM1114568-011 | D5 | EP068: alpha-BHC | 319-84-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Hexachlorobenzene (HCB) | 118-74-1 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: beta-BHC | 319-85-7 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: gamma-BHC | 58-89-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: delta-BHC | 319-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Heptachlor | 76-44-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Aldrin | 309-00-2 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Heptachlor epoxide | 1024-57-3 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: trans-Chlordane | 5103-74-2 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: alpha-Endosulfan | 959-98-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: cis-Chlordane | 5103-71-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Dieldrin | 60-57-1 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: 4.4`-DDE | 72-55-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Endrin | 72-20-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: beta-Endosulfan | 33213-65-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: 4.4`-DDD | 72-54-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Endrin aldehyde | 7421-93-4 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Endosulfan sulfate | 1031-07-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: Endrin ketone | 53494-70-5 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | | |
| | | EP068: 4.4`-DDT | 50-29-3 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | | |

Page : 5 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| ub-Matrix: SOIL | | | | Laboratory Duplicate (DUP) Report | | | | | | |
|---------------------|-------------------------|------------------------------|------------|-----------------------------------|-------|-----------------|------------------|---------|--------------------|--|
| aboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (% | |
| P068A: Organochi | orine Pesticides (OC) (| QC Lot: 2106215) - continued | | | | | | | | |
| M1114568-011 | D5 | EP068: Methoxychlor | 72-43-5 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | |
| 068B: Organopho | osphorus Pesticides (O | P) (QC Lot: 2106215) | | | | | | | | |
| M1114568-001 | A1 | EP068: Dichlorvos | 62-73-7 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Demeton-S-methyl | 919-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Dimethoate | 60-51-5 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Diazinon | 333-41-5 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Chlorpyrifos-methyl | 5598-13-0 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Malathion | 121-75-5 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Fenthion | 55-38-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Chlorpyrifos | 2921-88-2 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Pirimphos-ethyl | 23505-41-1 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Chlorfenvinphos | 470-90-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Bromophos-ethyl | 4824-78-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Fenamiphos | 22224-92-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Prothiofos | 34643-46-4 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Ethion | 563-12-2 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Carbophenothion | 786-19-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Azinphos Methyl | 86-50-0 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Monocrotophos | 6923-22-4 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | |
| | | EP068: Parathion-methyl | 298-00-0 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | |
| | | EP068: Parathion | 56-38-2 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | |
| И1114568-011 | D5 | EP068: Dichlorvos | 62-73-7 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Demeton-S-methyl | 919-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Dimethoate | 60-51-5 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Diazinon | 333-41-5 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Chlorpyrifos-methyl | 5598-13-0 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Malathion | 121-75-5 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Fenthion | 55-38-9 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Chlorpyrifos | 2921-88-2 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Pirimphos-ethyl | 23505-41-1 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Chlorfenvinphos | 470-90-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Bromophos-ethyl | 4824-78-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Fenamiphos | 22224-92-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Prothiofos | 34643-46-4 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Ethion | 563-12-2 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Carbophenothion | 786-19-6 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Azinphos Methyl | 86-50-0 | 0.05 | mg/kg | <0.05 | <0.05 | 0.0 | No Limit | |
| | | EP068: Monocrotophos | 6923-22-4 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | |
| | | EP068: Parathion-methyl | 298-00-0 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit | |

Page : 6 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: SOIL | | | | | | Laboratory L | Duplicate (DUP) Report | | |
|----------------------|------------------------|----------------------------------|------------|-----|---------|-----------------|------------------------|---------|---------------------|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) |
| EP068B: Organopho | osphorus Pesticides (O | P) (QC Lot: 2106215) - continued | | | | | | | |
| EM1114568-011 | D5 | EP068: Parathion | 56-38-2 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| EP080-SD / EP071-S | D: Total Petroleum Hyd | drocarbons (QC Lot: 2106035) | | | | | | | |
| EM1114568-002 B1 | B1 | EP071-SD: C10 - C14 Fraction | | 3 | mg/kg | <3 | <3 | 0.0 | No Limit |
| | | EP071-SD: C15 - C28 Fraction | | 3 | mg/kg | 22 | 22 | 0.0 | No Limit |
| | | EP071-SD: C29 - C36 Fraction | | 5 | mg/kg | 9 | 9 | 0.0 | No Limit |
| EM1114568-012 | B6 | EP071-SD: C10 - C14 Fraction | | 3 | mg/kg | <3 | <3 | 0.0 | No Limit |
| | | EP071-SD: C15 - C28 Fraction | | 3 | mg/kg | 31 | 30 | 0.0 | 0% - 50% |
| | | EP071-SD: C29 - C36 Fraction | | 5 | mg/kg | 12 | 9 | 24.3 | No Limit |
| EP080-SD / EP071-S | D: Total Petroleum Hyd | drocarbons (QC Lot: 2109665) | | | | | | | |
| EM1114568-001 | A1 | EP080-SD: C6 - C9 Fraction | | 3 | mg/kg | <3 | <3 | 0.0 | No Limit |
| EM1114568-011 | D5 | EP080-SD: C6 - C9 Fraction | | 3 | mg/kg | <3 | <3 | 0.0 | No Limit |
| EP080-SD: BTEXN | (QC Lot: 2109665) | | | | | | | | |
| EM1114568-001 | A1 | EP080-SD: Benzene | 71-43-2 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| | | EP080-SD: Toluene | 108-88-3 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| | | EP080-SD: Ethylbenzene | 100-41-4 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| | | EP080-SD: meta- & para-Xylene | 108-38-3 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| | | | 106-42-3 | | | | | | |
| | | EP080-SD: ortho-Xylene | 95-47-6 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| EM1114568-011 | D5 | EP080-SD: Benzene | 71-43-2 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| | | EP080-SD: Toluene | 108-88-3 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| | | EP080-SD: Ethylbenzene | 100-41-4 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| | | EP080-SD: meta- & para-Xylene | 108-38-3 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| | | | 106-42-3 | | | | | | |
| | | EP080-SD: ortho-Xylene | 95-47-6 | 0.2 | mg/kg | <0.2 | <0.2 | 0.0 | No Limit |
| EP090: Organotin C | ompounds (QC Lot: 21 | 05989) | | | | | | | |
| EM1114568-001 | A1 | EP090: Tributyltin | 56573-85-4 | 0.5 | μgSn/kg | <0.5 | <0.5 | 0.0 | No Limit |
| EM1114568-010 | C5 | EP090: Tributyltin | 56573-85-4 | 0.5 | μgSn/kg | <0.5 | <0.5 | 0.0 | No Limit |

Page : 7 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Method Blank (MB) and Laboratory Control Spike (LCS) Report

The quality control term Method / Laboratory Blank refers to an analyte free matrix to which all reagents are added in the same volumes or proportions as used in standard sample preparation. The purpose of this QC parameter is to monitor potential laboratory contamination. The quality control term Laboratory Control Sample (LCS) refers to a certified reference material, or a known interference free matrix spiked with target analytes. The purpose of this QC parameter is to monitor method precision and accuracy independent of sample matrix. Dynamic Recovery Limits are based on statistical evaluation of processed LCS.

| Sub-Matrix: SOIL | | | | Method Blank (MB) | Laboratory Control Spike (LCS) Report | | | | | |
|--|--------------------|----------|-------|-------------------|---------------------------------------|--------------------|----------|------------|--|--|
| | | | | Report | Spike | Spike Recovery (%) | Recovery | Limits (%) | | |
| Method: Compound | CAS Number | LOR Unit | | Result | Concentration | LCS | Low | High | | |
| EG005T: Total Metals by ICP-AES (QCLot: 211) | 5272) | | | | | | | | | |
| EG005T: Barium | 7440-39-3 | 10 | mg/kg | <10 | 137.41 mg/kg | 95.0 | 70 | 130 | | |
| EG005T: Beryllium | 7440-41-7 | 1 | mg/kg | <1 | 5.51 mg/kg | 94.5 | 70 | 130 | | |
| EG005T: Iron | 7439-89-6 | 50 | mg/kg | <50 | | | | | | |
| EG020-SD: Total Metals in Sediments by ICPMS | S (QCLot: 2115274) | | | | | | | | | |
| EG020-SD: Arsenic | 7440-38-2 | 1.0 | mg/kg | <1.00 | 13.1 mg/kg | 107 | 70 | 130 | | |
| EG020-SD: Cadmium | 7440-43-9 | 0.1 | mg/kg | <0.1 | 2.76 mg/kg | 98.1 | 70 | 130 | | |
| EG020-SD: Chromium | 7440-47-3 | 1.0 | mg/kg | <1.0 | 60.9 mg/kg | 102 | 70 | 130 | | |
| EG020-SD: Copper | 7440-50-8 | 1.0 | mg/kg | <1.0 | 54.7 mg/kg | 103 | 70 | 130 | | |
| EG020-SD: Cobalt | 7440-48-4 | 10 | mg/kg | <10.0 | 24.5 mg/kg | 106 | 70 | 130 | | |
| EG020-SD: Lead | 7439-92-1 | 1.0 | mg/kg | <1.0 | 54.8 mg/kg | 108 | 70 | 130 | | |
| EG020-SD: Manganese | 7439-96-5 | 10 | mg/kg | <10 | 136 mg/kg | 101 | 70 | 130 | | |
| EG020-SD: Nickel | 7440-02-0 | 1.0 | mg/kg | <1.0 | 55.2 mg/kg | 108 | 70 | 130 | | |
| EG020-SD: Vanadium | 7440-62-2 | 2 | mg/kg | <2.0 | 34 mg/kg | 90.5 | 70 | 130 | | |
| EG020-SD: Zinc | 7440-66-6 | 1.0 | mg/kg | <1.0 | 104 mg/kg | 99.7 | 70 | 130 | | |
| EG035T: Total Recoverable Mercury by FIMS | (QCLot: 2115273) | | | | | | | | | |
| EG035T-LL: Mercury | 7439-97-6 | 0.01 | mg/kg | <0.01 | 0.11 mg/kg | 84.7 | 74.2 | 126 | | |
| EP068A: Organochlorine Pesticides (OC) (QCL | _ot: 2106215) | | | | | | | | | |
| EP068: alpha-BHC | 319-84-6 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 100 | 60.8 | 116 | | |
| EP068: Hexachlorobenzene (HCB) | 118-74-1 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 90.0 | 59.4 | 115 | | |
| EP068: beta-BHC | 319-85-7 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 95.0 | 59.8 | 117 | | |
| EP068: gamma-BHC | 58-89-9 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 94.8 | 59.8 | 118 | | |
| EP068: delta-BHC | 319-86-8 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 82.5 | 65.8 | 114 | | |
| EP068: Heptachlor | 76-44-8 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 88.2 | 65.6 | 115 | | |
| EP068: Aldrin | 309-00-2 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 110 | 67 | 113 | | |
| EP068: Heptachlor epoxide | 1024-57-3 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 91.6 | 65.6 | 113 | | |
| EP068: trans-Chlordane | 5103-74-2 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 113 | 60.7 | 113 | | |
| EP068: alpha-Endosulfan | 959-98-8 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 110 | 65.8 | 116 | | |
| EP068: cis-Chlordane | 5103-71-9 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 74.6 | 57.3 | 120 | | |
| EP068: Dieldrin | 60-57-1 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 104 | 67.4 | 116 | | |
| EP068: 4.4`-DDE | 72-55-9 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 104 | 67.5 | 114 | | |
| EP068: Endrin | 72-20-8 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 94.3 | 63 | 121 | | |
| EP068: beta-Endosulfan | 33213-65-9 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 104 | 66.1 | 117 | | |
| EP068: 4.4`-DDD | 72-54-8 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 110 | 65.3 | 116 | | |
| EP068: Endrin aldehyde | 7421-93-4 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 85.0 | 57.3 | 115 | | |

Page : 8 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: SOIL | | | Method Blank (MB) | Laboratory Control Spike (LCS) Report | | | | | |
|---|------|-------|-------------------|---------------------------------------|--------------------|----------|------------|--|--|
| | | | Report | Spike | Spike Recovery (%) | Recovery | Limits (%) | | |
| Method: Compound CAS Number | LOR | Unit | Result | Concentration | LCS | Low | High | | |
| EP068A: Organochlorine Pesticides (OC) (QCLot: 2106215) - continued | | | | | | | | | |
| EP068: Endosulfan sulfate 1031-07-8 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 108 | 63.6 | 119 | | |
| EP068: 4.4`-DDT 50-29-3 | 0.2 | mg/kg | <0.2 | 0.5 mg/kg | 112 | 58.4 | 127 | | |
| EP068: Endrin ketone 53494-70-5 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 104 | 63.6 | 117 | | |
| EP068: Methoxychlor 72-43-5 | 0.2 | mg/kg | <0.2 | 0.5 mg/kg | 98.8 | 50.4 | 132 | | |
| EP068B: Organophosphorus Pesticides (OP) (QCLot: 2106215) | | | | | | | | | |
| EP068: Dichlorvos 62-73-7 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 103 | 25.5 | 124 | | |
| EP068: Demeton-S-methyl 919-86-8 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 103 | 10.1 | 159 | | |
| EP068: Monocrotophos 6923-22-4 | 0.2 | mg/kg | <0.2 | 0.5 mg/kg | 77.0 | 2.88 | 149 | | |
| EP068: Dimethoate 60-51-5 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 91.8 | 48.6 | 126 | | |
| EP068: Diazinon 333-41-5 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 96.8 | 64.9 | 111 | | |
| EP068: Chlorpyrifos-methyl 5598-13-0 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 110 | 65.1 | 111 | | |
| EP068: Parathion-methyl 298-00-0 | 0.2 | mg/kg | <0.2 | 0.5 mg/kg | 97.4 | 61.4 | 113 | | |
| EP068: Malathion 121-75-5 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 83.4 | 60.4 | 127 | | |
| EP068: Fenthion 55-38-9 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 102 | 64.7 | 110 | | |
| EP068: Chlorpyrifos 2921-88-2 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 104 | 64.2 | 111 | | |
| EP068: Parathion 56-38-2 | 0.2 | mg/kg | <0.2 | 0.5 mg/kg | 96.1 | 60 | 116 | | |
| EP068: Pirimphos-ethyl 23505-41-1 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 91.8 | 64.8 | 111 | | |
| EP068: Chlorfenvinphos 470-90-6 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 89.9 | 61.4 | 123 | | |
| EP068: Bromophos-ethyl 4824-78-6 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 81.7 | 64.3 | 114 | | |
| EP068: Fenamiphos 22224-92-6 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 109 | 45.5 | 128 | | |
| EP068: Prothiofos 34643-46-4 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | # 112 | 65.4 | 111 | | |
| EP068: Ethion 563-12-2 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 112 | 62 | 116 | | |
| EP068: Carbophenothion 786-19-6 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 112 | 59.5 | 119 | | |
| EP068: Azinphos Methyl 86-50-0 | 0.05 | mg/kg | <0.05 | 0.5 mg/kg | 78.3 | 29.8 | 137 | | |
| EP080-SD / EP071-SD: Total Petroleum Hydrocarbons (QCLot: 2106035) | | | | | | | | | |
| EP071-SD: C10 - C14 Fraction | 3 | mg/kg | <3 | 5 mg/kg | 87.0 | 75.2 | 116 | | |
| EP071-SD: C15 - C28 Fraction | 3 | mg/kg | <3 | 6.75 mg/kg | 99.2 | 75.3 | 113 | | |
| EP071-SD: C29 - C36 Fraction | 5 | mg/kg | <5 | 5 mg/kg | 79.0 | 72.6 | 117 | | |
| EP080-SD / EP071-SD: Total Petroleum Hydrocarbons (QCLot: 2109665) | | | | | | | | | |
| EP080-SD: C6 - C9 Fraction | 3 | mg/kg | <3 | 26 mg/kg | 114 | 61 | 133 | | |
| EP080-SD: BTEXN (QCLot: 2109665) | | | | | | | | | |
| EP080-SD: Benzene 71-43-2 | 0.2 | mg/kg | <0.2 | 1 mg/kg | 113 | 66 | 122 | | |
| EP080-SD: Toluene 108-88-3 | 0.2 | mg/kg | <0.2 | 1 mg/kg | 112 | 69 | 122 | | |
| EP080-SD: Ethylbenzene 100-41-4 | 0.2 | mg/kg | <0.2 | 1 mg/kg | 113 | 66 | 126 | | |
| EP080-SD: meta- & para-Xylene 108-38-3 | 0.2 | mg/kg | <0.2 | 2.5 mg/kg | 91.5 | 59 | 129 | | |
| 106-42-3 | | | | | | | | | |
| EP080-SD: ortho-Xylene 95-47-6 | 0.2 | mg/kg | <0.2 | 1 mg/kg | 109 | 66 | 126 | | |
| EP090: Organotin Compounds (QCLot: 2105989) | | | | | | | | | |

Page : 9 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: SOIL | | | Method Blank (MB) | Laboratory Control Spike (LCS) Report | | | | |
|---|----------------|--------|-------------------|---------------------------------------|---------------|------------|------|------|
| | | Report | Spike | Spike Recovery (%) | Recovery | Limits (%) | | |
| Method: Compound | CAS Number | LOR | Unit | Result | Concentration | LCS | Low | High |
| EP090: Organotin Compounds (QCLot: 210598 | 9) - continued | | | | | | | |
| EP090: Tributyltin | 56573-85-4 | 0.5 | μgSn/kg | <0.5 | 1.25 μgSn/kg | 46.4 | 19.5 | 129 |

Page : 10 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Matrix Spike (MS) Report

The quality control term Matrix Spike (MS) refers to an intralaboratory split sample spiked with a representative set of target analytes. The purpose of this QC parameter is to monitor potential matrix effects on analyte recoveries. Static Recovery Limits as per laboratory Data Quality Objectives (DQOs). Ideal recovery ranges stated may be waived in the event of sample matrix interference.

| Sub-Matrix: SOIL | | | Matrix Spike (MS) Repo | ort | | | |
|----------------------|---------------------------------|-------------------------------|------------------------|---------------|--------------------|----------|------------|
| | | | | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | Concentration | MS | Low | High |
| G020-SD: Total Met | tals in Sediments by ICPMS(Qu | CLot: 2115274) | | | | | |
| EM1114568-001 | A1 | EG020-SD: Arsenic | 7440-38-2 | 50 mg/kg | 105 | 70 | 130 |
| | | EG020-SD: Cadmium | 7440-43-9 | 50 mg/kg | 108 | 70 | 130 |
| | | EG020-SD: Chromium | 7440-47-3 | 50 mg/kg | 104 | 70 | 130 |
| | | EG020-SD: Copper | 7440-50-8 | 250 mg/kg | 98.2 | 70 | 130 |
| | | EG020-SD: Lead | 7439-92-1 | 250 mg/kg | 107 | 70 | 130 |
| | | EG020-SD: Nickel | 7440-02-0 | 50 mg/kg | 104 | 70 | 130 |
| | | EG020-SD: Zinc | 7440-66-6 | 250 mg/kg | 88.9 | 70 | 130 |
| G035T: Total Reco | verable Mercury by FIMS (QCL | ot: 2115273) | | | | | |
| EM1114568-001 | A1 | EG035T-LL: Mercury | 7439-97-6 | 0.50 mg/kg | 75.8 | 70 | 130 |
| P068A: Organochic | orine Pesticides (OC) (QCLot: 2 | 106215) | | | | | |
| EM1114568-001 | A1 | EP068: gamma-BHC | 58-89-9 | 0.5 mg/kg | 101 | 70 | 130 |
| | | EP068: Heptachlor | 76-44-8 | 0.5 mg/kg | 92.0 | 70 | 130 |
| | | EP068: Aldrin | 309-00-2 | 0.5 mg/kg | 80.6 | 70 | 130 |
| | | EP068: Dieldrin | 60-57-1 | 0.5 mg/kg | 95.6 | 70 | 130 |
| | | EP068: Endrin | 72-20-8 | 2 mg/kg | 102 | 70 | 130 |
| | | EP068: 4.4`-DDT | 50-29-3 | 2 mg/kg | 76.9 | 70 | 130 |
| P068B: Organopho | sphorus Pesticides (OP) (QCLo | ot: 2106215) | | | <u>'</u> | | |
| EM1114568-001 | A1 | EP068: Diazinon | 333-41-5 | 0.5 mg/kg | 109 | 70 | 130 |
| | | EP068: Chlorpyrifos-methyl | 5598-13-0 | 0.5 mg/kg | 104 | 70 | 130 |
| | | EP068: Pirimphos-ethyl | 23505-41-1 | 0.5 mg/kg | 104 | 70 | 130 |
| | | EP068: Bromophos-ethyl | 4824-78-6 | 0.5 mg/kg | 80.1 | 70 | 130 |
| | | EP068: Prothiofos | 34643-46-4 | 0.5 mg/kg | 72.0 | 70 | 130 |
| P080-SD / FP071-SI | D: Total Petroleum Hydrocarbo | | | | | | |
| M1114568-002 | B1 | EP071-SD: C10 - C14 Fraction | | 19.75 mg/kg | 76.7 | 70 | 130 |
| | | EP071-SD: C15 - C28 Fraction | | 87.25 mg/kg | 92.9 | 70 | 130 |
| | | EP071-SD: C29 - C36 Fraction | | 60 mg/kg | 95.6 | 70 | 130 |
| P080-SD / FP071-SI | D: Total Petroleum Hydrocarbo | | | 5 5 | | | |
| EM1114568-001 | A1 | EP080-SD: C6 - C9 Fraction | | 25 mg/kg | # 130 | 70 | 130 |
| P080-SD: BTEXN (| • • • | El 000-0B. Co - Co i faction | | _0gg | 100 | . • | |
| EM1114568-001 | | EP080-SD: Benzene | 71-43-2 | 2.5 mg/kg | 117 | 70 | 130 |
| _1011117000-001 | A1 | | 108-88-3 | 2.5 mg/kg | 110 | 70 | 130 |
| | | EP080-SD: Toluene | 100-68-3 | 2.5 mg/kg | 107 | 70 | 130 |
| | | EP080-SD: Ethylbenzene | | 2.5 mg/kg | 107 | 70 | 130 |
| | | EP080-SD: meta- & para-Xylene | 108-38-3 106-42-3 | 2.0 mg/kg | 109 | 70 | 130 |

Page : 11 of 11

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: SOIL | | Matrix Spike (MS) Report | | | | | |
|----------------------|-----------------------------|--------------------------|------------|---------------|--------------------|----------|------------|
| | | | | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | Concentration | MS | Low | High |
| EP080-SD: BTEXN (| QCLot: 2109665) - continued | | | | | | |
| EM1114568-001 | A1 | EP080-SD: ortho-Xylene | 95-47-6 | 2.5 mg/kg | 109 | 70 | 130 |
| EP090: Organotin Co | ompounds (QCLot: 2105989) | | | | | | |
| EM1114568-002 | B1 | EP090: Tributyltin | 56573-85-4 | 1.25 μgSn/kg | 52.5 | 20 | 130 |

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

CERTIFICATE OF ANALYSIS

Work Order : **EM1114568** Page : 1 of 12

Amendment : 1

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : DR GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

 Telephone
 : +61 08 8424 3800
 Telephone
 : +61-3-8549 9608

 Facsimile
 : +61 08 8424 3810
 Facsimile
 : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

 Order number
 : --

 C-O-C number
 : --

 Sampler
 : GB

 Issue Date
 : 22-DEC-2011

 Issue Date
 : 28-FEB-2012

Sampler : GB Site : ----

No. of samples received : 14

Quote number : EN/003/10 No. of samples analysed : 14

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for release.

This Certificate of Analysis contains the following information:

- General Comments
- Analytical Results
- Surrogate Control Limits



NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

| Signatories | Position | Accreditation Category | |
|-------------------|-------------------------------|------------------------|--|
| Di-An Dao | | Sydney Inorganics | |
| Matt Frost | Senior Organic Chemist | Brisbane Organics | |
| Phalak Inthaksone | Laboratory Manager - Organics | Sydney Organics | |
| Wisam Marassa | Inorganics Coordinator | Sydney Inorganics | |
| | | | |

Environmental Division Melbourne
Part of the ALS Laboratory Group

4 Westall Rd Springvale VIC Australia 3171

Tel. +61-3-8549 9600 Fax. +61-3-8549 9601 www.alsglobal.com

A Campbell Brothers Limited Company

Page : 2 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

When sampling time information is not provided by the client, sampling dates are shown without a time component. In these instances, the time component has been assumed by the laboratory for processing purposes.

Key: CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

^ = This result is computed from individual analyte detections at or above the level of reporting

- 27/2/12 This report has been amended and re-released to allow the reporting of additional analytical data.
- All analysis with the exception of TBT conducted by ALS Sydney, NATA accreditation no. 825, site no 10911.
- EG020: LCS recoveries for particular element(s) fall outside ALS Dynamic control limit, however, they are within the acceptance criteria based on ALS DQO. No further action is required.
- TBT conducted by ALS Brisbane, NATA Site No. 818.
- Tributyltin: Sample C6 shows poor surrogate recovery due to matrix interference. Confirmed by re-extraction and re-analysis.

Page : 3 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: SEDIMENT | | Clie | ent sample ID | A 1 | B1 | C1 | D1 | A4 |
|---------------------------------------|------------|--------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Cl | ient samplii | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114568-001 | EM1114568-002 | EM1114568-003 | EM1114568-004 | EM1114568-005 |
| EA055: Moisture Content | | | | | | | | |
| Moisture Content (dried @ 103°C) | | 1.0 | % | 29.0 | 34.5 | 35.6 | 42.5 | 29.8 |
| EG005T: Total Metals by ICP-AES | | | | | | | | |
| Barium | 7440-39-3 | 10 | mg/kg | <10 | <10 | <10 | <10 | <10 |
| Beryllium | 7440-41-7 | 1 | mg/kg | <1 | <1 | <1 | <1 | <1 |
| Iron | 7439-89-6 | 50 | mg/kg | 2460 | 3520 | 4070 | 5020 | 3210 |
| EG020-SD: Total Metals in Sediments I | by ICPMS | | | | | | | |
| Arsenic | 7440-38-2 | 1.00 | mg/kg | 2.54 | 2.08 | 3.76 | 2.88 | 4.64 |
| Cadmium | 7440-43-9 | 0.1 | mg/kg | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 |
| Chromium | 7440-47-3 | 1.0 | mg/kg | 5.5 | 10.6 | 10.4 | 16.1 | 6.7 |
| Copper | 7440-50-8 | 1.0 | mg/kg | 1.3 | 2.1 | 1.5 | 2.2 | <1.0 |
| Cobalt | 7440-48-4 | 0.5 | mg/kg | <0.5 | 0.6 | 0.6 | 0.8 | <0.5 |
| Lead | 7439-92-1 | 1.0 | mg/kg | 1.1 | 2.1 | 2.0 | 2.9 | 1.4 |
| Manganese | 7439-96-5 | 10 | mg/kg | 10 | 12 | 11 | 14 | 11 |
| Nickel | 7440-02-0 | 1.0 | mg/kg | <1.0 | 2.0 | 1.9 | 3.2 | 1.1 |
| Vanadium | 7440-62-2 | 2.0 | mg/kg | <2.0 | <2.0 | 3.7 | 8.0 | 5.2 |
| Zinc | 7440-66-6 | 1.0 | mg/kg | 1.4 | 2.7 | 3.2 | 3.6 | 1.9 |
| EG035T: Total Recoverable Mercury b | y FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.01 | mg/kg | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EP068A: Organochlorine Pesticides (C | OC) | | | | | | | |
| alpha-BHC | 319-84-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Hexachlorobenzene (HCB) | 118-74-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| beta-BHC | 319-85-7 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| gamma-BHC | 58-89-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| delta-BHC | 319-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Heptachlor | 76-44-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Aldrin | 309-00-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Heptachlor epoxide | 1024-57-3 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| trans-Chlordane | 5103-74-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| alpha-Endosulfan | 959-98-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| cis-Chlordane | 5103-71-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Dieldrin | 60-57-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| 4.4`-DDE | 72-55-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Endrin | 72-20-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| beta-Endosulfan | 33213-65-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| 4.4`-DDD | 72-54-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Endrin aldehyde | 7421-93-4 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Endosulfan sulfate | 1031-07-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| 4.4`-DDT | 50-29-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |

Page : 4 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: SEDIMENT | | Clie | ent sample ID | A 1 | B1 | C1 | D1 | A4 |
|---------------------------------------|--------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Cl | ent samplir | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114568-001 | EM1114568-002 | EM1114568-003 | EM1114568-004 | EM1114568-005 |
| EP068A: Organochlorine Pesticides (| OC) - Continued | | | | | | | |
| Endrin ketone | 53494-70-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Methoxychlor | 72-43-5 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| EP068B: Organophosphorus Pesticid | les (OP) | | | | | | | |
| Dichlorvos | 62-73-7 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Demeton-S-methyl | 919-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Monocrotophos | 6923-22-4 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Dimethoate | 60-51-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Diazinon | 333-41-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Chlorpyrifos-methyl | 5598-13-0 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Parathion-methyl | 298-00-0 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Malathion | 121-75-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Fenthion | 55-38-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Chlorpyrifos | 2921-88-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Parathion | 56-38-2 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Pirimphos-ethyl | 23505-41-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Chlorfenvinphos | 470-90-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Bromophos-ethyl | 4824-78-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Fenamiphos | 22224-92-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Prothiofos | 34643-46-4 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Ethion | 563-12-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Carbophenothion | 786-19-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Azinphos Methyl | 86-50-0 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EP080/071: Total Recoverable Hydroc | carbons - NEPM 201 | 0 Draft | | | | | | |
| >C10 - C16 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| >C16 - C34 Fraction | | 3 | mg/kg | 30 | 32 | 26 | 56 | 27 |
| >C34 - C40 Fraction | | 5 | mg/kg | <5 | 8 | <5 | 12 | <5 |
| >C10 - C40 Fraction (sum) | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| EP080-SD / EP071-SD: Total Petroleum | m Hydrocarbons | | | | | | | |
| C6 - C9 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| C10 - C14 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| C15 - C28 Fraction | | 3 | mg/kg | 23 | 22 | 19 | 39 | 20 |
| C29 - C36 Fraction | | 5 | mg/kg | 5 | 9 | 6 | 14 | <5 |
| [^] C10 - C36 Fraction (sum) | | 3 | mg/kg | 28 | 31 | 25 | 53 | 20 |
| EP080-SD / EP071-SD: Total Recovera | able Hydrocarbons | | | | | | | |
| C6 - C10 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| EP080-SD: BTEXN | | | | | | | | |
| Benzene | 71-43-2 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| | - | | | | | | | |

Page : 5 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: SEDIMENT | | Cli | ent sample ID | A 1 | B1 | C1 | D1 | A4 |
|----------------------------------|-------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | CI | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114568-001 | EM1114568-002 | EM1114568-003 | EM1114568-004 | EM1114568-005 |
| EP080-SD: BTEXN - Continued | | | | | | | | |
| Toluene | 108-88-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Ethylbenzene | 100-41-4 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| meta- & para-Xylene | 108-38-3 106-42-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| ortho-Xylene | 95-47-6 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| ^ Total Xylenes | 1330-20-7 | 0.5 | mg/kg | <0.5 | <0.5 | <0.5 | <0.5 | <0.5 |
| ^ Sum of BTEX | | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Naphthalene | 91-20-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| EP090: Organotin Compounds | | | | | | | | |
| Tributyltin | 56573-85-4 | 0.5 | μgSn/kg | <0.5 | <0.5 | <0.5 | <0.5 | <0.5 |
| EP068S: Organochlorine Pesticide | e Surrogate | | | | | | | |
| Dibromo-DDE | 21655-73-2 | 0.1 | % | 94.5 | 85.5 | 84.6 | 91.0 | 72.0 |
| EP068T: Organophosphorus Pest | icide Surrogate | | | | | | | |
| DEF | 78-48-8 | 0.1 | % | 93.0 | 82.8 | 81.7 | 90.1 | 63.8 |
| EP080-SD: TPH(V)/BTEX Surrogat | es | | | | | | | |
| 1.2-Dichloroethane-D4 | 17060-07-0 | 0.1 | % | 114 | 126 | 107 | 119 | 107 |
| Toluene-D8 | 2037-26-5 | 0.1 | % | 104 | 108 | 98.4 | 103 | 99.7 |
| 4-Bromofluorobenzene | 460-00-4 | 0.1 | % | 88.4 | 94.4 | 87.8 | 91.1 | 86.4 |
| EP090S: Organotin Surrogate | | | | | | | | |
| Tripropyltin | | 0.1 | % | 46.4 | 92.8 | 66.5 | 108 | 121 |

Page : 6 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: SEDIMENT | | Clie | ent sample ID | B4 | C4 | D4 | A5 | C5 |
|---------------------------------------|------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Cl | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114568-006 | EM1114568-007 | EM1114568-008 | EM1114568-009 | EM1114568-010 |
| EA055: Moisture Content | | | | | | | | |
| Moisture Content (dried @ 103°C) | | 1.0 | % | 32.3 | 34.6 | 37.3 | 20.2 | 29.6 |
| EG005T: Total Metals by ICP-AES | | | | | | | | |
| Barium | 7440-39-3 | 10 | mg/kg | <10 | <10 | <10 | <10 | <10 |
| Beryllium | 7440-41-7 | 1 | mg/kg | <1 | <1 | <1 | <1 | <1 |
| Iron | 7439-89-6 | 50 | mg/kg | 3940 | 5830 | 5400 | 3000 | 5060 |
| EG020-SD: Total Metals in Sediments b | by ICPMS | | | | | | | |
| Arsenic | 7440-38-2 | 1.00 | mg/kg | 3.74 | 7.36 | 5.06 | 6.71 | 7.38 |
| Cadmium | 7440-43-9 | 0.1 | mg/kg | <0.1 | <0.1 | <0.1 | <0.1 | <0.1 |
| Chromium | 7440-47-3 | 1.0 | mg/kg | 9.9 | 13.5 | 13.7 | 5.8 | 8.4 |
| Copper | 7440-50-8 | 1.0 | mg/kg | 1.2 | 1.6 | 2.0 | <1.0 | <1.0 |
| Cobalt | 7440-48-4 | 0.5 | mg/kg | 0.6 | 0.6 | 0.6 | <0.5 | <0.5 |
| Lead | 7439-92-1 | 1.0 | mg/kg | 1.9 | 2.3 | 2.6 | <1.0 | 1.6 |
| Manganese | 7439-96-5 | 10 | mg/kg | 12 | 14 | 12 | <10 | <10 |
| Nickel | 7440-02-0 | 1.0 | mg/kg | 1.7 | 1.9 | 2.1 | <1.0 | <1.0 |
| Vanadium | 7440-62-2 | 2.0 | mg/kg | 5.5 | 12.3 | 14.2 | 10.1 | 10.9 |
| Zinc | 7440-66-6 | 1.0 | mg/kg | 2.6 | 2.9 | 3.1 | 1.7 | 1.4 |
| EG035T: Total Recoverable Mercury b | y FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.01 | mg/kg | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EP068A: Organochlorine Pesticides (O | C) | | | | | | | |
| alpha-BHC | 319-84-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Hexachlorobenzene (HCB) | 118-74-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| beta-BHC | 319-85-7 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| gamma-BHC | 58-89-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| delta-BHC | 319-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Heptachlor | 76-44-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Aldrin | 309-00-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Heptachlor epoxide | 1024-57-3 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| trans-Chlordane | 5103-74-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| alpha-Endosulfan | 959-98-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| cis-Chlordane | 5103-71-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Dieldrin | 60-57-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| 4.4`-DDE | 72-55-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Endrin | 72-20-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| beta-Endosulfan | 33213-65-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| 4.4`-DDD | 72-54-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Endrin aldehyde | 7421-93-4 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Endosulfan sulfate | 1031-07-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| 4.4`-DDT | 50-29-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |

Page : 7 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: SEDIMENT | | Clie | ent sample ID | B4 | C4 | D4 | A5 | C5 |
|------------------------------------|---------------------|--------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | CI | ient samplii | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114568-006 | EM1114568-007 | EM1114568-008 | EM1114568-009 | EM1114568-010 |
| EP068A: Organochlorine Pesticides | s (OC) - Continued | | | | | | | |
| Endrin ketone | 53494-70-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Methoxychlor | 72-43-5 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| EP068B: Organophosphorus Pestic | cides (OP) | | | | | | | |
| Dichlorvos | 62-73-7 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Demeton-S-methyl | 919-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Monocrotophos | 6923-22-4 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Dimethoate | 60-51-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Diazinon | 333-41-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Chlorpyrifos-methyl | 5598-13-0 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Parathion-methyl | 298-00-0 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Malathion | 121-75-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Fenthion | 55-38-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Chlorpyrifos | 2921-88-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Parathion | 56-38-2 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Pirimphos-ethyl | 23505-41-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Chlorfenvinphos | 470-90-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Bromophos-ethyl | 4824-78-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Fenamiphos | 22224-92-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Prothiofos | 34643-46-4 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Ethion | 563-12-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Carbophenothion | 786-19-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Azinphos Methyl | 86-50-0 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EP080/071: Total Recoverable Hydr | ocarbons - NEPM 201 | 0 Draft | | | | | | |
| >C10 - C16 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| >C16 - C34 Fraction | | 3 | mg/kg | 53 | 31 | 30 | 21 | 28 |
| >C34 - C40 Fraction | | 5 | mg/kg | 11 | <5 | 7 | <5 | 5 |
| >C10 - C40 Fraction (sum) | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| EP080-SD / EP071-SD: Total Petrole | eum Hydrocarbons | | | | | | | |
| C6 - C9 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| C10 - C14 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| C15 - C28 Fraction | | 3 | mg/kg | 37 | 22 | 20 | 16 | 20 |
| C29 - C36 Fraction | | 5 | mg/kg | 14 | 7 | 9 | <5 | 7 |
| C10 - C36 Fraction (sum) | | 3 | mg/kg | 51 | 29 | 29 | 16 | 27 |
| EP080-SD / EP071-SD: Total Recove | erable Hydrocarbons | | | | | | | |
| C6 - C10 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | <3 |
| EP080-SD: BTEXN | | | | | | | | |
| Benzene | 71-43-2 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |

Page : 8 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: SEDIMENT | | Cli | ent sample ID | B4 | C4 | D4 | A5 | C5 |
|----------------------------------|-------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | CI | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114568-006 | EM1114568-007 | EM1114568-008 | EM1114568-009 | EM1114568-010 |
| EP080-SD: BTEXN - Continued | | | | | | | | |
| Toluene | 108-88-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Ethylbenzene | 100-41-4 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| meta- & para-Xylene | 108-38-3 106-42-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| ortho-Xylene | 95-47-6 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| ^ Total Xylenes | 1330-20-7 | 0.5 | mg/kg | <0.5 | <0.5 | <0.5 | <0.5 | <0.5 |
| ^ Sum of BTEX | | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| Naphthalene | 91-20-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 |
| EP090: Organotin Compounds | | | | | | | | |
| Tributyltin | 56573-85-4 | 0.5 | μgSn/kg | <0.5 | <0.5 | <0.5 | <0.5 | <0.5 |
| EP068S: Organochlorine Pesticide | Surrogate | | | | | | | |
| Dibromo-DDE | 21655-73-2 | 0.1 | % | 90.3 | 85.8 | 96.1 | 93.9 | 95.9 |
| EP068T: Organophosphorus Pestic | cide Surrogate | | | | | | | |
| DEF | 78-48-8 | 0.1 | % | 86.4 | 81.1 | 81.4 | 90.4 | 94.8 |
| EP080-SD: TPH(V)/BTEX Surrogate | s | | | | | | | |
| 1.2-Dichloroethane-D4 | 17060-07-0 | 0.1 | % | 105 | 109 | 123 | 121 | 115 |
| Toluene-D8 | 2037-26-5 | 0.1 | % | 96.1 | 102 | 118 | 111 | 106 |
| 4-Bromofluorobenzene | 460-00-4 | 0.1 | % | 82.2 | 87.0 | 100 | 96.2 | 90.3 |
| EP090S: Organotin Surrogate | | | | | | | | |
| Tripropyltin | | 0.1 | % | 106 | 106 | 87.5 | 111 | 109 |

Page : 9 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: SEDIMENT | | Clie | ent sample ID | D5 | В6 | C6 | D6 | |
|---|------------------------|--------------|----------------|--------------------|-------------------|-------------------|-------------------|--|
| | CI | ient samplii | ng date / time | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | |
| Compound | CAS Number | LOR | Unit | EM1114568-011 | EM1114568-012 | EM1114568-013 | EM1114568-014 | |
| EA055: Moisture Content | CAS Number | | 0 | | | | | |
| | | 1.0 | % | 25.7 | 32.9 | 38.8 | 31.3 | |
| Moisture Content (dried @ 103°C) | | 1.0 | 70 | 25.1 | 32.3 | 30.0 | 31.3 | |
| EG005T: Total Metals by ICP-AES Barium | 7440.00.0 | 10 | ma/ka | <10 | <10 | <10 | <10 | |
| Beryllium | 7440-39-3 | 10 | mg/kg | <1 | <10 | <10 | <1 | |
| Iron | 7440-41-7 7439-89-6 | 50 | mg/kg mg/kg | 6540 | 3480 | 14000 | 13500 | |
| • | | 30 | mg/kg | 0040 | 3400 | 14000 | 13500 | |
| EG020-SD: Total Metals in Sediments b | | 4.00 | | 0.00 | 40.0 | F4.4 | 20.0 | |
| Arsenic | 7440-38-2 | 1.00 | mg/kg | 8.99 | 10.3 | 51.1 | 29.6 | |
| Cadmium | 7440-43-9 | 0.1 | mg/kg | <0.1 | <0.1 | <0.1 | <0.1 | |
| Conner | 7440-47-3 | 1.0 | mg/kg | 13.5 | 8.4 | 23.2 <1.0 | 24.0 | |
| Copper Cobalt | 7440-50-8 | 1.0 0.5 | mg/kg | <1.0 0.5 | 2.4 <0.5 | <1.0 | <1.0 <0.5 | |
| | 7440-48-4 | 1.0 | mg/kg | | 1.1 | 2.1 | | |
| Lead | 7439-92-1 | | mg/kg | 1.8 14 | 1.1 <10 | 10 | 1.8 <10 | |
| Manganese | 7439-96-5 | 1.0 | mg/kg | 1.3 | <1.0 | <1.0 | <1.0 | |
| Nickel Vanadium | 7440-02-0 | 2.0 | mg/kg mg/kg | 19.1 | 13.6 | 57.4 | 50.7 | |
| Zinc | 7440-62-2 | 1.0 | | 2.1 | 2.0 | 1.2 | 1.4 | |
| | 7440-66-6 | 1.0 | mg/kg | Z. I | 2.0 | 1.2 | 1.4 | |
| EG035T: Total Recoverable Mercury by | | 0.04 | | 10.04 | -0.04 | .0.04 | 10.04 | |
| Mercury | 7439-97-6 | 0.01 | mg/kg | <0.01 | <0.01 | <0.01 | <0.01 | |
| EP068A: Organochlorine Pesticides (O | · | | | | 1 | | | |
| alpha-BHC | 319-84-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Hexachlorobenzene (HCB) | 118-74-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| beta-BHC | 319-85-7 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| gamma-BHC | 58-89-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| delta-BHC | 319-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Heptachlor | 76-44-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Aldrin | 309-00-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Heptachlor epoxide | 1024-57-3 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| trans-Chlordane | 5103-74-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| alpha-Endosulfan | 959-98-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| cis-Chlordane | 5103-71-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Dieldrin | 60-57-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| 4.4`-DDE | 72-55-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Endrin | 72-20-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| beta-Endosulfan | 33213-65-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| 4.4`-DDD | 72-54-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Endrin aldehyde | 7421-93-4 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Endosulfan sulfate | 1031-07-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| 4.4`-DDT | 50-29-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |

Page : 10 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Compound | Cli | ient samnlir | | | | | | |
|--------------------------------------|-------------------|----------------|----------------|-------------------|-------------------|-------------------|-------------------|--|
| Compound | | ciit sairipiii | ng date / time | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | |
| | CAS Number | LOR | Unit | EM1114568-011 | EM1114568-012 | EM1114568-013 | EM1114568-014 | |
| EP068A: Organochlorine Pesticides (C | DC) - Continued | | | | | | | |
| Endrin ketone | 53494-70-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Methoxychlor | 72-43-5 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| EP068B: Organophosphorus Pesticide | es (OP) | | | | | | | |
| Dichlorvos | 62-73-7 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Demeton-S-methyl | 919-86-8 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Monocrotophos | 6923-22-4 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| Dimethoate | 60-51-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Diazinon | 333-41-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Chlorpyrifos-methyl | 5598-13-0 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Parathion-methyl | 298-00-0 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| Malathion | 121-75-5 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Fenthion | 55-38-9 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Chlorpyrifos | 2921-88-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Parathion | 56-38-2 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| Pirimphos-ethyl | 23505-41-1 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Chlorfenvinphos | 470-90-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Bromophos-ethyl | 4824-78-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Fenamiphos | 22224-92-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Prothiofos | 34643-46-4 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Ethion | 563-12-2 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Carbophenothion | 786-19-6 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| Azinphos Methyl | 86-50-0 | 0.05 | mg/kg | <0.05 | <0.05 | <0.05 | <0.05 | |
| EP080/071: Total Recoverable Hydroca | arbons - NEPM 201 | 0 Draft | | | | | | |
| >C10 - C16 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | |
| >C16 - C34 Fraction | | 3 | mg/kg | 24 | 36 | 35 | 35 | |
| >C34 - C40 Fraction | | 5 | mg/kg | <5 | <5 | 5 | <5 | |
| >C10 - C40 Fraction (sum) | | 3 | mg/kg | <3 | <3 | <3 | <3 | |
| EP080-SD / EP071-SD: Total Petroleun | n Hydrocarbons | | | | | | | |
| C6 - C9 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | |
| C10 - C14 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | |
| C15 - C28 Fraction | | 3 | mg/kg | 17 | 31 | 26 | 26 | |
| C29 - C36 Fraction | | 5 | mg/kg | 6 | 12 | 6 | 6 | |
| ^ C10 - C36 Fraction (sum) | | 3 | mg/kg | 23 | 43 | 32 | 32 | |
| EP080-SD / EP071-SD: Total Recovera | ble Hydrocarbons | | | | | | | |
| C6 - C10 Fraction | | 3 | mg/kg | <3 | <3 | <3 | <3 | |
| EP080-SD: BTEXN | | | | | | | | |
| Benzene | 71-43-2 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |

Page : 11 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: SEDIMENT | | Cli | ent sample ID | D5 | B6 | C6 | D6 | |
|--------------------------------|-------------------|-------------|-----------------|-------------------|-------------------|-------------------|-------------------|--|
| | Cl | ient sampli | ing date / time | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | |
| Compound | CAS Number | LOR | Unit | EM1114568-011 | EM1114568-012 | EM1114568-013 | EM1114568-014 | |
| EP080-SD: BTEXN - Continued | | | | | | | | |
| Toluene | 108-88-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| Ethylbenzene | 100-41-4 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| meta- & para-Xylene | 108-38-3 106-42-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| ortho-Xylene | 95-47-6 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| ^ Total Xylenes | 1330-20-7 | 0.5 | mg/kg | <0.5 | <0.5 | <0.5 | <0.5 | |
| ^ Sum of BTEX | | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| Naphthalene | 91-20-3 | 0.2 | mg/kg | <0.2 | <0.2 | <0.2 | <0.2 | |
| EP090: Organotin Compounds | | | | | | | | |
| Tributyltin | 56573-85-4 | 0.5 | μgSn/kg | <0.5 | <0.5 | <0.5 | <0.5 | |
| EP068S: Organochlorine Pestici | de Surrogate | | | | | | | |
| Dibromo-DDE | 21655-73-2 | 0.1 | % | 90.1 | 94.4 | 97.1 | 83.9 | |
| EP068T: Organophosphorus Pe | sticide Surrogate | | | | | | | |
| DEF | 78-48-8 | 0.1 | % | 87.6 | 100 | 90.6 | 75.4 | |
| EP080-SD: TPH(V)/BTEX Surroga | ates | | | | | | | |
| 1.2-Dichloroethane-D4 | 17060-07-0 | 0.1 | % | 121 | 111 | 119 | 130 | |
| Toluene-D8 | 2037-26-5 | 0.1 | % | 103 | 105 | 110 | 117 | |
| 4-Bromofluorobenzene | 460-00-4 | 0.1 | % | 89.0 | 87.8 | 95.1 | 102 | |
| EP090S: Organotin Surrogate | | | | | | | | |
| Tripropyltin | | 0.1 | % | 110 | 104 | 146 | 110 | |

Page : 12 of 12

Work Order : EM1114568 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609

ALS

Surrogate Control Limits

| Sub-Matrix: SEDIMENT | | Recovery | Limits (%) |
|--|------------|----------|------------|
| Compound | CAS Number | Low | High |
| EP068S: Organochlorine Pesticide Surrogate | | | |
| Dibromo-DDE | 21655-73-2 | 19.5 | 167.0 |
| EP068T: Organophosphorus Pesticide Surrogate | | | |
| DEF | 78-48-8 | 22.7 | 163.5 |
| EP080-SD: TPH(V)/BTEX Surrogates | | | |
| 1.2-Dichloroethane-D4 | 17060-07-0 | 67 | 137 |
| Toluene-D8 | 2037-26-5 | 74 | 134 |
| 4-Bromofluorobenzene | 460-00-4 | 73 | 137 |
| EP090S: Organotin Surrogate | | | |
| Tripropyltin | | 35 | 130 |

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

INTERPRETIVE QUALITY CONTROL REPORT

Work Order : **EM1114567** Page : 1 of 13

Amendment : 1

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

Telephone : +61 08 8424 3800 Telephone : +61-3-8549 9608 Facsimile : +61 08 8424 3810 Facsimile : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

: 60

Site : ----

 C-O-C number
 :--- Date Samples Received
 : 22-DEC-2011

 Sampler
 : GB
 Issue Date
 : 28-FEB-2012

Order number : ---No. of samples received

Quote number : ADBQ/037/11 No. of samples analysed : 60

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for release.

This Interpretive Quality Control Report contains the following information:

- Analysis Holding Time Compliance
- Quality Control Parameter Frequency Compliance
- Brief Method Summaries
- Summary of Outliers

Page : 2 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Analysis Holding Time Compliance

The following report summarises extraction / preparation and analysis times and compares with recommended holding times. Dates reported represent first date of extraction or analysis and precludes subsequent dilutions and reruns. Information is also provided re the sample container (preservative) from which the analysis aliquot was taken. Elapsed period to analysis represents number of days from sampling where no extraction / digestion is involved or period from extraction / digestion where this is present. For composite samples, sampling date is assumed to be that of the oldest sample contributing to the composite. Sample date for laboratory produced leachates is assumed as the completion date of the leaching process. Outliers for holding time are based on USEPA SW 846, APHA, AS and NEPM (1999). A listing of breaches is provided in the Summary of Outliers.

Holding times for leachate methods (excluding elutriates) vary according to the analytes being determined on the resulting solution. For non-volatile analytes, the holding time compliance assessment compares the leach date with the shortest analyte holding time for the equivalent soil method. These soil holding times are: Organics (14 days); Mercury (28 days) & other metals (180 days). A recorded breach therefore does not guarantee a breach for all non-volatile parameters.

Matrix: WATER Evaluation: ▼ = Holding time breach; ✓ = Within holding time.

| Matrix: WATER | | | | | Evaluation | . × = notating time | breach; ∨ = withir | i nolaling tir |
|---------------------------------|------|-------------|----------------|-------------------------|------------|---------------------|--------------------|----------------|
| flethod | | Sample Date | E | ktraction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| A025: Suspended Solids | | | | | | | | |
| Clear Plastic Bottle - Natural | | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | | | 23-DEC-2011 | 26-DEC-2011 | 1 |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| B4S, | C4S, | | | | | | | |
| D4S, | X4S, | | | | | | | |
| Y4S, | AaS, | | | | | | | |
| BbS, | CcS, | | | | | | | |
| DdS, | A5S, | | | | | | | |
| B5S, | C5S, | | | | | | | |
| D5S, | A6S, | | | | | | | |
| B6S, | C6S, | | | | | | | |
| D6S, | A1B, | | | | | | | |
| B1B, | C1B, | | | | | | | |
| D1B, | A2B, | | | | | | | |
| B2B, | C2B, | | | | | | | |
| D2B, | A3B, | | | | | | | |
| B3B, | C3B, | | | | | | | |
| D3B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 3 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Matrix: WATER | | | | | Evaluation | : x = Holding time | breach ; ✓ = Within | n holding tim |
|----------------------------------|--|-------------|----------------|-------------------------|------------|--------------------|---------------------|---------------|
| Method | | Sample Date | E) | ktraction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EG020T: Total Metals by ICP-M | NS CONTRACTOR OF THE CONTRACTO | | | | | | | |
| Clear Plastic Bottle - Unfiltere | ed; Lab-acidified | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | 23-DEC-2011 | 16-JUN-2012 | ✓ | 30-DEC-2011 | 16-JUN-2012 | 1 |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B | | | | | | | |
| Clear Plastic Bottle - Unfiltere | | | | | | | | |
| C3B, | D3B, | 19-DEC-2011 | 28-DEC-2011 | 16-JUN-2012 | ✓ | 30-DEC-2011 | 16-JUN-2012 | 1 |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 4 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Matrix: WATER | | | | | Evaluation: | : x = Holding time | breach ; ✓ = Within | n holding time |
|-----------------------------------|------------------|-------------|----------------|------------------------|-------------|--------------------|---------------------|----------------|
| Method | | Sample Date | Ex | traction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EG035T: Total Recoverable Me | ercury by FIMS | | | | | | | |
| Clear Plastic Bottle - Unfiltered | d; Lab-acidified | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | | | 05-JAN-2012 | 16-JAN-2012 | 1 |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 5 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Matrix: WATER | | | | | Evaluation | x = Holding time | breach ; ✓ = Within | holding time |
|------------------------------------|---------------|-------------|----------------|------------------------|------------|------------------|---------------------|--------------|
| Method | | Sample Date | Ex | traction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EK055G: Ammonia as N by Discr | rete Analyser | | | | | | | |
| Clear Plastic Bottle - Sulfuric Ad | cid | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | 16-JAN-2012 | | 06-JAN-2012 | 16-JAN-2012 | ✓ |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 6 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Matrix: WATER | | | | | Evaluation | : x = Holding time | breach ; ✓ = Within | n holding time |
|---------------------------------|-------------|-------------|----------------|------------------------|------------|--------------------|---------------------|----------------|
| Method | | Sample Date | Ex | traction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EK057G: Nitrite as N by Discret | te Analyser | | | | | | | |
| Clear Plastic Bottle - Natural | | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | 21-DEC-2011 | | 22-DEC-2011 | 21-DEC-2011 | * |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

: 7 of 13 Page

: EM1114567 Amendment 1 Work Order : SINCLAIR KNIGHT MERZ Client



| Matrix: WATER | | | | | Evaluation: | × = Holding time | breach ; ✓ = Withir | n holding time |
|---------------------------------|---------------------------------|-------------|----------------|------------------------|-------------|------------------|---------------------|----------------|
| Method | | Sample Date | Ex | traction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EK059G: Nitrite plus Nitrate | as N (NOx) by Discrete Analyser | | | | | | | |
| Clear Plastic Bottle - Sulfuri | ic Acid | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | 16-JAN-2012 | | 05-JAN-2012 | 16-JAN-2012 | ✓ |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 8 of 13

A3B,

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

B3B



| Date extracted Due for extraction Evaluation Date analysed Due for analysis Evaluation Evaluation Date analysed Due for analysis Evaluation Evaluation Evaluation Date analysed Due for analysis Evaluation Evaluation Evaluation Evaluation Date analysed Due for analysis Evaluation Evaluation Date analysed Due for analysed Due for analysed Evaluation Evaluation Date analysed Due for analysed Evaluation Evaluation Date analysed Da | Matrix: WATER | | _ | _ | | LvaiuatiOII | | breach ; ✓ = Within | i noluling til |
|---|-------------------------------------|----------------------|-------------|----------------|--------------------|-------------|---------------|---------------------|----------------|
| Total Kjeldahl Nitrogen By Discrete Analyser Clear Plastic Bottle - Sulfuric Acid | Method | | Sample Date | | | <u> </u> | | | 1 |
| Clear Plastic Bottle - Sulfuric Acid C3B. D3B. A4B. B4B. C4B. D4B. X4B. A4B. B4B. C6B. D4B. A5B. B5B. C6B. D6B. C6B. D6B. C6B. D6B. C6B. D6B. A6B. B6B. C6B. D6B. A6B. A6B. B1S. C1Car Plastic Bottle - Sulfuric Acid A1S. B1S. C1S. B1S. C2S. C2S. C2S. C2S. C2S. C2S. C2S. C2 | Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| C3B, D3B, B4B, B4B, B4B, C4B, D4B, Waller, Wa | EK061G: Total Kjeldahl Nitrogen I | By Discrete Analyser | | | | | | | |
| A4B, | | | | | | | | | |
| C4B, D4B, Y4B. Y4B. AB. Y4B. AB. BbB. CCB. ABB. BBB. CCB. ABB. BBB. CSB. ABB. BBB. CSB. | | | 19-DEC-2011 | 03-JAN-2012 | 16-JAN-2012 | ✓ | 03-JAN-2012 | 16-JAN-2012 | ✓ |
| X4B, | | | | | | | | | |
| ABB, | | | | | | | | | |
| CCB, DdB, A5B, B5B, CSB, DSB, A6B, B6B, C6B, D6B Clear Plastic Bottle - Sulfuric Acid A1S, B1S, D1S, D1S, A2S, B2S, C2S, D2S, A3S, B3S, C3S, D4S, A4S, B4S, C4S, D4S, X4S, B4S, C4S, D4S, X4S, B5S, C5S, D4S, A6S, B5S, C6S, D6S, A6S, B6S, C6S, D6S, A6S, B6S, C6S, D6S, A6S, B6S, C6S, D6S, A1B, B1B, C1B, D1B, A2B, B2B, | X4B, | Y4B, | | | | | | | |
| A5B, | AaB, | BbB, | | | | | | | |
| C5B, A6B, B6B, B6B, B6B, B6B, B6B, B6B, B6 | CcB, | DdB, | | | | | | | |
| A6B, C6B, D6B C6B, D6B C6B, D6B C6B, D6B D6B D6B D6B D6B D6B D6B D6B | A5B, | B5B, | | | | | | | |
| C6B, D6B Instruction of Decembration | C5B, | D5B, | | | | | | | |
| Clear Plastic Bottle - Sulfuric Acid A15, B1S, D15, A2S, B2S, C2S, D2S, B3S, A4S, B4S, C4S, D4S, A2S, B5S, C5S, D5S, B5S, C5S, D5S, B6S, C5S, D5S, B6S, C6S, C6S, C6S, C6S, C6S, C6S, C6S, C | A6B, | B6B, | | | | | | | |
| A1S, B1S, D1S, D1S, B2S, C2S, B2S, C2S, B2S, C3S, B3S, C3S, B4S, C4S, C5S, B5S, C5S, B5S, C5S, B5S, C5S, B5S, C6S, B6S, C6S, C6S, C6S, C6S, C6S, C6S, C6S, C | C6B, | D6B | | | | | | | |
| C1S, D1S, B2S, C2S, B2S, C2S, D2S, A3S, B3S, C3S, D3S, A4S, D4S, C4S, D4S, C4S, D4S, C5S, D6S, C5S, D6S, A6S, C5S, D6S, A6S, C6S, D6S, A6S, D6S, C6S, D6S, A6S, D6S, C6S, D6S, A1B, B1B, C1B, A2B, B2B, C1B, C1B, C1B, C1B, C1B, C1B, C1B, C1 | Clear Plastic Bottle - Sulfuric Aci | id | | | | | | | |
| A2S, | A1S, | B1S, | 19-DEC-2011 | 30-DEC-2011 | 16-JAN-2012 | ✓ | 30-DEC-2011 | 16-JAN-2012 | 1 |
| C2S, | C1S, | D1S, | | | | | | | |
| A3S, C3S, D3S, A4S, D4S, C4S, D4S, A8S, C6S, B6S, C6S, C6S, B6S, C6S, C6S, C6S, C6S, C6S, C6S, C6S, C | A2S, | B2S, | | | | | | | |
| C3S, | C2S, | D2S, | | | | | | | |
| A4S, C4S, D4S, D4S, X4S, A8S, BBS, CcS, DdS, B5S, C5S, B6S, B6S, C6S, B6S, B6S, C6S, B6S, B6S, C6S, B6S, B6S, C6S, B6S, B6S, B6S, C6S, B6S, B6S, B6S, B6S, B6S, B6S, B6S, B | A3S, | B3S, | | | | | | | |
| C4S, D4S, X4S, X4S, Y4S, AaS, BbS, CcS, DdS, B5S, C5S, D5S, A6S, B6S, C6S, D6S, A1B, B1B, B1B, B1B, B1B, B1B, B2B, B2B, B | C3S, | D3S, | | | | | | | |
| X4S, Y4S, BbS, CcS, DdS, A6S, B5S, C5S, D5S, A6S, B6S, C6S, B6S, B6S, C6S, C6S, C6S, C6S, C6S, C6S, C6S, C | A4S, | B4S, | | | | | | | |
| AaS, BbS, CcS, DdS, A5S, B5S, C5S, D5S, A6S, B6S, C6S, B6B, C6B, C6S, B6B, C6B, C6B, B6B, C6B, C6B, B6B, C6B, C | C4S, | D4S, | | | | | | | |
| AaS, BbS, CcS, DdS, A5S, B5S, C5S, D5S, A6S, B6S, C6S, B6B, C6B, C6S, B6B, C6B, C6B, B6B, C6B, C6B, B6B, C6B, C | X4S, | Y4S, | | | | | | | |
| A5S, B5S, C5S, D5S, A6S, B6S, C6S, B6S, B6S, C6S, B6S, C6S, D6S, A1B, B1B, C1B, B1B, B2B, B2B, | | BbS, | | | | | | | |
| C5S, D5S, A6S, B6S, C6S, D6S, D6S, A1B, B1B, C1B, D1B, A2B, B2B, | CcS, | DdS, | | | | | | | |
| A6S, B6S, D6S, D6S, A1B, B1B, C1B, D1B, B2B, B2B, | A5S, | B5S, | | | | | | | |
| A6S, B6S, D6S, D6S, A1B, B1B, C1B, D1B, B2B, B2B, | C5S, | D5S, | | | | | | | |
| A1B, B1B, C1B, D1B, A2B, B2B, | | B6S, | | | | | | | |
| C1B, D1B, A2B, B2B, | C6S, | D6S, | | | | | | | |
| C1B, D1B, A2B, B2B, | A1B, | B1B, | | | | | | | |
| | C1B, | | | | | | | | |
| | A2B, | B2B, | | | | | | | |
| | | | | | | | | | |

Page : 9 of 13

A2B,

C2B,

A3B,

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

B2B,

D2B, B3B



| Matrix: WATER | | | | | ⊏valuation | . ~ - Holding time | breach; ✓ = Withir | i noluling tin |
|------------------------------------|---------------------|-------------|----------------|------------------------|------------|--------------------|--------------------|----------------|
| Method | | Sample Date | E | traction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluatio |
| EK067G: Total Phosphorus as P t | y Discrete Analyser | | | | | | | |
| Clear Plastic Bottle - Sulfuric Ac | id | | | | | | | |
| C3B, | D3B, | 19-DEC-2011 | 03-JAN-2012 | 16-JAN-2012 | ✓ | 03-JAN-2012 | 16-JAN-2012 | ✓ |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |
| Clear Plastic Bottle - Sulfuric Ac | id | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | 30-DEC-2011 | 16-JAN-2012 | 1 | 30-DEC-2011 | 16-JAN-2012 | 1 |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| 015, | 515, | | | | | | | |

Page : 10 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609

Total Phosphorus as P By Discrete Analyser



Quality Control Parameter Frequency Compliance

The following report summarises the frequency of laboratory QC samples analysed within the analytical lot(s) in which the submitted sample(s) was(where) processed. Actual rate should be greater than or equal to the expected rate. A listing of breaches is provided in the Summary of Outliers.

EK067G

3

60

5.0

5.0

Evaluation: × = Quality Control frequency not within specification: ✓ = Quality Control frequency within specification. Matrix: WATER Quality Control Sample Type Quality Control Specification Count Rate (%) Method Evaluation Analytical Methods QC Regular Actual Expected Laboratory Duplicates (DUP) Ammonia as N by Discrete analyser EK055G 6 60 10.0 10.0 1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Nitrite and Nitrate as N (NOx) by Discrete Analyser 60 EK059G 6 10.0 10.0 ✓ NEPM 1999 Schedule B(3) and ALS QCS3 requirement Nitrite as N by Discrete Analyser EK057G 6 60 10.0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 Suspended Solids (High Level) EA025H 8 69 11.6 10.0 1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Kjeldahl Nitrogen as N By Discrete Analyser EK061G 6 60 10 0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 EG035T 6 60 10.0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Mercury by FIMS Total Metals by ICP-MS - Suite A EG020A-T 6 60 10.0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Phosphorus as P By Discrete Analyser EK067G 6 60 10.0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Laboratory Control Samples (LCS) Ammonia as N by Discrete analyser EK055G 60 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 3 5.0 5.0 Nitrite and Nitrate as N (NOx) by Discrete Analyser 3 60 5.0 EK059G 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 Nitrite as N by Discrete Analyser EK057G 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Suspended Solids (High Level) FA025H 4 69 5.8 5.0 1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Kjeldahl Nitrogen as N By Discrete Analyser EK061G 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 3 Total Mercury by FIMS EG035T 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement ✓ Total Metals by ICP-MS - Suite A EG020A-T 3 60 5.0 5.0 1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Phosphorus as P By Discrete Analyser EK067G 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Method Blanks (MB) 3 Ammonia as N by Discrete analyser EK055G 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 3 60 Nitrite and Nitrate as N (NOx) by Discrete Analyser EK059G 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 3 Nitrite as N by Discrete Analyser EK057G 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement EA025H 69 Suspended Solids (High Level) 4 5.8 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 5.0 1 Total Kjeldahl Nitrogen as N By Discrete Analyser 3 EK061G 60 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 5.0 Total Mercury by FIMS EG035T 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Metals by ICP-MS - Suite A EG020A-T 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 Total Phosphorus as P By Discrete Analyser EK067G 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Matrix Spikes (MS) Ammonia as N by Discrete analyser EK055G 3 60 ALS QCS3 requirement 5.0 5.0 Nitrite and Nitrate as N (NOx) by Discrete Analyser EK059G 3 60 5.0 5.0 ALS QCS3 requirement 3 Nitrite as N by Discrete Analyser EK057G 60 5.0 ALS QCS3 requirement 5.0 1 3 Total Kjeldahl Nitrogen as N By Discrete Analyser EK061G 60 5.0 5.0 ALS QCS3 requirement ✓ EG035T 3 60 Total Mercury by FIMS 5.0 5.0 1 ALS QCS3 requirement 3 Total Metals by ICP-MS - Suite A EG020A-T 60 5.0 5.0 ALS QCS3 requirement

ALS QCS3 requirement

Page : 11 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Brief Method Summaries

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the US EPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request. The following report provides brief descriptions of the analytical procedures employed for results reported in the Certificate of Analysis. Sources from which ALS methods have been developed are provided within the Method Descriptions.

| Analytical Methods | Method | Matrix | Method Descriptions |
|---|-------------|--------|---|
| Suspended Solids (High Level) | EA025H | WATER | APHA 21st ed., 2540D A gravimetric procedure employed to determine the amount of `non-filterable` residue in a aqueous sample. The prescribed GFC (1.2um) filter is rinsed with deionised water, oven dried and weighed prior to analysis. A well-mixed sample is filtered through a glass fibre filter (1.2um). The residue on the filter paper is dried at 104+/-2C. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Total Metals by ICP-MS - Suite A | EG020A-T | WATER | (APHA 21st ed., 3125; USEPA SW846 - 6020, ALS QWI-EN/EG020): The ICPMS technique utilizes a highly efficient argon plasma to ionize selected elements. Ions are then passed into a high vacuum mass spectrometer, which separates the analytes based on their distinct mass to charge ratios prior to their measurement by a discrete dynode ion detector. |
| Total Mercury by FIMS | EG035T | WATER | AS 3550, APHA 21st ed. 3112 Hg - B (Flow-injection (SnCl2)(Cold Vapour generation) AAS) FIM-AAS is an automated flameless atomic absorption technique. A bromate/bromide reagent is used to oxidise any organic mercury compounds in the unfiltered sample. The ionic mercury is reduced online to atomic mercury vapour by SnCl2 which is then purged into a heated quartz cell. Quantification is by comparing absorbance against a calibration curve. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Ammonia as N by Discrete analyser | EK055G | WATER | APHA 21st ed., 4500-NH3 G Ammonia is determined by direct colorimetry by Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Nitrite as N by Discrete Analyser | EK057G | WATER | APHA 21st ed., 4500-NO2- B. Nitrite is determined by direct colourimetry by Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Nitrate as N by Discrete Analyser | EK058G | WATER | APHA 21st ed., 4500-NO3- F. Nitrate is reduced to nitrite by way of a cadmium reduction column followed by quantification by Discrete Analyser. Nitrite is determined seperately by direct colourimetry and result for Nitrate calculated as the difference between the two results. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Nitrite and Nitrate as N (NOx) by Discrete Analyser | EK059G | WATER | APHA 21st ed., 4500-NO3- F. Combined oxidised Nitrogen (NO2+NO3) is determined by Cadmium Reduction and direct colourimetry by Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Total Kjeldahl Nitrogen as N By Discrete Analyser | EK061G | WATER | APHA 21st ed., 4500-Norg D. 25mL water samples are digested using a traditional Kjeldahl digestion followed by determination by Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Total Nitrogen as N (TKN + Nox) By Discrete Analyser | EK062G | WATER | APHA 21st ed., 4500-Norg / 4500-NO3 This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Total Phosphorus as P By Discrete Analyser | EK067G | WATER | APHA 21st ed., 4500-P B&F This procedure involves sulphuric acid digestion of a 100mL sample to break phosphorus down to orthophosphate. The orthophosphate reacts with ammonium molybdate and antimony potassium tartrate to form a complex which is then reduced and its concentration measured at 880nm using Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Preparation Methods | Method | Matrix | Method Descriptions |
| TKN/TP Digestion | EK061/EK067 | WATER | APHA 21st ed., 4500 Norg - D; APHA 21st ed., 4500 P - H. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Digestion for Total Recoverable Metals | EN25 | WATER | USEPA SW846-3005 Method 3005 is a Nitric/Hydrochloric acid digestion procedure used to prepare surface and ground water samples for analysis by ICPAES or ICPMS. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |

Page : 12 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Summary of Outliers

Outliers: Quality Control Samples

The following report highlights outliers flagged in the Quality Control (QC) Report. Surrogate recovery limits are static and based on USEPA SW846 or ALS-QWI/EN/38 (in the absence of specific USEPA limits). This report displays QC Outliers (breaches) only.

Duplicates, Method Blanks, Laboratory Control Samples and Matrix Spikes

- For all matrices, no Method Blank value outliers occur.
- For all matrices, no Duplicate outliers occur.
- For all matrices, no Laboratory Control outliers occur.
- For all matrices, no Matrix Spike outliers occur.

Regular Sample Surrogates

• For all regular sample matrices, no surrogate recovery outliers occur.

Outliers: Analysis Holding Time Compliance

This report displays Holding Time breaches only. Only the respective Extraction / Preparation and/or Analysis component is/are displayed.

Matrix: WATER

| Method | | Ex | traction / Preparation | | | Analysis | |
|---------------------------------|------|----------------|------------------------|---------|---------------|------------------|---------|
| Container / Client Sample ID(s) | | Date extracted | Due for extraction | Days | Date analysed | Due for analysis | Days |
| | | | | overdue | | | overdue |
| EA025: Suspended Solids | | | | | | | |
| Clear Plastic Bottle - Natural | | | | | | | |
| A1S, | B1S, | | | | 28-DEC-2011 | 26-DEC-2011 | 2 |
| C1S, | D1S, | | | | | | |
| A2S, | B2S, | | | | | | |
| C2S, | D2S, | | | | | | |
| A3S, | B3S, | | | | | | |
| C3S, | D3S, | | | | | | |
| B4S, | C4S, | | | | | | |
| D4S, | X4S, | | | | | | |
| Y4S, | AaS, | | | | | | |
| BbS, | CcS, | | | | | | |
| DdS, | A5S, | | | | | | |
| B5S, | C5S, | | | | | | |
| D5S | | | | | | | |

Page : 13 of 13

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Matrix: WATER

| Method | | E | extraction / Preparation | | Analysis | | | |
|---------------------------------------|---|----------------|--------------------------|-----------------|---------------|------------------|-----------------|--|
| Container / Client Sample ID(s) | | Date extracted | Due for extraction | Days overdue | Date analysed | Due for analysis | Days overdue | |
| EK057G: Nitrite as N by Discrete Anal | yser - Analysis Holding Time Compliance | | | | | | | |
| Clear Plastic Bottle - Natural | | | | | | | | |
| A1S, | B1S, | | | | 22-DEC-2011 | 21-DEC-2011 | 1 | |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Outliers : Frequency of Quality Control Samples

The following report highlights breaches in the Frequency of Quality Control Samples.

No Quality Control Sample Frequency Outliers exist.

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

QUALITY CONTROL REPORT

Work Order : **EM1114567** Page : 1 of 12

Amendment : 1

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

 Telephone
 : +61 08 8424 3800
 Telephone
 : +61-3-8549 9608

 Facsimile
 : +61 08 8424 3810
 Facsimile
 : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

Site : ----

 C-O-C number
 : -- Date Samples Received
 : 22-DEC-2011

 Sampler
 : GB
 Issue Date
 : 28-FEB-2012

No. of samples received : 60

Quote number : ADBQ/037/11 No. of samples analysed : 60

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for release.

This Quality Control Report contains the following information:

: ----

- Laboratory Duplicate (DUP) Report; Relative Percentage Difference (RPD) and Acceptance Limits
- Method Blank (MB) and Laboratory Control Spike (LCS) Report; Recovery and Acceptance Limits
- Matrix Spike (MS) Report; Recovery and Acceptance Limits



Order number

NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

| Signatories | Position | Accreditation Category |
|----------------|------------------------|------------------------|
| Eric Chau | Metals Team Leader | Melbourne Inorganics |
| Herman Lin | Laboratory Coordinator | Melbourne Inorganics |
| Varsha Ho Wing | Non-Metals Team Leader | Melbourne Inorganics |

Page : 2 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

Key: Anonymous = Refers to samples which are not specifically part of this work order but formed part of the QC process lot

CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

RPD = Relative Percentage Difference

= Indicates failed QC

Page : 3 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Laboratory Duplicate (DUP) Report

The quality control term Laboratory Duplicate refers to a randomly selected intralaboratory split. Laboratory duplicates provide information regarding method precision and sample heterogeneity. The permitted ranges for the Relative Percent Deviation (RPD) of Laboratory Duplicates are specified in ALS Method QWI-EN/38 and are dependent on the magnitude of results in comparison to the level of reporting: Result < 10 times LOR:-No Limit; Result between 10 and 20 times LOR:-0% - 50%; Result > 20 times LOR:-0% - 20%.

| Sub-Matrix: WATER | | | | Laboratory Duplicate (DUP) Report | | | | | | | | |
|----------------------|--------------------------|-------------------------------|------------|-----------------------------------|------|-----------------|------------------|---------|---------------------|--|--|--|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) | | | |
| EA025: Suspended | Solids (QC Lot: 2105373) |) | | | | | | | | | | |
| EM1114567-027 | A6S | EA025H: Suspended Solids (SS) | | 5 | mg/L | 223 | 240 | 7.3 | 0% - 20% | | | |
| EM1114567-036 | B2B | EA025H: Suspended Solids (SS) | | 5 | mg/L | 129 | 130 | 0.8 | 0% - 20% | | | |
| EA025: Suspended | Solids (QC Lot: 2105374) |) | | | | | | | | | | |
| EM1114567-048 | Y4B | EA025H: Suspended Solids (SS) | | 5 | mg/L | 128 | 128 | 0.0 | 0% - 20% | | | |
| EM1114567-057 | A6B | EA025H: Suspended Solids (SS) | | 5 | mg/L | 111 | 114 | 2.7 | 0% - 20% | | | |
| EA025: Suspended | Solids (QC Lot: 2107718) |) | | | | | | | | | | |
| EM1114567-001 | A1S | EA025H: Suspended Solids (SS) | | 5 | mg/L | 50 | 69 | 31.9 | 0% - 50% | | | |
| EM1114567-010 | B3S | EA025H: Suspended Solids (SS) | | 5 | mg/L | 60 | 63 | 4.9 | 0% - 50% | | | |
| EA025: Suspended | Solids (QC Lot: 2107719) | | | | | | | | | | | |
| EM1114567-022 | DdS | EA025H: Suspended Solids (SS) | | 5 | mg/L | 62 | 66 | 6.2 | 0% - 50% | | | |
| EM1114604-001 | Anonymous | EA025H: Suspended Solids (SS) | | 5 | mg/L | 835 | 865 | 3.5 | 0% - 20% | | | |
| FG020T: Total Meta | ls by ICP-MS (QC Lot: 21 | | | | | | | | | | | |
| EM1114567-001 | A1S | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.0 | No Limit | | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.008 | 0.009 | 21.7 | No Limit | | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.001 | 0.002 | 0.0 | No Limit | | | |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.005 | 0.0 | No Limit | | | |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | | |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.0 | No Limit | | | |
| EM1114567-010 | B3S | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.0 | No Limit | | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.009 | 0.0 | No Limit | | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | | |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.003 | 0.001 | 64.5 | No Limit | | | |

Page : 4 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | | Laboratory Duplicate (DUP) Report | | | | | | | |
|----------------------|----------------------|----------------------|------------|-----------------------------------|------|-----------------|------------------|---------|---------------------|--|--|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) | | |
| EG020T: Total Meta | s by ICP-MS (QC Lot: | 2105258) - continued | | | | | | | | | |
| EM1114567-010 | B3S | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.0 | No Limit | | |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.0 | No Limit | | |
| EG020T: Total Meta | s by ICP-MS (QC Lot: | 2105259) | | | | | | | | | |
| EM1114567-021 | CcS | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.0 | No Limit | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.0 | 0% - 50% | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.0 | No Limit | | |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.0 | No Limit | | |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.0 | No Limit | | |
| EM1114567-030 | D6S | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.0 | No Limit | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.0 | 0% - 50% | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.0 | No Limit | | |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.0 | No Limit | | |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.0 | No Limit | | |
| EG020T: Total Metal | s by ICP-MS (QC Lot: | 2107796) | | | | | | | | | |
| EM1114566-001 | Anonymous | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | 0.0001 | 0.0001 | 0.0 | No Limit | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | 0.013 | 0.014 | 11.3 | 0% - 50% | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.058 | 0.061 | 5.0 | 0% - 20% | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | 0.002 | 0.002 | 0.0 | No Limit | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | 0.001 | 0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.017 | 0.017 | 0.0 | 0% - 50% | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | 0.015 | 0.016 | 7.4 | 0% - 50% | | |

Page : 5 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | | Laboratory Duplicate (DUP) Report | | | | | | | |
|----------------------|-----------------------------|----------------------|------------|-----------------------------------|------|-----------------|------------------|---------|---------------------|--|--|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) | | |
| EG020T: Total Metal | s by ICP-MS (QC Lot: 2 | 107796) - continued | | | | | | | | | |
| EM1114566-001 | Anonymous | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | 0.082 | 0.087 | 5.1 | 0% - 20% | | |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.016 | 0.017 | 0.0 | 0% - 50% | | |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | 0.163 | 0.187 | 13.7 | 0% - 20% | | |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | 0.06 | 0.06 | 0.0 | No Limit | | |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | 1.67 | 1.71 | 2.4 | 0% - 20% | | |
| EM1114567-050 | BbB | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.0001 | 0.0 | No Limit | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.005 | 0.006 | 0.0 | No Limit | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.0 | 0% - 50% | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.005 | 0.005 | 0.0 | No Limit | | |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.009 | 32.8 | No Limit | | |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | 0.06 | 0.10 | 42.2 | No Limit | | |
| EG035T: Total Reco | verable Mercury by FIM | S (QC Lot: 2115265) | | | | | | | | | |
| EM1114567-001 | A1S | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| EM1114567-010 | B3S | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | 0.0007 | 151 | No Limit | | |
| EG035T: Total Reco | verable Mercury by FIM | S (QC Lot: 2115266) | | | | | | | | | |
| EM1114567-021 | CcS | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| EM1114567-030 | D6S | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| FG035T: Total Reco | verable Mercury by FIM | S (QC Lot: 2115267) | | | | | | | | | |
| EM1114567-041 | C3B | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| EM1114567-050 | BbB | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| | s N by Discrete Analyse | | | | 3 | | | | | | |
| EM1114567-001 | A1S | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.10 | 0.0 | 0% - 50% | | |
| EM1114567-010 | B3S | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.11 | 10.1 | 0% - 50% | | |
| | | | 7004 41 7 | 0.01 | mg/L | 0.12 | 0.11 | 10.1 | 070 0070 | | |
| EM1114567-021 | S N by Discrete Analyse CcS | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.11 | 0.10 | 0.0 | 0% - 50% | | |
| EM1114567-021 | D6S | | 7664-41-7 | 0.01 | mg/L | 0.11 | 0.09 | 0.0 | No Limit | | |
| | | EK055G: Ammonia as N | 7004-41-7 | 0.01 | my/L | 0.10 | 0.09 | 0.0 | INO LIITIIL | | |
| | s N by Discrete Analyse | | 7004 44 7 | 0.04 | | 0.40 | 0.45 | 40.0 | 00/ 500/ | | |
| EM1114567-041 | C3B | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.15 | 16.8 | 0% - 50% | | |
| EM1114567-050 | BbB | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.11 | 0.13 | 16.8 | 0% - 50% | | |
| | by Discrete Analyser | • | | | - | | | | | | |
| EM1114567-001 | A1S | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| EM1114567-010 | B3S | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |

Page : 6 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | | | | | | | |
|----------------------|----------------------------------|--------------------------------------|------------|------|------|-----------------|------------------|---------|---------------------|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) |
| EK057G: Nitrite as | N by Discrete Analyser | (QC Lot: 2105294) | | | | | | | |
| EM1114567-021 | CcS | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-030 | D6S | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EK057G: Nitrite as | N by Discrete Analyser | (QC Lot: 2105295) | | | | | | | |
| EM1114567-041 | C3B | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-050 | BbB | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EK059G: Nitrite plu | s Nitrate as N (NOx) by | Discrete Analyser (QC Lot: 2115293) | | | | | | | |
| EM1114567-001 | A1S | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-010 | B3S | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EK059G: Nitrite plu | s Nitrate as N (NOx) by | Discrete Analyser (QC Lot: 2115296) | | | | | | | |
| EM1114567-021 | CcS | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-030 | D6S | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EK059G: Nitrite plu | s Nitrate as N (NOx) by | Discrete Analyser (QC Lot: 2115298) | | | | | | | |
| EM1114567-041 | СЗВ | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-050 | BbB | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EK061G: Total Kjelo | dahl Nitrogen By Discrete | e Analyser (QC Lot: 2109542) | | | | | | | |
| EM1114567-001 | A1S | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.3 | 0.0 | No Limit |
| EM1114567-010 | B3S | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.2 | 0.0 | No Limit |
| EK061G: Total Kjelo | dahl Nitrogen By Discrete | e Analyser (QC Lot: 2109544) | | | | | | | |
| EM1114567-021 | CcS | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.1 | 0.0 | No Limit |
| EM1114567-030 | D6S | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.3 | 0.2 | 52.8 | No Limit |
| EK061G: Total Kjelo | dahl Nitrogen By Discrete | e Analyser (QC Lot: 2110810) | | | | | | | |
| EM1114567-041 | СЗВ | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | 0.2 | 0.0 | No Limit |
| EM1114567-050 | BbB | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | 0.0 | No Limit |
| EK067G: Total Phos | sphorus as P by Discrete | e Analyser (QC Lot: 2109543) | | | | | | | |
| EM1114567-001 | A1S | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-010 | B3S | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.01 | 0.0 | No Limit |
| EK067G: Total Phos | sphorus as P by Di <u>screte</u> | e Analyser (QC Lot: 2109545) | | | | | | | |
| EM1114567-021 | CcS | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-030 | D6S | EK067G: Total Phosphorus as P | | 0.01 | mg/L | 0.05 | 0.04 | 29.9 | No Limit |
| EK067G: Total Phos | sphorus as P by Di <u>screte</u> | Analyser (QC Lot: 2110811) | | | | | | | |
| EM1114567-041 | C3B | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.01 | 0.0 | No Limit |
| EM1114567-050 | BbB | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |

Page : 7 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Method Blank (MB) and Laboratory Control Spike (LCS) Report

The quality control term Method / Laboratory Blank refers to an analyte free matrix to which all reagents are added in the same volumes or proportions as used in standard sample preparation. The purpose of this QC parameter is to monitor potential laboratory contamination. The quality control term Laboratory Control Sample (LCS) refers to a certified reference material, or a known interference free matrix spiked with target analytes. The purpose of this QC parameter is to monitor method precision and accuracy independent of sample matrix. Dynamic Recovery Limits are based on statistical evaluation of processed LCS.

| Sub-Matrix: WATER | | | | Method Blank (MB) | Laboratory Control Spike (LCS) Report | | | | | |
|--|------------|--------|------|-------------------|---------------------------------------|--------------------|-----|------------|--|--|
| | | | | Report | Spike | Spike Recovery (%) | | Limits (%) | | |
| Method: Compound | CAS Number | LOR | Unit | Result | Concentration | LCS | Low | High | | |
| EA025: Suspended Solids (QCLot: 2105373) | | | | | | | | | | |
| EA025H: Suspended Solids (SS) | | 5 | mg/L | <5 | 150 mg/L | 101 | 96 | 112 | | |
| EA025: Suspended Solids (QCLot: 2105374) | | | | | | ' | | | | |
| EA025H: Suspended Solids (SS) | | 5 | mg/L | <5 | 150 mg/L | 101 | 96 | 112 | | |
| EA025: Suspended Solids (QCLot: 2107718) | | | | | | | | | | |
| EA025H: Suspended Solids (SS) | | 5 | mg/L | <5 | 150 mg/L | 101 | 96 | 112 | | |
| EA025: Suspended Solids (QCLot: 2107719) | | | 3 | | | | | | | |
| EA025. Suspended Solids (QCL0t. 2107719) EA025H: Suspended Solids (SS) | | 5 | mg/L | <5 | 150 mg/L | 105 | 96 | 112 | | |
| | | | mg/L | .0 | 100 mg/L | 100 | | 112 | | |
| EG020T: Total Metals by ICP-MS (QCLot: 2105258) | 7440-38-2 | 0.001 | ma/l | <0.001 | 0.1 mg/l | 106 | 86 | 110 | | |
| EG020A-T: Arsenic | 7440-36-2 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 68 | 128 | | |
| EG020A-T: Beryllium | | | mg/L | | 0.1 mg/L | | | | | |
| EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 87 | 115 | | |
| EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.1 mg/L | 102 | 87 | 111 | | |
| EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 103 | 87 | 115 | | |
| EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 101 | 88 | 114 | | |
| EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 88 | 110 | | |
| EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 102 | 90 | 114 | | |
| EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 87 | 113 | | |
| EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 107 | 89 | 113 | | |
| EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | 0.1 mg/L | 103 | 86 | 114 | | |
| EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | 0.1 mg/L | 104 | 82 | 116 | | |
| EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.5 mg/L | 101 | 81 | 125 | | |
| EG020T: Total Metals by ICP-MS (QCLot: 2105259) | | | | | | | | | | |
| EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 102 | 86 | 110 | | |
| EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 124 | 68 | 128 | | |
| EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 101 | 87 | 115 | | |
| EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.1 mg/L | 98.9 | 87 | 111 | | |
| EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 87 | 115 | | |
| EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 102 | 88 | 114 | | |
| EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 102 | 88 | 110 | | |
| EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 99.2 | 90 | 114 | | |
| EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 100 | 87 | 113 | | |
| EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 105 | 89 | 113 | | |
| EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | 0.1 mg/L | 108 | 86 | 114 | | |
| EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | 0.1 mg/L | 101 | 82 | 116 | | |

Page : 8 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | | Method Blank (MB) | Laboratory Control Spike (LCS) Report | | | | |
|---|----------------|----------|--------|-------------------|---------------------------------------|--------------------|----------|------------|--|
| | | | | Report | Spike | Spike Recovery (%) | Recovery | Limits (%) | |
| Method: Compound | CAS Number | LOR | Unit | Result | Concentration | LCS | Low | High | |
| G020T: Total Metals by ICP-MS (QCLot: 2105259) - contin | ued | | | | | | | | |
| G020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.5 mg/L | 112 | 81 | 125 | |
| G020T: Total Metals by ICP-MS (QCLot: 2107796) | | | | | | | | | |
| G020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 110 | 86 | 110 | |
| G020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 105 | 68 | 128 | |
| G020A-T: Barium | 7440-39-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 110 | 87 | 115 | |
| G020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.1 mg/L | 103 | 87 | 111 | |
| G020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 87 | 115 | |
| G020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 107 | 88 | 114 | |
| G020A-T: Copper | 7440-50-8 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 108 | 88 | 110 | |
| G020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 107 | 90 | 114 | |
| G020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 108 | 87 | 113 | |
| G020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 109 | 89 | 113 | |
| G020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | 0.1 mg/L | 103 | 86 | 114 | |
| G020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | 0.1 mg/L | 110 | 82 | 116 | |
| G020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.5 mg/L | 108 | 81 | 125 | |
| G035T: Total Recoverable Mercury by FIMS (QCLot: 2115 | | | | | | | | | |
| G035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | 0.0100 mg/L | 89.2 | 69 | 125 | |
| G035T: Total Recoverable Mercury by FIMS (QCLot: 2115 | 266) | | | | | | | | |
| G035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | 0.0100 mg/L | 85.6 | 69 | 125 | |
| G035T: Total Recoverable Mercury by FIMS (QCLot: 2115 | 267) | | | | | | | | |
| G035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | 0.0100 mg/L | 80.5 | 69 | 125 | |
| K055G: Ammonia as N by Discrete Analyser (QCLot: 2115 | 294) | | | | | | | | |
| :K055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | <0.01 | 0.5 mg/L | 113 | 76 | 122 | |
| K055G: Ammonia as N by Discrete Analyser (QCLot: 2115 | 297) | | | | | | | | |
| K055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | <0.01 | 0.5 mg/L | 114 | 76 | 122 | |
| :K055G: Ammonia as N by Discrete Analyser (QCLot: 2115 | 299) | | | | _ | | | | |
| K055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | <0.01 | 0.5 mg/L | 110 | 76 | 122 | |
| K057G: Nitrite as N by Discrete Analyser (QCLot: 210529; | | | 3 | | J. J | | | | |
| :R057G: Nitrite as N by Discrete Analyser (QCL0t. 210525) | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 100 | 84 | 112 | |
| | | 0.01 | ilig/L | 10.01 | 0.0 mg/L | 100 | 04 | 112 | |
| K057G: Nitrite as N by Discrete Analyser (QCLot: 2105294 | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 99.1 | 84 | 112 | |
| K057G: Nitrite as N | | 0.01 | mg/L | ~ 0.01 | 0.5 mg/L | 33. I | 04 | 112 | |
| K057G: Nitrite as N by Discrete Analyser (QCLot: 210529 | | 0.04 | - n | 40.04 | 0.5 " | 00.7 | 0.4 | 110 | |
| K057G: Nitrite as N | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 99.7 | 84 | 112 | |
| K059G: Nitrite plus Nitrate as N (NOx) by Discrete Analys | er (QCLot: 211 | | | | | | | | |
| K059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 95.0 | 73 | 127 | |
| K059G: Nitrite plus Nitrate as N (NOx) by Discrete Analys | er (QCLot: 211 | <u> </u> | | | | | | | |
| K059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 104 | 73 | 127 | |

Page : 9 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | | | Method Blank (MB) | | Laboratory Control Spike (LC | S) Report | |
|---|---------------------------|-------|------|-------------------|---------------|------------------------------|-----------|------------|
| | | | | Report | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| Method: Compound | CAS Number | LOR | Unit | Result | Concentration | LCS | Low | High |
| EK059G: Nitrite plus Nitrate as N (NOx) by Disc | rete Analyser (QCLot: 211 | 5298) | | | | | | |
| EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 101 | 73 | 127 |
| EK061G: Total Kjeldahl Nitrogen By Discrete Ana | alyser (QCLot: 2109542) | | | | | | | |
| EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | 10 mg/L | 104 | 63 | 117 |
| EK061G: Total Kjeldahl Nitrogen By Discrete Ana | alyser (QCLot: 2109544) | | | | | | | |
| EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | 10 mg/L | 106 | 63 | 117 |
| EK061G: Total Kjeldahl Nitrogen By Discrete Ana | alyser (QCLot: 2110810) | | | | | | | |
| EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | 10 mg/L | 87.1 | 63 | 117 |
| EK067G: Total Phosphorus as P by Discrete Ana | lyser (QCLot: 2109543) | | | | | | | |
| EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 4.42 mg/L | 99.5 | 73 | 117 |
| EK067G: Total Phosphorus as P by Discrete Ana | lyser (QCLot: 2109545) | | | | | | | |
| EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 4.42 mg/L | 89.7 | 73 | 117 |
| EK067G: Total Phosphorus as P by Discrete Ana | llyser (QCLot: 2110811) | | | | | | | |
| EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 4.42 mg/L | 98.0 | 73 | 117 |

Page : 10 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Matrix Spike (MS) Report

The quality control term Matrix Spike (MS) refers to an intralaboratory split sample spiked with a representative set of target analytes. The purpose of this QC parameter is to monitor potential matrix effects on analyte recoveries. Static Recovery Limits as per laboratory Data Quality Objectives (DQOs). Ideal recovery ranges stated may be waived in the event of sample matrix interference.

| ub-Matrix: WATER | | | | | Matrix Spike (MS) Repo | ort | |
|---------------------|------------------------------|---------------------------------------|------------|---------------|------------------------|----------|------------|
| | | | | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| aboratory sample ID | Client sample ID | Method: Compound | CAS Number | Concentration | MS | Low | High |
| G020T: Total Metal | s by ICP-MS (QCLot: 2105258) | | | | | | |
| M1114567-001 | A1S | EG020A-T: Arsenic | 7440-38-2 | 1 mg/L | 110 | 72 | 146 |
| | | EG020A-T: Beryllium | 7440-41-7 | 1 mg/L | 85.8 | 61 | 139 |
| | | EG020A-T: Barium | 7440-39-3 | 1 mg/L | 98.8 | 78 | 126 |
| | | EG020A-T: Cadmium | 7440-43-9 | 0.25 mg/L | 83.3 | 73 | 131 |
| | | EG020A-T: Chromium | 7440-47-3 | 1 mg/L | 85.1 | 65 | 131 |
| | | EG020A-T: Cobalt | 7440-48-4 | 1 mg/L | 101 | 68 | 132 |
| | | EG020A-T: Copper | 7440-50-8 | 1 mg/L | 84.6 | 71 | 125 |
| | | EG020A-T: Lead | 7439-92-1 | 1 mg/L | 95.5 | 68 | 130 |
| | | EG020A-T: Manganese | 7439-96-5 | 1 mg/L | 88.8 | 63 | 129 |
| | | EG020A-T: Nickel | 7440-02-0 | 1 mg/L | 89.5 | 72 | 128 |
| | | EG020A-T: Vanadium | 7440-62-2 | 1 mg/L | 95.6 | 66 | 130 |
| | | EG020A-T: Zinc | 7440-66-6 | 1 mg/L | 79.4 | 67 | 129 |
| G020T: Total Metal | s by ICP-MS (QCLot: 2105259) | | | - | | | |
| M1114567-021 | CcS | EG020A-T: Arsenic | 7440-38-2 | 1 mg/L | 104 | 72 | 146 |
| | 57-021 CCS | EG020A-T: Alsenic | 7440-41-7 | 1 mg/L | 94.9 | 61 | 139 |
| | | EG020A-T: Berymum | 7440-39-3 | 1 mg/L | 109 | 78 | 126 |
| | | EG020A-T: Cadmium | 7440-43-9 | 0.25 mg/L | 93.9 | 73 | 131 |
| | | EG020A-T: Cadmidiff | 7440-47-3 | 1 mg/L | 105 | 65 | 131 |
| | | EG020A-T: Coholit | 7440-48-4 | 1 mg/L | 111 | 68 | 132 |
| | | EG020A-T: Copper | 7440-50-8 | 1 mg/L | 93.9 | 71 | 125 |
| | | EG020A-T: Copper | 7439-92-1 | 1 mg/L | 105 | 68 | 130 |
| | | EG020A-T: Lead EG020A-T: Manganese | 7439-96-5 | 1 mg/L | 99.6 | 63 | 129 |
| | | EG020A-T: Nickel | 7440-02-0 | 1 mg/L | 101 | 72 | 128 |
| | | EG020A-T: Vanadium | 7440-62-2 | 1 mg/L | 110 | 66 | 130 |
| | | EG020A-T: Variation | 7440-66-6 | 1 mg/L | 88.4 | 67 | 129 |
| C020T: Total Matel | s by ICP-MS (QCLot: 2107796) | LOUZUA-1. ZIIIC | 111000 | · ···g·= | 33.1 | <u> </u> | |
| M1114567-041 | C3B | FC0000A T. Assertia | 7440-38-2 | 1 mg/L | 117 | 72 | 146 |
| WITT14307-041 | C3B | EG020A-T. Providence | 7440-41-7 | 1 mg/L | 94.9 | 61 | 139 |
| | | EG020A-T. Beryllium | 7440-39-3 | 1 mg/L | 120 | 78 | 126 |
| | | EG020A-T: Barium | 7440-43-9 | 0.25 mg/L | 104 | 73 | 131 |
| | | EG020A-T. Clauseiner | 7440-43-9 | | 112 | 65 | 131 |
| | | EG020A-T: Chromium | 7440-47-3 | 1 mg/L | 125 | | 131 |
| | | EG020A-T. Cobalt | 7440-48-4 | 1 mg/L | 125 | 68 71 | 132 |
| | | EG020A-T: Copper | | 1 mg/L | | | |
| | | EG020A-T: Lead | 7439-92-1 | 1 mg/L | 112 | 68 | 130 |
| | | EG020A-T: Manganese | 7439-96-5 | 1 mg/L | 108 | 63 | 129 |

Page : 11 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | | | | Matrix Spike (MS) Report | | | |
|---------------------------------------|--------------------------------------|--------------------------------------|-------------|---------------|--------------------------|----------|------------|--|
| | | | | Spike | Spike Recovery (%) | Recovery | Limits (%) | |
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | Concentration | MS | Low | High | |
| G020T: Total Metal | s by ICP-MS (QCLot: 2107796) | - continued | | | | | | |
| EM1114567-041 | C3B | EG020A-T: Nickel | 7440-02-0 | 1 mg/L | 110 | 72 | 128 | |
| | | EG020A-T: Vanadium | 7440-62-2 | 1 mg/L | 118 | 66 | 130 | |
| | | EG020A-T: Zinc | 7440-66-6 | 1 mg/L | 99.9 | 67 | 129 | |
| G035T: Total Reco | verable Mercury by FIMS (QC | Lot: 2115265) | | | | | | |
| EM1114567-002 | B1S | EG035T: Mercury | 7439-97-6 | 0.0100 mg/L | 84.8 | 70 | 130 | |
| G035T: Total Reco | verable Mercury by FIMS (QC | Lot: 2115266) | | | | | | |
| EM1114567-022 | DdS | EG035T: Mercury | 7439-97-6 | 0.0100 mg/L | 84.9 | 70 | 130 | |
| G035T: Total Reco | verable Mercury by FIMS (QC | Lot: 2115267) | | | | | | |
| EM1114567-042 | D3B | EG035T: Mercury | 7439-97-6 | 0.0100 mg/L | 82.0 | 70 | 130 | |
| K055G: Ammonia a | s N by Discrete Analyser (QC | | | | | | | |
| EM1114567-002 | B1S | EK055G: Ammonia as N | 7664-41-7 | 0.5 mg/L | 94.4 | 70 | 130 | |
| K055G: Ammonia | s N by Discrete Analyser (QC | | | | | | | |
| EM1114567-022 | DdS | EK055G: Ammonia as N | 7664-41-7 | 0.5 mg/L | 98.3 | 70 | 130 | |
| | is N by Discrete Analyser (QC | | | | 77.7 | | | |
| EM1114567-042 | D3B | EK055G: Ammonia as N | 7664-41-7 | 0.5 mg/L | 90.3 | 70 | 130 | |
| | N by Discrete Analyser (QCLo | | 7.00.7.17.1 | 0.0g/_ | 55.5 | | | |
| EM1114567-002 | B1S | EK057G: Nitrite as N | | 0.5 mg/L | 118 | 70 | 130 | |
| | N by Discrete Analyser (QCLo | | | 0.0 mg/L | 110 | 70 | 100 | |
| EM1114567-022 | DdS | EK057G: Nitrite as N | | 0.5 mg/L | 118 | 70 | 130 | |
| | | | | 0.5 Hig/L | 110 | 70 | 130 | |
| :K057G: Nitrite as I EM1114567-042 | N by Discrete Analyser (QCLor D3B | | | 0.5/ | 115 | 70 | 400 | |
| | | EK057G: Nitrite as N | | 0.5 mg/L | 115 | 70 | 130 | |
| | Nitrate as N (NOx) by Discre | | | 0.5 " | 440 | | 100 | |
| EM1114567-002 | B1S | EK059G: Nitrite + Nitrate as N | | 0.5 mg/L | 119 | 70 | 130 | |
| | Nitrate as N (NOx) by Discre | te Analyser (QCLot: 2115296) | | | | | | |
| EM1114567-022 | DdS | EK059G: Nitrite + Nitrate as N | | 0.6 mg/L | 110 | 70 | 130 | |
| | s Nitrate as N (NOx) by Discre | te Analyser (QCLot: 2115298) | | | | | | |
| EM1114567-042 | D3B | EK059G: Nitrite + Nitrate as N | | 0.5 mg/L | 117 | 70 | 130 | |
| K061G: Total Kjeld | ahl Nitrogen By Discrete Analy | vser (QCLot: 2109542) | | | | | | |
| EM1114567-002 | B1S | EK061G: Total Kjeldahl Nitrogen as N | | 25 mg/L | 107 | 70 | 130 | |
| K061G: Total Kjeld | ahl Nitrogen By Discrete Analy | vser (QCLot: 2109544) | | | | | | |
| M1114567-022 | DdS | EK061G: Total Kjeldahl Nitrogen as N | | 25 mg/L | 76.7 | 70 | 130 | |
| K061G: Total Kjeld | ahl Nitrogen By Discrete Analy | vser (QCLot: 2110810) | | | | | | |
| EM1114567-042 | D3B | EK061G: Total Kjeldahl Nitrogen as N | | 25 mg/L | 102 | 70 | 130 | |
| K067G: Total Phos | phorus as P by Discrete Analy | | | | | | | |
| EM1114567-002 | B1S | EK067G: Total Phosphorus as P | | 5 mg/L | 79.7 | 70 | 130 | |

Page : 12 of 12

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | | | | Matrix Spike (MS) Repo | ort | |
|----------------------|--|-------------------------------|------------|---------------|------------------------|----------|------------|
| | | | | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | Concentration | MS | Low | High |
| EK067G: Total Phos | phorus as P by Discrete Analyser (QCLot: 2 | 2109545) | | | | | |
| EM1114567-022 | DdS | EK067G: Total Phosphorus as P | | 5 mg/L | 88.0 | 70 | 130 |
| EK067G: Total Phos | phorus as P by Discrete Analyser (QCLot: : | 2110811) | | | | | |
| EM1114567-042 | D3B | EK067G: Total Phosphorus as P | | 5 mg/L | 95.2 | 70 | 130 |

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

CERTIFICATE OF ANALYSIS

Work Order : **EM1114567** Page : 1 of 14

Amendment : 1

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

 Telephone
 : +61 08 8424 3800
 Telephone
 : +61-3-8549 9608

 Facsimile
 : +61 08 8424 3810
 Facsimile
 : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

Order number : ----

 C-O-C number
 : -- Date Samples Received
 : 22-DEC-2011

 Sampler
 : GB
 Issue Date
 : 28-FEB-2012

Site : ----

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for release.

This Certificate of Analysis contains the following information:

- General Comments
- Analytical Results



NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

| Signatories | Position | Accreditation Category |
|----------------|------------------------|------------------------|
| Eric Chau | Metals Team Leader | Melbourne Inorganics |
| Herman Lin | Laboratory Coordinator | Melbourne Inorganics |
| Varsha Ho Wing | Non-Metals Team Leader | Melbourne Inorganics |

Page : 2 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

When sampling time information is not provided by the client, sampling dates are shown without a time component. In these instances, the time component has been assumed by the laboratory for processing purposes.

Key: CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

^ = This result is computed from individual analyte detections at or above the level of reporting

- 27/2/12 This report has been amended and re-released to allow the reporting of additional analytical data.
- EA025: Total Suspended Solids has been conducted outside of holding times due to laboratory constraints. Results should be scrutinised accordingly.
- It is recognised that Total Kjeldahl Nitrogen is less than Ammonia for sample EM1114567. However, the difference is within experimental variation of the methods.

Page : 3 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | A1S | B1S | C1S | D1S | A2S |
|---------------------------------|--------------------------|------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Clie | ent sampli | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-001 | EM1114567-002 | EM1114567-003 | EM1114567-004 | EM1114567-005 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 50 | 85 | 69 | 31 | 52 |
| EG020T: Total Metals by ICP-MS | ; | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.0001 | 0.0001 | <0.0001 | 0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.008 | 0.020 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.001 | 0.002 | 0.003 | 0.002 | 0.002 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.022 | <0.005 | <0.005 | 0.006 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mei | cury by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Disc | rete Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.11 | 0.10 | 0.09 | 0.10 |
| EK057G: Nitrite as N by Discret | e Analyser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discre | te Analyser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as | | | 3 | | | | | |
| Nitrite + Nitrate as N | N (NOX) by Discrete Anal | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitroger | | 0.01 | mg/L | 70.01 | 10.01 | -0.01 | 10.01 | -0.01 |
| Total Kjeldahl Nitrogen as N | т Бу Discrete Analyser | 0.1 | mg/L | 0.2 | 0.3 | 0.3 | 0.4 | 0.2 |
| | | | mg/L | U. <u>L</u> | 0.0 | 0.0 | V. T | V.E |
| EK062G: Total Nitrogen as N (Th | (N + NOx) by Discrete An | | ma/l | 0.0 | 0.2 | 0.2 | 0.4 | 0.0 |
| ` Total Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.3 | 0.3 | 0.4 | 0.2 |
| EK067G: Total Phosphorus as P | by Discrete Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | 0.03 | <0.01 | <0.01 |

Page : 4 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | B2S | C2S | D2S | A3S | B3S |
|--------------------------------------|---------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Clie | ent samplii | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-006 | EM1114567-007 | EM1114567-008 | EM1114567-009 | EM1114567-010 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 26 | 30 | 44 | 60 | 60 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.007 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | 0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.002 | 0.002 | 0.003 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.005 | 0.006 | <0.005 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercury | y by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete | Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.11 | 0.10 | 0.10 | 0.11 | 0.12 |
| EK057G: Nitrite as N by Discrete Ar | nalvser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete A | nalvser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | 0.02 | 0.01 | <0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (N | | veor | J | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | 0.02 | 0.01 | <0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen By | | | | | 5.5 | 3.3. | 3.3. | 5.5. |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.4 | 0.3 | 0.2 | 0.2 | 0.1 |
| , | | | | VI-7 | 0.0 | | VI.2 | 5 11 |
| EK062G: Total Nitrogen as N (TKN + | NOX) by Discrete An | 0.1 | mg/L | 0.4 | 0.3 | 0.2 | 0.2 | 0.1 |
| ^ Total Nitrogen as N | | 0.1 | IIIg/L | U.4 | U.3 | U.Z | U.Z | U. I |
| EK067G: Total Phosphorus as P by | | 0.04 | | -0.04 | 40.04 | 40.04 | 0.04 | 40.04 |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | 0.01 | <0.01 |

Page : 5 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | C3S | D3S | A4S | B4S | C4S |
|-------------------------------------|------------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Cli | ent samplii | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-011 | EM1114567-012 | EM1114567-013 | EM1114567-014 | EM1114567-015 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 28 | 42 | | 50 | 40 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.007 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | 0.0006 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.014 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | 0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.007 | <0.005 | <0.005 | <0.005 | 0.028 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercu | ry by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discret | e Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.09 | 0.10 | 0.10 | 0.10 |
| EK057G: Nitrite as N by Discrete A | nalvser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete A | Analyser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | 0.01 | <0.01 | 0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (| | vser | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | 0.01 | <0.01 | 0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen B | v Discrete Analyser | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.2 | 0.2 | 0.6 | 0.2 |
| EK062G: Total Nitrogen as N (TKN | | | | | | | | |
| ↑ Total Nitrogen as N | - NOX) by Discrete All | 0.1 | mg/L | 0.1 | 0.2 | 0.2 | 0.6 | 0.2 |
| | Dicerete Analyses | V. 1 | g, ⊏ | | | | -10 | 712 |
| EK067G: Total Phosphorus as P by | | 0.01 | mg/L | 0.01 | 0.05 | 0.03 | 0.03 | 0.04 |
| Total Phosphorus as P | | 0.01 | IIIg/L | 0.01 | 0.05 | 0.03 | 0.03 | 0.04 |

Page : 6 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609

ALS

| Sub-Matrix: WATER | | Cli | ent sample ID | D4S | X4S | Y4S | AaS | BbS |
|--------------------------------------|-----------------------|------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Cli | ent sampli | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-016 | EM1114567-017 | EM1114567-018 | EM1114567-019 | EM1114567-020 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 51 | 36 | 45 | 47 | 63 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.011 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.131 | <0.005 | 0.010 | <0.005 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercury | y by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete | Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.09 | 0.08 | 0.10 | 0.10 |
| EK057G: Nitrite as N by Discrete Ar | nalyser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete A | nalvser | | | | | | | • |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (N | Ox) by Discrete Anal | lvser | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen By | Discrete Analyser | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| EK062G: Total Nitrogen as N (TKN + | · NOx) by Discrete An | alvser_ | | | | | | |
| ↑ Total Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| EK067G: Total Phosphorus as P by | Discrete Analyser | | J. Company | | | | | 1 |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.01 | 0.04 | 0.02 | 0.02 |
| Total Filospilorus as F | | 0.01 | g, = | -0.01 | 0.01 | V.V T | 0.02 | 0.02 |

Page : 7 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | CcS | DdS | A5S | B5S | C5S |
|--|---------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Clie | ent samplii | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-021 | EM1114567-022 | EM1114567-023 | EM1114567-024 | EM1114567-025 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 47 | 62 | 60 | 47 | 48 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.003 | 0.002 | 0.004 | 0.003 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | <0.005 | <0.005 | <0.005 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercury b | y FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete A | nalyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.11 | 0.09 | 0.11 | 0.08 | 0.10 |
| EK057G: Nitrite as N by Discrete Anal | vser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete Ana | lyeor | | J. Company | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| | | | 9 | 0.01 | 0.01 | 0.0. | 3.3. | 0.01 |
| EK059G: Nitrite plus Nitrate as N (NO) | x) by discrete Anai | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Nitrite + Nitrate as N | | 0.01 | IIIg/L | ~ 0.01 | ~0.01 | N.01 | \0.01 | ~0.01 |
| EK061G: Total Kjeldahl Nitrogen By Di | | 0.1 | ma/l | 0.4 | 0.0 | | 0.0 | |
| Total Kjeldahl Nitrogen as N | | | mg/L | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| EK062G: Total Nitrogen as N (TKN + N | Ox) by Discrete An | | | | | | | |
| ^ Total Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| EK067G: Total Phosphorus as P by Di | screte Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.04 | 0.05 | 0.09 | 0.03 |

Page : 8 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Suspended Solids (SS) | Sub-Matrix: WATER | | Clie | ent sample ID | D5S | A6S | B6S | C6S | D6S |
|--|--------------------------------------|---------------------------|-----------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| Application Color | | CI | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Suspanded Solids (SS) | Compound | CAS Number | LOR | Unit | EM1114567-026 | EM1114567-027 | EM1114567-028 | EM1114567-029 | EM1114567-030 |
| Suspended Solids (SS) 5 mg/L | EA025: Suspended Solids | | | | | | | | |
| Color Total Metals by ICP-MS T440-39-2 0.001 mg/L 0.006 | Suspended Solids (SS) | | 5 | mg/L | 67 | | | | |
| Asenic 7440-38-2 0.001 mg/L 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0. | Suspended Solids (SS) | | 5 | mg/L | | 223 | 114 | 128 | 154 |
| Barlum | EG020T: Total Metals by ICP-MS | | | | | | | | |
| Beryllium | Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Cobat | Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Copper | Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Manganese 7439-96-5 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 | Chromium | 7440-47-3 | 0.001 | mg/L | | <0.001 | | <0.001 | |
| Nickel 7440-02-0 0.001 mg/L 0.002 0.003 0.002 0.002 0.002 0.002 0.002 0.002 0.002 0.002 0.002 0.002 0.003 0.002 0.002 0.002 0.002 0.003 0. | Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.010 |
| Lead 7439-92-1 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.0 | Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.003 | 0.002 | 0.002 | 0.002 |
| Zinc 7440-66-6 0.005 mg/L < 0.005 0.006 0.006 < 0.005 < 0.005 < 0.005 Iron 7439-89-6 0.05 mg/L < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 CEG035T: Total Recoverable Mercury by FIMS ***Total Recoverable Mercury by FIMS** ***Total Recoverable Mercury 7439-97-6 0.0001 mg/L < 0.0001 | Lead | 7439-92-1 | 0.001 | mg/L | | <0.001 | | <0.001 | |
| Iron 7439-99-6 0.05 mg/L < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < | Vanadium | 7440-62-2 | | mg/L | | <0.01 | | | 1.1 |
| Mercury 7439-97-6 0.0001 mg/L <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 | Zinc | 7440-66-6 | | mg/L | | 0.006 | 0.006 | <0.005 | |
| Mercury 7439-97-6 0.0001 mg/L <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.0 | Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EK055G: Ammonia as N by Discrete Analyser Ammonia as N | EG035T: Total Recoverable Mercury | y by FIMS | | | | | | | |
| Ammonia as N 7664-41-7 0.01 mg/L 0.10 0.10 0.10 0.11 0.11 0.11 0.10 EK057G: Nitrite as N by Discrete Analyser Nitrite as N | Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Nitrite as N by Discrete Analyser Nitrite as N by Discrete Analyser Nitrite as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 | EK055G: Ammonia as N by Discrete | Analyser | | | | | | | |
| Nitrite as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0 | Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.10 | 0.11 | 0.11 | 0.10 |
| Nitrate as N by Discrete Analyser Sknots as N by Discrete Analyser Sknots as N 14797-55-8 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.0 | EK057G: Nitrite as N by Discrete Ar | nalyser | | | | | | | |
| Nitrate as N | Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Nitrate as N | EK058G: Nitrate as N by Discrete A | nalvser | | | | | | | |
| Nitrite + Nitrate as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.0 | Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Nitrite + Nitrate as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.0 | EK059G: Nitrite plus Nitrate as N (N | IOx) by Discrete Ana | lvser | | | | | | |
| EK061G: Total Kjeldahl Nitrogen By Discrete Analyser Total Kjeldahl Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.1 0.3 EK062G: Total Nitrogen as N (TKN + NOx) by Discrete Analyser Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.1 0.3 EK067G: Total Phosphorus as P by Discrete Analyser | Nitrite + Nitrate as N | | | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Total Kjeldahl Nitrogen as N | | Discrete Analyser | | | | | | | |
| Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.1 0.1 0.3 EK067G: Total Phosphorus as P by Discrete Analyser | Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.2 | 0.2 | 0.1 | 0.3 |
| Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.1 0.1 0.3 EK067G: Total Phosphorus as P by Discrete Analyser | EK062G: Total Nitrogen as N (TKN + | NOx) by Discrete Ar | nalys <u>er</u> | | | | | | |
| | ` Total Nitrogen as N | | | mg/L | 0.2 | 0.2 | 0.2 | 0.1 | 0.3 |
| | EK067G: Total Phosphorus as P by | Discrete Analys <u>er</u> | | | | | | | |
| | Total Phosphorus as P | | 0.01 | mg/L | 0.02 | 0.02 | 0.02 | <0.01 | 0.05 |

Page : 9 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Cli | ent sample ID | A1B | B1B | C1B | D1B | A2B |
|--------------------------------------|-----------------------|--------------|-----------------|-------------------|-------------------|-------------------|---------------------------------------|-------------------|
| | CI | lient sampli | ing date / time | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-031 | EM1114567-032 | EM1114567-033 | EM1114567-034 | EM1114567-035 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 177 | 216 | 143 | 147 | 146 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.003 | 0.002 | 0.002 | 0.002 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.013 | 0.009 | <0.005 | <0.005 | 0.006 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercui | ry by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete | e Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.12 | 0.11 | 0.09 | 0.12 |
| EK057G: Nitrite as N by Discrete A | nalvser | | | | | | | • |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete A | \nalvser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (I | | lvser | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen By | | | 5 | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.1 | 0.2 | 0.2 | 0.2 |
| EK062G: Total Nitrogen as N (TKN | + NOx) by Discrete Ar | nalyser | | | | | | |
| ^ Total Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.1 | 0.2 | 0.2 | 0.2 |
| EK067G: Total Phosphorus as P by | Discrete Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.01 | 0.02 | <0.01 | 0.04 |
| | | | 3 | · · | 1.4. | 1 | · · · · · · · · · · · · · · · · · · · | |

Page : 10 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | B2B | C2B | D2B | A3B | ВЗВ |
|--|----------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Clie | ent samplii | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-036 | EM1114567-037 | EM1114567-038 | EM1114567-039 | EM1114567-040 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 129 | 183 | 99 | 129 | 161 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | 0.0001 | <0.0001 | <0.0001 | 0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.011 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.003 | 0.003 | 0.004 | 0.002 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | <0.005 | 0.006 | <0.005 | <0.005 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercury b | y FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete A | nalyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.13 | 0.10 | 0.09 | 0.12 | 0.09 |
| EK057G: Nitrite as N by Discrete Anal | vser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete Ana | lyser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (NO) | | veor | J | | | | | |
| Nitrite + Nitrate as N | L) Dy Discrete Aliai | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen By Di | | 3.01 | y. = | -0.01 | -0.01 | -0.01 | -0.01 | .0.01 |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | 0.2 | 0.1 | 0.1 | 0.2 |
| | | | 9, = | •••• | V.2 | V.1 | V. 1 | V.2 |
| EK062G: Total Nitrogen as N (TKN + N | Ox) by Discrete An | 0.1 | mg/L | <0.1 | 0.2 | 0.1 | 0.1 | 0.2 |
| ↑ Total Nitrogen as N | | 0.1 | IIIg/L | ~ U. I | U.Z | U.1 | U.1 | U.Z |
| EK067G: Total Phosphorus as P by Dis | screte Analyser | 0.04 | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | 0.02 | 0.01 | 0.06 | 0.02 | 0.04 |

Page : 11 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | C3B | D3B | A4B | B4B | C4B |
|---------------------------------------|----------------------|-------------|----------------|---|-------------------|-------------------|-------------------|-------------------|
| | Cli | ent samplii | ng date / time | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-041 | EM1114567-042 | EM1114567-043 | EM1114567-044 | EM1114567-045 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 114 | 188 | | 158 | 94 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.005 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.009 | 0.009 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.004 | 0.004 | 0.005 | 0.005 | 0.005 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.007 | 0.008 | 0.007 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.08 | <0.05 | 0.06 |
| EG035T: Total Recoverable Mercury | by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete | Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.13 | 0.12 | 0.10 | 0.09 |
| EK057G: Nitrite as N by Discrete Ana | alvser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete An | alveor | | J. | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| | | | 9, _ | 0.01 | 3.01 | 0.01 | 3.3. | 0.01 |
| EK059G: Nitrite plus Nitrate as N (NO | OX) by Discrete Anal | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Nitrite + Nitrate as N | | 0.01 | my/L | ~ 0.01 | ~0.01 | ~0.01 | \0.01 | ~0.01 |
| EK061G: Total Kjeldahl Nitrogen By I | | 0.1 | ma/l | <0.1 | -0.1 | <0.1 | 0.4 | <0.1 |
| Total Kjeldahl Nitrogen as N | | | mg/L | <u.1< td=""><td><0.1</td><td><0.1</td><td>0.1</td><td><0.1</td></u.1<> | <0.1 | <0.1 | 0.1 | <0.1 |
| EK062G: Total Nitrogen as N (TKN + | NOx) by Discrete An | | | | | | | |
| ^ Total Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | <0.1 | 0.1 | <0.1 |
| EK067G: Total Phosphorus as P by D | Discrete Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |

Page : 12 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | D4B | X4B | Y4B | AaB | BbB |
|---------------------------------------|-----------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Cli | ent samplii | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-046 | EM1114567-047 | EM1114567-048 | EM1114567-049 | EM1114567-050 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 163 | 141 | 128 | 116 | 137 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.005 | 0.007 | 0.006 | 0.005 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | 0.0001 | <0.0001 | <0.0001 | 0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.011 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.005 | 0.005 | 0.004 | 0.004 | 0.005 |
| Lead | 7439-92-1 | 0.001 | mg/L | 0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.010 | 0.006 | 0.008 | 0.007 | 0.006 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.06 | 0.06 | 0.10 | 0.06 |
| EG035T: Total Recoverable Mercury | by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete A | Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.08 | 0.16 | 0.11 | 0.12 | 0.11 |
| EK057G: Nitrite as N by Discrete Ana | llvser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete Ana | alvser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | 0.04 | <0.01 | <0.01 | 0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (NO | | lveor | J. | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | 0.04 | <0.01 | <0.01 | 0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen By D | | 0.01 | g, = | U.U. | -0.01 | 0.01 | 0.01 | .0.01 |
| Total Kjeldahl Nitrogen as N | Jiscrete Analyser | 0.1 | mg/L | <0.1 | <0.1 | <0.1 | 0.1 | <0.1 |
| | | | mg/L | ~0.1 | 50.1 | 50.1 | U. I | 50.1 |
| EK062G: Total Nitrogen as N (TKN + N | NOx) by Discrete An | | | 40.4 | 40.4 | 40.4 | 0.4 | 40.4 |
| ↑ Total Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | <0.1 | 0.1 | <0.1 |
| EK067G: Total Phosphorus as P by D | iscrete Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.02 | <0.01 | 0.03 | <0.01 |

Page : 13 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | СсВ | DdB | A5B | B5B | C5B |
|-----------------------------------|---------------------------|------------|------------------|-------------------|--|-------------------|-------------------|-------------------|
| | Clie | ent sampli | ng date / time | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-051 | EM1114567-052 | EM1114567-053 | EM1114567-054 | EM1114567-055 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 166 | 104 | 174 | 187 | 155 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.007 | 0.005 | 0.005 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0001 | <0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.011 | 0.010 | 0.011 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | 0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.005 | 0.005 | 0.005 | 0.005 | 0.004 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | 0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | <0.005 | 0.011 | 0.006 | 0.006 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.09 | 0.17 | 0.14 | 0.12 |
| EG035T: Total Recoverable Mer | cury by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Disci | rete Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.13 | 0.08 | 0.12 | 0.10 |
| EK057G: Nitrite as N by Discrete | Analyser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discret | ο Analyser | | , and the second | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| | | | 9-2 | | | | | |
| EK059G: Nitrite plus Nitrate as N | N (NOX) by discrete Analy | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| | | 0.01 | my/L | -0.01 | \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\ | N.U I | 70.01 | 70.01 |
| EK061G: Total Kjeldahl Nitrogen | | 0.1 | mg/L | <0.1 | 0.1 | <0.1 | 0.2 | 1.0 |
| Total Kjeldahl Nitrogen as N | | | mg/L | ~ U. I | U.T | ~ 0.1 | U.Z | 1.0 |
| EK062G: Total Nitrogen as N (TK | N + NOx) by Discrete Ana | | | -0.4 | | -0.4 | | 4.0 |
| Total Nitrogen as N | | 0.1 | mg/L | <0.1 | 0.1 | <0.1 | 0.2 | 1.0 |
| EK067G: Total Phosphorus as P | by Discrete Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | 0.02 | 0.14 | 0.02 | 0.05 | 0.05 |

Page : 14 of 14

Work Order : EM1114567 Amendment 1
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | D5B | A6B | B6B | C6B | D6B |
|--|---------------------|-------------|----------------|---|--|-------------------|-------------------|-------------------|
| | Cli | ent samplii | ng date / time | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-056 | EM1114567-057 | EM1114567-058 | EM1114567-059 | EM1114567-060 |
| EA025: Suspended Solids | | | | | | | | |
| Suspended Solids (SS) | | 5 | mg/L | 99 | 111 | 156 | 19 | 43 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.007 | 0.005 | 0.006 | 0.005 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0001 | 0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.012 | 0.011 | 0.010 | 0.012 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.006 | 0.005 | 0.005 | 0.005 | 0.006 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | 0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.025 | 0.008 | 0.009 | 0.014 |
| Iron | 7439-89-6 | 0.05 | mg/L | 0.10 | 0.16 | 0.17 | 0.16 | 0.14 |
| EG035T: Total Recoverable Mercury b | y FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete A | nalyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.09 | 0.08 | 0.08 | 0.11 | 0.10 |
| EK057G: Nitrite as N by Discrete Analy | vser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete Anal | lyeor | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| | | | 9 | 0.01 | 0.01 | 0.0. | 3.3. | 0.0. |
| EK059G: Nitrite plus Nitrate as N (NOx | t) by Discrete Anal | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Nitrite + Nitrate as N | | 0.01 | my/L | ~ 0.01 | \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\ | ~ 0.01 | ~ U.U1 | ~0.01 |
| EK061G: Total Kjeldahl Nitrogen By Di | | 0.1 | ma/l | <0.1 | 40.1 | 0.4 | 0.4 | <0.1 |
| Total Kjeldahl Nitrogen as N | | | mg/L | <u.1< td=""><td><0.1</td><td>0.1</td><td>0.1</td><td><0.1</td></u.1<> | <0.1 | 0.1 | 0.1 | <0.1 |
| EK062G: Total Nitrogen as N (TKN + N | Ox) by Discrete An | | | | | | | |
| ^ Total Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | 0.1 | 0.1 | <0.1 |
| EK067G: Total Phosphorus as P by Dis | screte Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | 0.04 | 0.02 | 0.04 | 0.04 | 0.02 |



Appendix C Geochemistry of Metal Scavenging: Technical Note

To Iron Road Ltd. Date 26 June 2014

From Land Quality and Sediment Chemistry, Jacobs Project No VE23730

Subject Controls On Trace Metal and Metalloid Geochemistry

Introduction

There are a number of process which control the geochemistry of metals and metalloids at the seawater/sediment interface. These controls are related to redox and, as a consequence the changes from anoxic to oxic geochemical environment. The relatively straight forward geochemistry can be complicated by the trace metal interaction with inorganic and organic ligands arising under anoxic conditions. Common inorganic complexes forming in anoxic pore waters include phosphate, carbonate, bicarbonate, bisulphite, sulphur and chloride. In addition marine organism can accumulate metals and metalloids (Langston, 1984) with the bioaccumulation within species of alga (Luoma et al., 1982) and marine invertebrates such as polycheates (Waring and Maher, 2005). While anaerobic microorganisms may play an important part in actively scavenging trace metals from the water column and surrounding sediments in anoxic sediments (Gaillard and Webb, 1998). Vanadium (V) has also been identified as an important element in the blood chemistry of sea squirts of the ascidian family (Michibata and Sakurai, 1990) and numerous studies have shown these marine filter feeders concentrate vanadium in their bodies to a level one million times higher than the concentration of vanadium in seawater. Finally organic ligands such as those which form colloidal complexes can increase solubility by a factor of ten beyond the theoretical inorganic solubility (Kersten, 1998). All of these interactions can either together or on their own create conditions for a trace metal "sink" which actively draws metals and metalloids, such as arsenic (As) and V into the sediment.

The data from the Jacobs survey at Cape Hardy suggest that the south to north transects appear to show evidence of a more increased redox in the south with the most anoxic sediment in the northern reaches (Jacobs, 2014). The concentrations of V, chromium (Cr) and, in particular iron (Fe) and As, increase in the sediment as they become more anoxic.

To understand the elevated concentrations of metals identified in sediments across the study area it is important to understand that the chemical or valence state of these elements drives their accumulation. Across the study area the highest concentration of As and Fe are noted to occur in the sediment with the lowest redox potential. Conversely, Zn concentrations are noted to decrease in the sediment with decreasing redox. These accumulation behaviours would appear counter intuitive to the anticipated elemental geochemistry, which would dictate that all these three elements should display similar positive ion speciation's. The behaviour of these metals and metalloids and the controls of their valence state and physical state within the water column and surficial sediments are discussed below.

<u>Arsenic</u>

Arsenic (As) concentrations were identified in elevated concentrations in sediments from several samples in the north eastern extent of the study area. As is a metalloid which can display a high mobility in anoxic sediment-water systems. The valence state of As is important as arsenite (As ³⁺) is more toxic for biological species and more mobile than arsenate (As ⁵⁺). The arsenate-arsenite interconversion in response to redox (Eh) reduction across oxic-anoxic interfaces can change the speciation from valence As ³⁺ to As ⁵⁺. Kersten (1988) described the following reaction:

$$HAsO_4^{2-} + 2H^+ + 2e^- = HAsO_3^{2-} + H_2O$$

Arsenate Arsenite

JACOBS

At pH 7.3 (which is typical of anoxic sediments) and no significant complexation by As oxyanions, the ration can be expressed as a function of redox:

$$\log \left(\frac{As^{3+}}{As^{5+}}\right) = \left(\frac{-Eh(mV)}{29.6}\right) - 6.46$$

Whilst an assessment of the redox conditions of the sediments was not undertaken during sampling, a strong sulphurous odour was reported. Typically the onset of sulfate reduction occurs from redox -0.1V and can still be occurring at 0.5V. So assuming a redox 0.1V and a mid point redox of 0.25V, will impact the ratio and the more toxic form of As (i.e. As ³⁺) will dominate in decreasing redox. Studies have shown that under these conditions As will precipitate in the presence of sulfide and adsorption on Fe (Peterson and Carpenter, 1986).

Arsenic may substitute for sulfide in pyrite (FeS₂) and thus the presence of As in reduced (i.e. anoxic) sediments is likely to be represented by arseno-pyrite (Appelo and Postma, 2005). The oxidation of pyrite and subsequent release of arsenic can be one of the major causes of arsenic in groundwater or sediment pore water (Smedley and Kinniburgh, 2002). A separate process associated with accumulation of As in sediments is the sorption of arsenic to iron oxide particles during deposition (Ravenscroft *et al.*, 2001). These theorems when considered with the high concentrations of Fe recorded from sediment samples collected in the northeastern corner of the study area, may help to explain the elevated measurements of As in the same samples.

Anaerobic sediments elevated in As that become aerobic through re-suspending sediment will impact the arsenic speciation and has the potential to increase arsenic levels in the overlying water column. However due to the high concentrations of Fe in the sediment within the study area the arsenic would be readily reabsorbed back into the sediment.

Vanadium and Chromium

The increased concentration of V and Cr in anoxic sediment, as identified across the study area is consistent with the geochemistry and thermodynamics of these trace elements. Calvert and Peterson (1986) have shown that a change in valency means V and Cr compounds are precipitated under anoxic conditions. Hence, the enrichment of trace elements indicates that the host sediments accumulated the metals under anoxic conditions. This is supported by the Eh-pH diagrams for V and Cr presented below in Figure 1. . With increased redox V favours a solid species and tends to precipitate, and with a change of redox such as that associated with resuspension of sediment in the water column, V is more likely to be present as a dissolved species. The same relationship is evident the Eh-pH diagram for chromium examined in Figure 2:

JACOBS




Figure 1 Vanadium redox state at varying pH (Takeno (2005)




Figure 2 Chromium redox state at varying pH (Takeno, 2005)



Zinc

In accordance with the above noted elemental geochemistry it would be expected that with increased redox that Zn would also be present as an insoluble sulphide precipitate in the sediment. It is noted however that Zn concentrations were observed to decrease with decreasing sediment redox across the study area.

A possible explanation for this observation is that colloidal chemistry is rendering Zn more 'soluble' resulting in it being present in the abundant organic matter (algal) or detritus. For this to be a viable consideration the Zn would need to be present in a Zn-acetate complex and acetate would need to be approximately four orders of magnitude more abundant than bisulfide (Gammons and Frandsen, 2001). Notwithstanding, it is considered likely that Zn is bound as sulfide phase in the sediments and will therefore maintain its bioavailability because of the continued cycling between pore waters and surface sediments due to physical mixing and bioturbation (Atkinson *et al.*, 2007).

Conclusions

As noted during the video tow survey of the study area the regions of anoxic sediment showed signs of seasquirts (known to accumulate V) and bioturbation indicating the presence of polychaete worms, several species of which are known to bioaccumulate As (Waring and Maher, 2005). Therefore it is considered possible the elevated levels of As and V identified in the sediments across the northern region of the study area are a result of the naturally prevailing chemical properties of the sediment (high iron levels and low redox) acting together with bioaccumulation by invertebrates. While low concentrations of Zn reported in the sediment are potentially the result of these same bioturbation and biological processes. Aerobic conditions would be created by the localised resuspension of sediment creating allowing the release of Zn (As and V) back into the water column while seasquirt and polychaetes then scavenge back the As and V into the sediment. This type of selective scavenging could account for the differences in trace metal concentrations across the study area.



References

Appelo CAJ and Postma D (2005) Geochemistry, groundwater and pollution. 2nd edition, Balkema, Leidan

Atkinson CA, Jolley DF and Simpson SL (2007) Effect of overlying water pH, dissolved oxygen, salinity and sediment disturbances on metal release and sequestration from metal contaminated marine sediments

Calvert SE and Peterson ML (1993) *Geochemistry of Recent oxic and anoxic marine sediments: Implications for the geological record* Marine Geology 113: 67-88

Gaillard JF and Webb SM (1998) *Biogeochemical Aspects of Metal-Microbes Interactions*. In: Transport, Fate and Effects of Silver in the Environment, 5th International Argentum Conference Proceedings, A. W. Andren and T. W. Bober, Eds. University of Wisconsin Sea Grant Publication No. WISCU-W-97001, 77-86.

Gammons CH and Frandsen AK (2001) Fate and Transport of metals in H2S-rich waters at a treatment wetland. Geochemical Transactions 2001 2:1

Jacobs (2014) *Central Eyre Iron Project; Port Site Marine Assessment* IRRT-RPT-00089 produced by Jacobs Adelaide Office for Iron Road Limited

Kersten M (1988) Geochemistry of Priority Pollutants in Anoxic Sludges: Cadmium, Arsenic, Methyl Mercury and Chlorinated Organics In Chemistry and Biology of Solid Waste ed Salomons and Forstner p189-195

Langston WJ (1984) Availability of arsenic to estuarine and marine organisms: a field and laboratory evaluation. Marine Biology 80: 143-154

Luoma SN, Bryan GW and Langston WJ (1982) *Scavenging of Heavy Metals from Particulates by Brown Seaweed*, Marine Pollution Bulletin 13: 394-396

Michibata H and Sakurai H (1990) *Vanadium in Ascidians*. In ND Chasteen (eds), Vanadium Biological Systems, Physiology and Biochemistry pp 153-171

Peterson ML and Carpenter R (1986) *Biogeochemical processes affecting total arsenic and arsenic species distribution in an intermittently anoxic fjord.* Mar Chem 12:295-321

Ravenscroft P, McArthur JM and Hoque BA (2001). *Geochemical and paleohydrological on pollution of groundwater by arsenic*. In W.R. Chappell *et al.*(eds), *Arsenic exposure and health effects IV*. Elsevier, Amsterdam.

Smedley PL and Kinniburgh DG (2002) A review of the source, behaviour and distribution of arsenic in natural waters. Appl. Geochem. 17, 517-568.

Takeno N (2005) *Atlas of Eh-pH diagrams. Intercomparison of thermodynamic databases*. Geological Survey of Japan Open File Report No.419. National Institute of AdvancedIndustrial Science and Technology Research Center for Deep Geological Environments.

Waring J and W Maher (2005) *Arsenic bioaccumulation and species in marine Poylchaeta*. Applied Organometallic Chemistry **19**:917-929



Appendix D Water Quality Laboratory Results

| | | | CHAIN OF CUSTODY FORM | TODY FORM | | | | | 7 |
|---|-----------------|---------------------------|-----------------------------------|----------------|----------|-------------|--------|--------------------------|--------------------|
| From : SKM Pty Ltd ABN: 37 001 024 095 | 95. Ta | | ALS | ., | | | | · 1 | THE PARTY NAMED IN |
| Level 5, 33 King William St. Adelaide, SA 5000 pb: (08) 8424 3800 | William St, Ade | laide, SA 5000 14 3810 | | | _ | | Cont | Container Identification | tion |
| | | | | | Size | 500ml | 250 mi | EM 09 | |
| CHOTE NUMBER | | L. Lolect NO. AFERGACION | -Ederor Zor | | Type | Green Label | purple | red | |
| do o a sement | | _ | ve23610.60a | , f | Preserv | 25 | | Cr, R | |
| Job Code: | | Project Manag | Project Manager: Dr. Greg Barbara | | | s EA(| | , Cd, | |
| Custody seal intact? | | Sampler(s): Greg Barbara | reg Barbara | | Analytes | Sólid | | a, Be | |
| Received for Laborator | چ ر | Checked: | | | | nded | utrie | Meta As, E I, Mn, | |
| b. A | 7 | Date: 23/11/2011 | 3 | | | uspe | | tetal | |
| i ah id | | | Sample Identification | Comment | | | | 9 | |
| \ | 19/12/2011 | Water | A1S | Surface sample | | ٠ ٠ | < | < | |
| 2 | 19/12/2011 | Water | B1S | Surface sample | | < | < | < | |
| 7 | 19/12/2011 | Water | C1S | Surface sample | | < | < | < | |
| 7 | 19/12/2011 | Water | D18 | Surface sample | | < | < | ٧ | |
| 5 | 19/12/2011 | Water | A2S | Surface sample | | ~ | < | < | |
| 6 | 19/12/2011 | Water | B2S | Surface sample | | \ | < | < | |
| 7 | 19/12/2011 | Water | C2S | Surface sample | | < | < | < | |
| Ø | 19/12/2011 | Water | D2S | Surface sample | | < | < | < | - |
| 9 | 19/12/2011 | Water | A3S | Surface sample | | \ | < | < | |
| 0 | 19/12/2011 | Water | B3S | Surface sample | | < | < | < | |
| 7 | 19/12/2011 | Water | C3S | Surface sample | | < | < | < | |
| 12/ | 19/12/2011 | Water | D3S | Surface sample | | < | < | < | |
| ~ | 19/12/2011 | Water | A4 🔇 | Surface sample | | < | ۲ | <u> </u> | + |
| 7 | 19/12/2011 | Water | B4 . | Surface sample | | < - | < | < | |

Spoker to hreg artered to do Nothernal Division

Surface sample

Environmental Division
Melbourne
Work Order

EM1114567

Telephone : +61-3-8549 9600

Bottom Sample Bottom Samp Surface samp Bottom Sample Bottom Samp Bottom Sample Bottom Sami Bottom Samp Bottom Samp Bottom Sampl Surface sample Surface samp ottom Sami Surface sample ottom Sampl ottom Sam ottom Sampl urface sample ottom Samp

bed using a Petite Ponar and transferred into sample containers

mail results to gbarbara@globalskm.com for queries with samples call Greg Barbara on 0459826420

Alex Haines

From: Barbara, Greg (SKM) [GBarbara@globalskm.com]

Sent: Thursday, 22 December 2011 3:15 PM

Alex Haines

Subject: RE: VE23610.609 (EM1114567)

Please report as per the sample bottle eg. A6B etc

Thank you

Greg

× Please consider the environment before printing this e-mail

Dr Greg Barbara

Marine Scientist - SKM (VESA)

BSc (Marine Biology) (Hon) PhD (Marine Biology)

SINCLAIR KNIGHT MERZ (SKM)L5, 33 King William St, Adelaide SA 5000 **P:** +61 8 8424 6508 |**F:** +61 8 8424 3810 |**M:** 0459 826420 |**E:** GBarbara@globalskm.com

Sinclair Knight Merz

achieve outstanding client success

For further information, visit our website www.globalskm.com

From: Alex Haines [mailto:Alex.Haines@alsglobal.com]
Sent: Thursday, 22 December 2011 2:43 PM
To: Barbara, Greg (SKM)

Cc: Carol Walsh

Subject: VE23610.609 (EM1114567)

Hi Greg

this is also true for bottom water samples. Which IDs would you like us to report, the ones on the the COC is missing this S for numerous samples eg A4 on the CoC is written as A4S on the sample, I am logging this work order for you and have a question concerning the IDs on some of the samples Thank you samples or the ones on the CoC? All surface water samples are written with an S following the code on the samples themselves but

Regards

WORK ORDER CREATION/LOGIN OFFICER **Alex Haines**

ALS | Environmental (General Environmental Group)

Address

4 Westall Road Springvale VIC 3171

PHONE +61 3 8549 9600 FAX +61 3 8549 9601

Winner of the inaugural CARE Award 2011 – Sustainable Technology & Innovation: Reduction in Sample Volumes – Improving quality, safety, efficiency and sustainability in environmental practices environmental practices

×

www.alsglobal.com

P Please consider the environment before printing this email.

must not copy, distribute, print or otherwise use the information. Email may be stored by the you must notify the sender immediately by return email and then delete all copies of this email. You The information contained in this email is confidential. If the reader is not the intended recipient then Company to support operational activities. All information will be held in accordance with the

Click here to report this email as spam.

transferred information and should advise us immediately upon receipt of any discrepancy. All email lost by any mistaken transmission to you. If you have received this message in error please delete the SKM is committed to working with its clients to deliver a sustainable future for all. Please consider the environment before printing this e-mail. Notice - This message contains confidential information intended only for the exclusive use of the addressee named above. No confidentiality is waived or sent to SKM will be intercepted, screened and filtered by SKM or its approved Service Providers. document and notify us immediately. Any opinion, text, documentation or attachment received is valid as at the date of issue only. The recipient is responsible for reviewing the status of the

ALS Group: Click here to report this email as spam.

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

SAMPLE RECEIPT NOTIFICATION (SRN)

Comprehensive Report

: EM1114567 Work Order

Amendment : 2

Client SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : DR GREG BARBARA Contact : Carol Walsh

> : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia ADELAIDE SA, AUSTRALIA 5000

3171

F-mail : gbarbara@globalskm.com E-mail : carol.walsh@alsglobal.com

Telephone : +61 08 8424 3800 Telephone : +61-3-8549 9608 Facsimile Facsimile : +61-3-8549 9601 : +61 08 8424 3810

: VE23610 609 Project Page : 1 of 5

Order number C-O-C number Quote number : EA2011SINKNI0361 (ADBQ/037/11) : ----

Site : GB QC Level : NEPM 1999 Schedule B(3) and Sampler ALS

QCS3 requirement

Dates

Address

Date Samples Received : 22-DEC-2011 Issue Date : 30-MAR-2012 Client Requested Due Date Scheduled Reporting Date : 28-FFB-2012 28-FEB-2012

Delivery Details

Mode of Delivery Temperature : Carrier : 16-22 - Ice bricks present

No. of coolers/boxes No. of samples received . 3 : 60 Security Seal No. of samples analysed : 60 : Intact.

General Comments

- This report contains the following information:
 - Sample Container(s)/Preservation Non-Compliances
 - Summary of Sample(s) and Requested Analysis
 - Proactive Holding Time Report
 - Requested Deliverables
- Samples received in appropriately pretreated and preserved containers.
- Please refer to the Proactive Holding Time Report table below which summarises breaches of recommended holding times that have occurred prior to samples/instructions being received at the laboratory. The absence of this summary table indicates that all samples have been received within the recommended holding times for the analysis requested.
- Please direct any queries related to sample condition / numbering / breakages to Peter Ravlic.
- Analytical work for this work order will be conducted at ALS Melbourne.
- Sample Disposal Aqueous (14 days), Solid (90 days) from date of completion of work order.

: 30-MAR-2012 Issue Date Page : 2 of 5

Work Order

: EM1114567 Amendment 2 Client : SINCLAIR KNIGHT MERZ



Sample Container(s)/Preservation Non-Compliances

All comparisons are made against pretreatment/preservation AS, APHA, USEPA standards.

• No sample container / preservation non-compliance exist.

Summary of Sample(s) and Requested Analysis

| Some items described below may be part of a laboratory process necessary for the execution of client requested tasks. Packages may contain additional analyses, such as the determination of moisture content and preparation of moisture content and | | | | | |
|---|--|--|--|---------------------------|------------------|
| EM1114567-001 19-DEC-2011 15:00 A1S ✓ ✓ EM1114567-002 19-DEC-2011 15:00 B1S ✓ ✓ EM1114567-003 19-DEC-2011 15:00 C1S ✓ ✓ EM1114567-004 19-DEC-2011 15:00 D1S ✓ ✓ EM1114567-005 19-DEC-2011 15:00 A2S ✓ ✓ EM1114567-006 19-DEC-2011 15:00 B2S ✓ ✓ EM1114567-007 19-DEC-2011 15:00 C2S ✓ ✓ EM1114567-008 19-DEC-2011 15:00 D2S ✓ ✓ EM1114567-009 19-DEC-2011 15:00 B3S ✓ ✓ EM1114567-010 19-DEC-2011 15:00 C3S ✓ ✓ EM1114567-011 19-DEC-2011 15:00 D3S ✓ ✓ EM114567-013 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-014 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-016 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-0 | process neccess tasks. Packages the determinatio tasks, that are incli if no sampling default to 15:00 date is provided, laboratory for p | ary for the execut may contain addition of moisture couded in the package. time is provided, on the date of sail the sampling date processing purposes | tion of client requested ional analyses, such as ontent and preparation the sampling time will mpling. If no sampling will be assumed by the |)8 + NO2 + NO3 + NH3 + | 3T al) (NEPM) |
| EM1114567-001 19-DEC-2011 15:00 A1S ✓ ✓ EM1114567-002 19-DEC-2011 15:00 B1S ✓ ✓ EM1114567-003 19-DEC-2011 15:00 C1S ✓ ✓ EM1114567-004 19-DEC-2011 15:00 D1S ✓ ✓ EM1114567-005 19-DEC-2011 15:00 A2S ✓ ✓ EM1114567-006 19-DEC-2011 15:00 B2S ✓ ✓ EM1114567-007 19-DEC-2011 15:00 C2S ✓ ✓ EM1114567-008 19-DEC-2011 15:00 D2S ✓ ✓ EM1114567-009 19-DEC-2011 15:00 B3S ✓ ✓ EM1114567-010 19-DEC-2011 15:00 C3S ✓ ✓ EM1114567-011 19-DEC-2011 15:00 D3S ✓ ✓ EM114567-013 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-014 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-016 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-0 | Matrix: WATER | | | NT-(| W-0X |
| EM1114567-001 19-DEC-2011 15:00 A1S ✓ ✓ EM1114567-002 19-DEC-2011 15:00 B1S ✓ ✓ EM1114567-003 19-DEC-2011 15:00 C1S ✓ ✓ EM1114567-004 19-DEC-2011 15:00 D1S ✓ ✓ EM1114567-005 19-DEC-2011 15:00 A2S ✓ ✓ EM1114567-006 19-DEC-2011 15:00 B2S ✓ ✓ EM1114567-007 19-DEC-2011 15:00 C2S ✓ ✓ EM1114567-008 19-DEC-2011 15:00 D2S ✓ ✓ EM1114567-009 19-DEC-2011 15:00 B3S ✓ ✓ EM1114567-010 19-DEC-2011 15:00 C3S ✓ ✓ EM1114567-011 19-DEC-2011 15:00 D3S ✓ ✓ EM114567-013 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-014 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-016 19-DEC-2011 15:00 A4S ✓ ✓ EM114567-0 | | | Client sample ID | VATER - Fotal Nitro | NATER - |
| EM1114567-003 19-DEC-2011 15:00 C1S | EM1114567-001 | 19-DEC-2011 15:00 | A1S | | |
| EM1114567-004 19-DEC-2011 15:00 D1S | EM1114567-002 | 19-DEC-2011 15:00 | B1S | ✓ | ✓ |
| EM1114567-005 19-DEC-2011 15:00 A2S | EM1114567-003 | 19-DEC-2011 15:00 | C1S | ✓ | ✓ |
| EM1114567-006 19-DEC-2011 15:00 B2S | EM1114567-004 | 19-DEC-2011 15:00 | D1S | ✓ | ✓ |
| EM1114567-007 19-DEC-2011 15:00 C2S | EM1114567-005 | 19-DEC-2011 15:00 | A2S | ✓ | ✓ |
| EM1114567-008 19-DEC-2011 15:00 D2S ✓ ✓ EM1114567-009 19-DEC-2011 15:00 A3S ✓ ✓ EM1114567-010 19-DEC-2011 15:00 B3S ✓ ✓ EM1114567-011 19-DEC-2011 15:00 C3S ✓ ✓ EM1114567-012 19-DEC-2011 15:00 D3S ✓ ✓ EM1114567-013 19-DEC-2011 15:00 A4S ✓ ✓ EM1114567-014 19-DEC-2011 15:00 C4S ✓ ✓ EM1114567-015 19-DEC-2011 15:00 C4S ✓ ✓ EM1114567-016 19-DEC-2011 15:00 D4S ✓ ✓ EM1114567-017 19-DEC-2011 15:00 Y4S ✓ ✓ EM1114567-018 19-DEC-2011 15:00 AaS ✓ ✓ EM1114567-019 19-DEC-2011 15:00 BbS ✓ ✓ EM1114567-020 19-DEC-2011 15:00 BbS ✓ ✓ EM1114567-021 19-DEC-2011 15:00 B6S ✓ ✓ EM11145 | EM1114567-006 | 19-DEC-2011 15:00 | B2S | ✓ | ✓ |
| EM1114567-009 19-DEC-2011 15:00 B3S | EM1114567-007 | 19-DEC-2011 15:00 | C2S | ✓ | ✓ |
| EM1114567-010 19-DEC-2011 15:00 B3S | EM1114567-008 | 19-DEC-2011 15:00 | D2S | ✓ | ✓ |
| EM1114567-011 19-DEC-2011 15:00 C3S | EM1114567-009 | 19-DEC-2011 15:00 | A3S | ✓ | ✓ |
| EM1114567-012 19-DEC-2011 15:00 D3S | EM1114567-010 | 19-DEC-2011 15:00 | B3S | ✓ | ✓ |
| EM1114567-013 19-DEC-2011 15:00 A4S EM1114567-014 19-DEC-2011 15:00 B4S EM1114567-015 19-DEC-2011 15:00 C4S EM1114567-016 19-DEC-2011 15:00 D4S EM1114567-017 19-DEC-2011 15:00 X4S EM1114567-018 19-DEC-2011 15:00 Y4S EM1114567-019 19-DEC-2011 15:00 BbS EM1114567-020 19-DEC-2011 15:00 BbS EM1114567-021 19-DEC-2011 15:00 DdS EM1114567-022 19-DEC-2011 15:00 DdS EM1114567-023 19-DEC-2011 15:00 B5S EM1114567-024 19-DEC-2011 15:00 B5S EM1114567-025 19-DEC-2011 15:00 C5S EM1114567-026 19-DEC-2011 15:00 D5S EM1114567-027 19-DEC-2011 15:00 B6S EM1114567-028 19-DEC-2011 15:00 B6S EM1114567-030 19-DEC-2011 15:00 B6S EM1114567-031 19-DEC-2011 15:00 B6S EM1114567-031 19-DEC-2011 15:00 B6S EM1114567-030 19-DEC-2011 15:00 B6S EM1114567-031 19-DEC-2011 15:00 B6S EM1114567-031 19-DEC-2011 15:00 B1B EM1114567-032 19-DEC-2011 15:00 B1B EM1114567-033 19-DEC-2011 15:00 D1B EM1114567-034 19-DEC-2011 15:00 D1B | EM1114567-011 | 19-DEC-2011 15:00 | C3S | ✓ | ✓ |
| EM1114567-014 19-DEC-2011 15:00 B4S | EM1114567-012 | 19-DEC-2011 15:00 | D3S | ✓ | ✓ |
| EM1114567-015 19-DEC-2011 15:00 C4S | EM1114567-013 | 19-DEC-2011 15:00 | A4S | ✓ | ✓ |
| EM1114567-016 19-DEC-2011 15:00 D4S | EM1114567-014 | 19-DEC-2011 15:00 | B4S | ✓ | ✓ |
| EM1114567-017 19-DEC-2011 15:00 X4S | EM1114567-015 | 19-DEC-2011 15:00 | C4S | ✓ | ✓ |
| EM1114567-018 19-DEC-2011 15:00 Y4S | EM1114567-016 | 19-DEC-2011 15:00 | D4S | ✓ | ✓ |
| EM1114567-019 19-DEC-2011 15:00 AaS | EM1114567-017 | 19-DEC-2011 15:00 | X4S | ✓ | ✓ |
| EM1114567-020 19-DEC-2011 15:00 BbS | EM1114567-018 | 19-DEC-2011 15:00 | Y4S | ✓ | ✓ |
| EM1114567-021 19-DEC-2011 15:00 CcS | EM1114567-019 | 19-DEC-2011 15:00 | AaS | ✓ | ✓ |
| EM1114567-022 19-DEC-2011 15:00 DdS | EM1114567-020 | 19-DEC-2011 15:00 | BbS | ✓ | ✓ |
| EM1114567-023 19-DEC-2011 15:00 A5S ✓ ✓ EM1114567-024 19-DEC-2011 15:00 B5S ✓ ✓ EM1114567-025 19-DEC-2011 15:00 C5S ✓ ✓ EM1114567-026 19-DEC-2011 15:00 D5S ✓ ✓ EM1114567-027 19-DEC-2011 15:00 A6S ✓ ✓ EM1114567-028 19-DEC-2011 15:00 B6S ✓ ✓ EM1114567-029 19-DEC-2011 15:00 C6S ✓ ✓ EM1114567-030 19-DEC-2011 15:00 D6S ✓ ✓ EM1114567-031 19-DEC-2011 15:00 B1B ✓ ✓ EM1114567-033 19-DEC-2011 15:00 C1B ✓ ✓ EM1114567-034 19-DEC-2011 15:00 D1B ✓ ✓ | EM1114567-021 | 19-DEC-2011 15:00 | CcS | ✓ | ✓ |
| EM1114567-024 19-DEC-2011 15:00 B5S ✓ ✓ EM1114567-025 19-DEC-2011 15:00 C5S ✓ ✓ EM1114567-026 19-DEC-2011 15:00 D5S ✓ ✓ EM1114567-027 19-DEC-2011 15:00 A6S ✓ ✓ EM1114567-028 19-DEC-2011 15:00 B6S ✓ ✓ EM1114567-029 19-DEC-2011 15:00 C6S ✓ ✓ EM1114567-030 19-DEC-2011 15:00 D6S ✓ ✓ EM1114567-031 19-DEC-2011 15:00 A1B ✓ ✓ EM1114567-032 19-DEC-2011 15:00 B1B ✓ ✓ EM1114567-034 19-DEC-2011 15:00 D1B ✓ ✓ | EM1114567-022 | 19-DEC-2011 15:00 | DdS | ✓ | ✓ |
| EM1114567-025 19-DEC-2011 15:00 C5S ✓ ✓ EM1114567-026 19-DEC-2011 15:00 D5S ✓ ✓ EM1114567-027 19-DEC-2011 15:00 A6S ✓ ✓ EM1114567-028 19-DEC-2011 15:00 B6S ✓ ✓ EM1114567-029 19-DEC-2011 15:00 C6S ✓ ✓ EM1114567-030 19-DEC-2011 15:00 D6S ✓ ✓ EM1114567-031 19-DEC-2011 15:00 A1B ✓ ✓ EM1114567-032 19-DEC-2011 15:00 B1B ✓ ✓ EM1114567-033 19-DEC-2011 15:00 C1B ✓ ✓ EM1114567-034 19-DEC-2011 15:00 D1B ✓ ✓ | EM1114567-023 | 19-DEC-2011 15:00 | A5S | ✓ | ✓ |
| EM1114567-026 19-DEC-2011 15:00 D5S ✓ ✓ EM1114567-027 19-DEC-2011 15:00 A6S ✓ ✓ EM1114567-028 19-DEC-2011 15:00 B6S ✓ ✓ EM1114567-029 19-DEC-2011 15:00 C6S ✓ ✓ EM1114567-030 19-DEC-2011 15:00 D6S ✓ ✓ EM1114567-031 19-DEC-2011 15:00 A1B ✓ ✓ EM1114567-032 19-DEC-2011 15:00 B1B ✓ ✓ EM1114567-033 19-DEC-2011 15:00 C1B ✓ ✓ EM1114567-034 19-DEC-2011 15:00 D1B ✓ ✓ | EM1114567-024 | 19-DEC-2011 15:00 | B5S | ✓ | ✓ |
| EM1114567-027 19-DEC-2011 15:00 A6S | EM1114567-025 | 19-DEC-2011 15:00 | C5S | ✓ | ✓ |
| EM1114567-028 19-DEC-2011 15:00 B6S | EM1114567-026 | 19-DEC-2011 15:00 | D5S | ✓ | ✓ |
| EM1114567-029 19-DEC-2011 15:00 C6S ✓ ✓ EM1114567-030 19-DEC-2011 15:00 D6S ✓ ✓ EM1114567-031 19-DEC-2011 15:00 A1B ✓ ✓ EM1114567-032 19-DEC-2011 15:00 B1B ✓ ✓ EM1114567-033 19-DEC-2011 15:00 C1B ✓ ✓ EM1114567-034 19-DEC-2011 15:00 D1B ✓ ✓ | EM1114567-027 | 19-DEC-2011 15:00 | A6S | ✓ | ✓ |
| EM1114567-030 19-DEC-2011 15:00 D6S ✓ ✓ EM1114567-031 19-DEC-2011 15:00 A1B ✓ ✓ EM1114567-032 19-DEC-2011 15:00 B1B ✓ ✓ EM1114567-033 19-DEC-2011 15:00 C1B ✓ ✓ EM1114567-034 19-DEC-2011 15:00 D1B ✓ ✓ | EM1114567-028 | 19-DEC-2011 15:00 | B6S | ✓ | ✓ |
| EM1114567-031 19-DEC-2011 15:00 A1B | EM1114567-029 | 19-DEC-2011 15:00 | C6S | ✓ | ✓ |
| EM1114567-032 19-DEC-2011 15:00 B1B | EM1114567-030 | 19-DEC-2011 15:00 | D6S | ✓ | ✓ |
| EM1114567-033 19-DEC-2011 15:00 C1B | EM1114567-031 | 19-DEC-2011 15:00 | A1B | ✓ | ✓ |
| EM1114567-034 19-DEC-2011 15:00 D1B 🗸 🗸 | EM1114567-032 | 19-DEC-2011 15:00 | B1B | ✓ | ✓ |
| | EM1114567-033 | 19-DEC-2011 15:00 | C1B | ✓ | ✓ |
| EM1114567-035 19-DEC-2011 15:00 A2B | EM1114567-034 | 19-DEC-2011 15:00 | D1B | ✓ | ✓ |
| | EM1114567-035 | 19-DEC-2011 15:00 | A2B | ✓ | ✓ |

Issue Date : 30-MAR-2012

Page : 3 of 5
Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| | | | WATER - NT-08 Total Nitrogen + NO2 + NO3 + NH3 + Total P | WATER - W-03T 13 Metals (Total) (NEPM) |
|---------------|-------------------|-----|--|---|
| EM1114567-036 | 19-DEC-2011 15:00 | B2B | ✓ | ✓ |
| EM1114567-037 | 19-DEC-2011 15:00 | C2B | ✓ | ✓ |
| EM1114567-038 | 19-DEC-2011 15:00 | D2B | ✓ | ✓ |
| EM1114567-039 | 19-DEC-2011 15:00 | АЗВ | ✓ | ✓ |
| EM1114567-040 | 19-DEC-2011 15:00 | ВЗВ | ✓ | ✓ |
| EM1114567-041 | 19-DEC-2011 15:00 | СЗВ | ✓ | ✓ |
| EM1114567-042 | 19-DEC-2011 15:00 | D3B | ✓ | ✓ |
| EM1114567-043 | 19-DEC-2011 15:00 | A4B | ✓ | ✓ |
| EM1114567-044 | 19-DEC-2011 15:00 | B4B | ✓ | ✓ |
| EM1114567-045 | 19-DEC-2011 15:00 | C4B | ✓ | ✓ |
| EM1114567-046 | 19-DEC-2011 15:00 | D4B | ✓ | ✓ |
| EM1114567-047 | 19-DEC-2011 15:00 | X4B | ✓ | ✓ |
| EM1114567-048 | 19-DEC-2011 15:00 | Y4B | ✓ | ✓ |
| EM1114567-049 | 19-DEC-2011 15:00 | AaB | ✓ | ✓ |
| EM1114567-050 | 19-DEC-2011 15:00 | BbB | ✓ | ✓ |
| EM1114567-051 | 19-DEC-2011 15:00 | СсВ | ✓ | ✓ |
| EM1114567-052 | 19-DEC-2011 15:00 | DdB | ✓ | ✓ |
| EM1114567-053 | 19-DEC-2011 15:00 | A5B | ✓ | ✓ |
| EM1114567-054 | 19-DEC-2011 15:00 | B5B | ✓ | ✓ |
| EM1114567-055 | 19-DEC-2011 15:00 | C5B | ✓ | ✓ |
| EM1114567-056 | 19-DEC-2011 15:00 | D5B | ✓ | ✓ |
| EM1114567-057 | 19-DEC-2011 15:00 | A6B | ✓ | ✓ |
| EM1114567-058 | 19-DEC-2011 15:00 | B6B | ✓ | ✓ |
| EM1114567-059 | 19-DEC-2011 15:00 | C6B | ✓ | ✓ |
| EM1114567-060 | 19-DEC-2011 15:00 | D6B | ✓ | ✓ |

Issue Date : 30-MAR-2012
Page : 4 of 5

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



Proactive Holding Time Report

The following table summarises breaches of recommended holding times that have occurred prior to samples/instructions being received at the laboratory.

Matrix: WATER

Evaluation: **x** = Holding time breach ; ✓ = Within holding time.

| Matrix: WATER | | Dun for | Description | | | breach; ✓ = W | s Received |
|---------------------|--------------------------------|--------------------|---------------------|-------------|------------|---------------|------------|
| Method | | Due for extraction | Due for analysis | Samples R | | | |
| Client Sample ID(s) | Container | CAtraction | unuiyoio | Date | Evaluation | Date | Evaluation |
| | N by Discrete Analyser | | | | | | |
| A1B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | x | | |
| A1S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | 3¢ | | |
| A2B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | 3¢ | | |
| A2S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| A3B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| A3S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | ¥ € | | |
| A4B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| A4S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | JC . | | |
| A5B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | 3c | | |
| A5S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | JC . | | |
| A6B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| A6S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | 3C | | |
| AaB | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| AaS | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| B1B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| B1S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| B2B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| B2S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| B3B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | | | |
| B3S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | * | | |
| B4B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | * | | |
| B4S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | * | | |
| B5B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | * | | |
| B5S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | * | | |
| | | | | | * | | |
| B6B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| B6S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| BbB | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 |)x | | |
| BbS | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | JK . | | |
| C1B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | x | | |
| C1S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| C2B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | x | | |
| C2S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| C3B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| C3S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | * | | |
| C4B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| C4S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | JC . | | |
| C5B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | 3¢ | | |
| C5S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| C6B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| C6S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | ¥ € | | |
| CcB | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| CcS | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | 3¢ | | |
| D1B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | 3¢ | | |
| D1S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | x | | |
| D2B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | 3¢ | | |
| D2S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| D3B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| D3S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| D4B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| D4S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| D5B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| D5S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| D6B | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | | | |
| D6S | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| DdB | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | × | | |
| DdS | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | x | | |
| X4B | | | | | * | | + |
| ∧ 4 D | Clear Plastic Bottle - Natural | 21-DEC-2011 | | 22-DEC-2011 | . | | |

Issue Date : 30-MAR-2012

: 5 of 5

Page Work Order : EM1114567 Amendment 2 Client : SINCLAIR KNIGHT MERZ



| X4S | Clear Plastic Bottle - Natural | 21-DEC-2011 | 22-DEC-2011 | × | |
|-----|--------------------------------|-------------|-----------------|---|------|
| Y4B | Clear Plastic Bottle - Natural | 21-DEC-2011 | 22-DEC-2011 | × | |
| Y4S | Clear Plastic Bottle - Natural | 21-DEC-2011 | 22-DEC-2011 | × | |

Requested Deliverables

| ALL INVOICES | | |
|---|-------|-------------------------------|
| - A4 - AU Tax Invoice (INV) | Email | CLM-Adelaide@skm.com.au |
| ALL REPORTS | | |
| - *AU Certificate of Analysis - NATA | Email | CLM-Adelaide@globalskm.com.au |
| *AU Interpretive QC Report - DEFAULT (Anon QCI Rep) | Email | CLM-Adelaide@globalskm.com.au |
| - *AU QC Report - DEFAULT (Anon QC Rep) - NATA | Email | CLM-Adelaide@globalskm.com.au |
| - A4 - AU Sample Receipt Notification - Environmental HT | Email | CLM-Adelaide@globalskm.com.au |
| - A4 - AU Tax Invoice | Email | CLM-Adelaide@globalskm.com.au |
| - Chain of Custody (CoC) | Email | CLM-Adelaide@globalskm.com.au |
| - EDI Format - ENMRG | Email | CLM-Adelaide@globalskm.com.au |
| - EDI Format - ESDAT | Email | CLM-Adelaide@globalskm.com.au |
| - EDI Format - XTab | Email | CLM-Adelaide@globalskm.com.au |
| DR GREG BARBARA | | |
| - *AU Certificate of Analysis - NATA | Email | gbarbara@globalskm.com |
| *AU Interpretive QC Report - DEFAULT (Anon QCI Rep) | Email | gbarbara@globalskm.com |
| - *AU QC Report - DEFAULT (Anon QC Rep) - NATA | Email | gbarbara@globalskm.com |
| - A4 - AU Sample Receipt Notification - Environmental HT | Email | gbarbara@globalskm.com |
| - A4 - AU Tax Invoice | Email | gbarbara@globalskm.com |
| - Chain of Custody (CoC) | Email | gbarbara@globalskm.com |
| - EDI Format - ENMRG | Email | gbarbara@globalskm.com |
| - EDI Format - ESDAT | Email | gbarbara@globalskm.com |
| - EDI Format - XTab | Email | gbarbara@globalskm.com |
| GREG BARBARA | | |
| - *AU Certificate of Analysis - NATA (COA) | Email | gbarbara@skm.com.au |
| - *AU Interpretive QC Report - DEFAULT (Anon QCI Rep) (QCI) | Email | gbarbara@skm.com.au |
| - *AU QC Report - DEFAULT (Anon QC Rep) - NATA (QC) | Email | gbarbara@skm.com.au |
| - A4 - AU Sample Receipt Notification - Environmental HT (SRN) | Email | gbarbara@skm.com.au |
| - A4 - AU Tax Invoice (INV) | Email | gbarbara@skm.com.au |
| - Chain of Custody (CoC) (COC) | Email | gbarbara@skm.com.au |
| - EDI Format - ENMRG (ENMRG) | Email | gbarbara@skm.com.au |
| - EDI Format - ESDAT (ESDAT) | Email | gbarbara@skm.com.au |
| - EDI Format - XTab (XTAB) | Email | gbarbara@skm.com.au |
| | | |

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

INTERPRETIVE QUALITY CONTROL REPORT

Work Order : **EM1114567** Page : 1 of 12

Amendment : 2

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : DR GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

Telephone : +61 08 8424 3800 Telephone : +61-3-8549 9608 Facsimile : +61 08 8424 3810 Facsimile : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

Site : ----

 C-O-C number
 : --- Date Samples Received
 : 22-DEC-2011

 Sampler
 : GB
 Issue Date
 : 30-MAR-2012

Order number : ----

Quote number : ADBQ/037/11 No. of samples analysed : 60

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for release.

This Interpretive Quality Control Report contains the following information:

- Analysis Holding Time Compliance
- Quality Control Parameter Frequency Compliance
- Brief Method Summaries
- Summary of Outliers

Page : 2 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Analysis Holding Time Compliance

The following report summarises extraction / preparation and analysis times and compares with recommended holding times. Dates reported represent first date of extraction or analysis and precludes subsequent dilutions and reruns. Information is also provided re the sample container (preservative) from which the analysis aliquot was taken. Elapsed period to analysis represents number of days from sampling where no extraction / digestion is involved or period from extraction / digestion where this is present. For composite samples, sampling date is assumed to be that of the oldest sample contributing to the composite. Sample date for laboratory produced leachates is assumed as the completion date of the leaching process. Outliers for holding time are based on USEPA SW 846, APHA, AS and NEPM (1999). A listing of breaches is provided in the Summary of Outliers.

Holding times for leachate methods (excluding elutriates) vary according to the analytes being determined on the resulting solution. For non-volatile analytes, the holding time compliance assessment compares the leach date with the shortest analyte holding time for the equivalent soil method. These soil holding times are: Organics (14 days); Mercury (28 days) & other metals (180 days). A recorded breach therefore does not guarantee a breach for all non-volatile parameters.

Matrix: WATER Evaluation: ▼ = Holding time breach; ✓ = Within holding time.

| Method | Sample Date | Ex | traction / Preparation | | | Analysis | |
|--|-------------|----------------|------------------------|------------|---------------|------------------|------------|
| Container / Client Sample ID(s) | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EG020T: Total Metals by ICP-MS | | | | | | | |
| Clear Plastic Bottle - Unfiltered; Lab-acidified | | | | | | | |
| A1S, B1S, | 19-DEC-2011 | 23-DEC-2011 | 16-JUN-2012 | 1 | 30-DEC-2011 | 16-JUN-2012 | ✓ |
| C1S, D1S, | | | | | | | |
| A2S, B2S, | | | | | | | |
| C2S, D2S, | | | | | | | |
| A3S, B3S, | | | | | | | |
| C3S, D3S, | | | | | | | |
| A4S, B4S, | | | | | | | |
| C4S, D4S, | | | | | | | |
| X4S, Y4S, | | | | | | | |
| AaS, BbS, | | | | | | | |
| CcS, DdS, | | | | | | | |
| A5S, B5S, | | | | | | | |
| C5S, D5S, | | | | | | | |
| A6S, B6S, | | | | | | | |
| C6S, D6S, | | | | | | | |
| A1B, B1B, | | | | | | | |
| C1B, D1B, | | | | | | | |
| A2B, B2B, | | | | | | | |
| C2B, D2B, | | | | | | | |
| A3B, B3B | | | | | | | |
| Clear Plastic Bottle - Unfiltered; Lab-acidified | | | | | | | |
| C3B, D3B, | 19-DEC-2011 | 28-DEC-2011 | 16-JUN-2012 | 1 | 30-DEC-2011 | 16-JUN-2012 | ✓ |
| A4B, B4B, | | | | | | | |
| C4B, D4B, | | | | | | | |
| X4B, Y4B, | | | | | | | |
| AaB, BbB, | | | | | | | |
| CcB, DdB, | | | | | | | |
| A5B, B5B, | | | | | | | |
| C5B, D5B, | | | | | | | |
| A6B, B6B, | | | | | | | |
| C6B, D6B | | | | | | | |

Page : 3 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Matrix: WATER | | | | | Evaluation | : x = Holding time | breach ; ✓ = Within | n holding time |
|-----------------------------------|------------------|--------------------------|----------------|--------------------|------------|--------------------|---------------------|----------------|
| Method | Sample Date | Extraction / Preparation | | | Analysis | | | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EG035T: Total Recoverable Me | ercury by FIMS | | | | | | | |
| Clear Plastic Bottle - Unfiltered | d; Lab-acidified | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | | | 05-JAN-2012 | 16-JAN-2012 | 1 |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 4 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Matrix: WATER Evaluation: × = Holding time breach ; ✓ = W | | | | | | | breach ; ✓ = Within | in holding time |
|---|----------------|-------------|--------------------------|--------------------|------------|---------------|---------------------|-----------------|
| Method Container / Client Sample ID(s) | | Sample Date | Extraction / Preparation | | | Analysis | | |
| | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EK055G: Ammonia as N by Disc | crete Analyser | | | | | | | |
| Clear Plastic Bottle - Sulfuric A | Acid | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | 16-JAN-2012 | | 06-JAN-2012 | 16-JAN-2012 | 1 |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 5 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| Matrix: WATER | | | | | Evaluation: | x = Holding time | breach ; ✓ = Withir | holding tin |
|----------------------------------|----------|-------------|----------------|------------------------|-------------|------------------|---------------------|-------------|
| Method | | Sample Date | E) | traction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EK057G: Nitrite as N by Discrete | Analyser | | | | | | | |
| Clear Plastic Bottle - Natural | | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | 21-DEC-2011 | | 22-DEC-2011 | 21-DEC-2011 | * |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 6 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| Matrix: WATER | | | | | Evaluation: | × = Holding time | breach ; ✓ = Withir | n holding time |
|---------------------------------|---------------------------------|-------------|----------------|------------------------|-------------|------------------|---------------------|----------------|
| Method | | Sample Date | Ex | traction / Preparation | | | | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EK059G: Nitrite plus Nitrate | as N (NOx) by Discrete Analyser | | | | | | | |
| Clear Plastic Bottle - Sulfuri | ic Acid | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | | 16-JAN-2012 | | 05-JAN-2012 | 16-JAN-2012 | ✓ |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B, | | | | | | | |
| C3B, | D3B, | | | | | | | |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |

Page : 7 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| Matrix: WATER | | | | | Evaluation | : × = Holding time | breach ; ✓ = Within | n holding time. |
|------------------------------------|----------------------|-------------|----------------|------------------------|------------|--------------------|---------------------|-----------------|
| Method | | Sample Date | Ex | traction / Preparation | | | Analysis | |
| Container / Client Sample ID(s) | | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation |
| EK061G: Total Kjeldahl Nitrogen | By Discrete Analyser | | | | | | | |
| Clear Plastic Bottle - Sulfuric Ad | cid | | | | | | | |
| C3B, | D3B, | 19-DEC-2011 | 03-JAN-2012 | 16-JAN-2012 | 1 | 03-JAN-2012 | 16-JAN-2012 | 1 |
| A4B, | B4B, | | | | | | | |
| C4B, | D4B, | | | | | | | |
| X4B, | Y4B, | | | | | | | |
| AaB, | BbB, | | | | | | | |
| CcB, | DdB, | | | | | | | |
| A5B, | B5B, | | | | | | | |
| C5B, | D5B, | | | | | | | |
| A6B, | B6B, | | | | | | | |
| C6B, | D6B | | | | | | | |
| Clear Plastic Bottle - Sulfuric Ad | cid | | | | | | | |
| A1S, | B1S, | 19-DEC-2011 | 30-DEC-2011 | 16-JAN-2012 | 1 | 30-DEC-2011 | 16-JAN-2012 | 1 |
| C1S, | D1S, | | | | | | | |
| A2S, | B2S, | | | | | | | |
| C2S, | D2S, | | | | | | | |
| A3S, | B3S, | | | | | | | |
| C3S, | D3S, | | | | | | | |
| A4S, | B4S, | | | | | | | |
| C4S, | D4S, | | | | | | | |
| X4S, | Y4S, | | | | | | | |
| AaS, | BbS, | | | | | | | |
| CcS, | DdS, | | | | | | | |
| A5S, | B5S, | | | | | | | |
| C5S, | D5S, | | | | | | | |
| A6S, | B6S, | | | | | | | |
| C6S, | D6S, | | | | | | | |
| A1B, | B1B, | | | | | | | |
| C1B, | D1B, | | | | | | | |
| A2B, | B2B, | | | | | | | |
| C2B, | D2B, | | | | | | | |
| A3B, | B3B | | | | | | | |

Page : 8 of 12

Matrix: WATER

C5S,

A6S, C6S,

A1B,

C1B,

A2B,

C2B,

A3B,

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

D5S, B6S,

D6S,

B1B,

D1B,

B2B,

D2B,

B3B

Project : VE23610 609



Evaluation: **x** = Holding time breach ; ✓ = Within holding time.

| Method | Sample Date | Ex | traction / Preparation | | | Analysis | | |
|--|-------------|----------------|------------------------|------------|---------------|------------------|------------|--|
| Container / Client Sample ID(s) | | Date extracted | Due for extraction | Evaluation | Date analysed | Due for analysis | Evaluation | |
| EK067G: Total Phosphorus as P by Discrete Analyser | | | | | | | | |
| Clear Plastic Bottle - Sulfuric Acid | | | | | | | | |
| C3B, D3B, | 19-DEC-2011 | 03-JAN-2012 | 16-JAN-2012 | 1 | 03-JAN-2012 | 16-JAN-2012 | ✓ | |
| A4B, B4B, | | | | | | | | |
| C4B, D4B, | | | | | | | | |
| X4B, Y4B, | | | | | | | | |
| AaB, BbB, | | | | | | | | |
| CcB, DdB, | | | | | | | | |
| A5B, B5B, | | | | | | | | |
| C5B, D5B, | | | | | | | | |
| A6B, B6B, | | | | | | | | |
| C6B, D6B | | | | | | | | |
| Clear Plastic Bottle - Sulfuric Acid | | | | | | | | |
| A1S, B1S, | 19-DEC-2011 | 30-DEC-2011 | 16-JAN-2012 | 1 | 30-DEC-2011 | 16-JAN-2012 | ✓ | |
| C1S, D1S, | | | | | | | | |
| A2S, B2S, | | | | | | | | |
| C2S, D2S, | | | | | | | | |
| A3S, B3S, | | | | | | | | |
| C3S, D3S, | | | | | | | | |
| A4S, B4S, | | | | | | | | |
| C4S, D4S, | | | | | | | | |
| X4S, Y4S, | | | | | | | | |
| AaS, BbS, | | | | | | | | |
| CcS, DdS, | | | | | | | | |
| A5S, B5S, | | | | | | | | |

Page : 9 of 12

Matrix: WATER

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609

Total Phosphorus as P By Discrete Analyser

Nitrite and Nitrate as N (NOx) by Discrete Analyser

Total Kjeldahl Nitrogen as N By Discrete Analyser

Total Phosphorus as P By Discrete Analyser

Ammonia as N by Discrete analyser

Nitrite as N by Discrete Analyser

Total Metals by ICP-MS - Suite A

Matrix Spikes (MS)

Total Mercury by FIMS



Evaluation: × = Quality Control frequency not within specification: ✓ = Quality Control frequency within specification.

NEPM 1999 Schedule B(3) and ALS QCS3 requirement

Quality Control Parameter Frequency Compliance

The following report summarises the frequency of laboratory QC samples analysed within the analytical lot(s) in which the submitted sample(s) was(where) processed. Actual rate should be greater than or equal to the expected rate. A listing of breaches is provided in the Summary of Outliers.

EK067G

EK055G

EK059G

EK057G

EK061G

EG035T

EK067G

EG020A-T

3

3

3

3

3

3

3

3

60

60

60

60

60

60

60

60

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

5.0

1

✓

1

Quality Control Sample Type Quality Control Specification Count Rate (%) Method Evaluation Analytical Methods QC Regular Actual Expected Laboratory Duplicates (DUP) Ammonia as N by Discrete analyser EK055G 6 60 10.0 10.0 1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Nitrite and Nitrate as N (NOx) by Discrete Analyser 60 EK059G 6 10.0 10.0 ✓ NEPM 1999 Schedule B(3) and ALS QCS3 requirement Nitrite as N by Discrete Analyser EK057G 6 60 10.0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 Suspended Solids (High Level) EA025H 2 11 18.2 10.0 1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Kjeldahl Nitrogen as N By Discrete Analyser 60 EK061G 6 10 0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 Total Mercury by FIMS EG035T 6 60 10.0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Metals by ICP-MS - Suite A EG020A-T 6 60 10.0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Phosphorus as P By Discrete Analyser EK067G 6 60 10.0 10.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Laboratory Control Samples (LCS) Ammonia as N by Discrete analyser EK055G 3 60 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 5.0 5.0 Nitrite and Nitrate as N (NOx) by Discrete Analyser 3 60 5.0 EK059G 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 Nitrite as N by Discrete Analyser EK057G 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Suspended Solids (High Level) FA025H 1 11 9.1 5.0 1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Kjeldahl Nitrogen as N By Discrete Analyser EK061G 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1 3 Total Mercury by FIMS EG035T 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement ✓ Total Metals by ICP-MS - Suite A EG020A-T 3 60 5.0 5.0 1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Phosphorus as P By Discrete Analyser EK067G 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Method Blanks (MB) 3 Ammonia as N by Discrete analyser EK055G 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 3 60 Nitrite and Nitrate as N (NOx) by Discrete Analyser EK059G 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 3 Nitrite as N by Discrete Analyser EK057G 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement EA025H Suspended Solids (High Level) 1 11 9.1 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 5.0 1 Total Kjeldahl Nitrogen as N By Discrete Analyser 3 EK061G 60 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 5.0 Total Mercury by FIMS EG035T 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement Total Metals by ICP-MS - Suite A EG020A-T 3 60 5.0 5.0 NEPM 1999 Schedule B(3) and ALS QCS3 requirement 1

Page : 10 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Brief Method Summaries

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the US EPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request. The following report provides brief descriptions of the analytical procedures employed for results reported in the Certificate of Analysis. Sources from which ALS methods have been developed are provided within the Method Descriptions.

| Analytical Methods | Method | Matrix | Method Descriptions |
|---|-------------|--------|---|
| Suspended Solids (High Level) | EA025H | WATER | APHA 21st ed., 2540D A gravimetric procedure employed to determine the amount of `non-filterable` residue in a aqueous sample. The prescribed GFC (1.2um) filter is rinsed with deionised water, oven dried and weighed prior to analysis. A well-mixed sample is filtered through a glass fibre filter (1.2um). The residue on the filter paper is dried at 104+/-2C. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Total Metals by ICP-MS - Suite A | EG020A-T | WATER | (APHA 21st ed., 3125; USEPA SW846 - 6020, ALS QWI-EN/EG020): The ICPMS technique utilizes a highly efficient argon plasma to ionize selected elements. Ions are then passed into a high vacuum mass spectrometer, which separates the analytes based on their distinct mass to charge ratios prior to their measurement by a discrete dynode ion detector. |
| Total Mercury by FIMS | EG035T | WATER | AS 3550, APHA 21st ed. 3112 Hg - B (Flow-injection (SnCl2)(Cold Vapour generation) AAS) FIM-AAS is an automated flameless atomic absorption technique. A bromate/bromide reagent is used to oxidise any organic mercury compounds in the unfiltered sample. The ionic mercury is reduced online to atomic mercury vapour by SnCl2 which is then purged into a heated quartz cell. Quantification is by comparing absorbance against a calibration curve. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Ammonia as N by Discrete analyser | EK055G | WATER | APHA 21st ed., 4500-NH3 G Ammonia is determined by direct colorimetry by Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Nitrite as N by Discrete Analyser | EK057G | WATER | APHA 21st ed., 4500-NO2- B. Nitrite is determined by direct colourimetry by Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Nitrate as N by Discrete Analyser | EK058G | WATER | APHA 21st ed., 4500-NO3- F. Nitrate is reduced to nitrite by way of a cadmium reduction column followed by quantification by Discrete Analyser. Nitrite is determined seperately by direct colourimetry and result for Nitrate calculated as the difference between the two results. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Nitrite and Nitrate as N (NOx) by Discrete Analyser | EK059G | WATER | APHA 21st ed., 4500-NO3- F. Combined oxidised Nitrogen (NO2+NO3) is determined by Cadmium Reduction and direct colourimetry by Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Total Kjeldahl Nitrogen as N By Discrete Analyser | EK061G | WATER | APHA 21st ed., 4500-Norg D. 25mL water samples are digested using a traditional Kjeldahl digestion followed by determination by Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Total Nitrogen as N (TKN + Nox) By Discrete Analyser | EK062G | WATER | APHA 21st ed., 4500-Norg / 4500-NO3 This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Total Phosphorus as P By Discrete Analyser | EK067G | WATER | APHA 21st ed., 4500-P B&F This procedure involves sulphuric acid digestion of a 100mL sample to break phosphorus down to orthophosphate. The orthophosphate reacts with ammonium molybdate and antimony potassium tartrate to form a complex which is then reduced and its concentration measured at 880nm using Discrete Analyser. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Preparation Methods | Method | Matrix | Method Descriptions |
| TKN/TP Digestion | EK061/EK067 | WATER | APHA 21st ed., 4500 Norg - D; APHA 21st ed., 4500 P - H. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |
| Digestion for Total Recoverable Metals | EN25 | WATER | USEPA SW846-3005 Method 3005 is a Nitric/Hydrochloric acid digestion procedure used to prepare surface and ground water samples for analysis by ICPAES or ICPMS. This method is compliant with NEPM (1999) Schedule B(3) (Appdx. 2) |

Page : 11 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Summary of Outliers

Outliers: Quality Control Samples

The following report highlights outliers flagged in the Quality Control (QC) Report. Surrogate recovery limits are static and based on USEPA SW846 or ALS-QWI/EN/38 (in the absence of specific USEPA limits). This report displays QC Outliers (breaches) only.

Duplicates, Method Blanks, Laboratory Control Samples and Matrix Spikes

- For all matrices, no Method Blank value outliers occur.
- For all matrices, no Duplicate outliers occur.
- For all matrices, no Laboratory Control outliers occur.
- For all matrices, no Matrix Spike outliers occur.

Regular Sample Surrogates

• For all regular sample matrices, no surrogate recovery outliers occur.

Outliers: Analysis Holding Time Compliance

This report displays Holding Time breaches only. Only the respective Extraction / Preparation and/or Analysis component is/are displayed.

Matrix: WATER

| Method Method | Ex | traction / Preparation | | Analysis | | |
|---|----------------|------------------------|---------|---------------|------------------|---------|
| Container / Client Sample ID(s) | Date extracted | Due for extraction | Days | Date analysed | Due for analysis | Days |
| | | | overdue | | | overdue |
| EK057G: Nitrite as N by Discrete Analyser | | | | | | |

Page : 12 of 12

Work Order : EM1114567 Amendment 2 Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Matrix: WATER

| Method | | E | extraction / Preparation | | | Analysis | |
|---------------------------------------|---|----------------|--------------------------|-----------------|---------------|------------------|-----------------|
| Container / Client Sample ID(s) | | Date extracted | Due for extraction | Days overdue | Date analysed | Due for analysis | Days overdue |
| EK057G: Nitrite as N by Discrete Anal | yser - Analysis Holding Time Compliance | | | | | | |
| Clear Plastic Bottle - Natural | | | | | | | |
| A1S, | B1S, | | | | 22-DEC-2011 | 21-DEC-2011 | 1 |
| C1S, | D1S, | | | | | | |
| A2S, | B2S, | | | | | | |
| C2S, | D2S, | | | | | | |
| A3S, | B3S, | | | | | | |
| C3S, | D3S, | | | | | | |
| A4S, | B4S, | | | | | | |
| C4S, | D4S, | | | | | | |
| X4S, | Y4S, | | | | | | |
| AaS, | BbS, | | | | | | |
| CcS, | DdS, | | | | | | |
| A5S, | B5S, | | | | | | |
| C5S, | D5S, | | | | | | |
| A6S, | B6S, | | | | | | |
| C6S, | D6S, | | | | | | |
| A1B, | B1B, | | | | | | |
| C1B, | D1B, | | | | | | |
| A2B, | B2B, | | | | | | |
| C2B, | D2B, | | | | | | |
| A3B, | B3B, | | | | | | |
| C3B, | D3B, | | | | | | |
| A4B, | B4B, | | | | | | |
| C4B, | D4B, | | | | | | |
| X4B, | Y4B, | | | | | | |
| AaB, | BbB, | | | | | | |
| CcB, | DdB, | | | | | | |
| A5B, | B5B, | | | | | | |
| C5B, | D5B, | | | | | | |
| A6B, | B6B, | | | | | | |
| C6B, | D6B | | | | | | |

Outliers : Frequency of Quality Control Samples

The following report highlights breaches in the Frequency of Quality Control Samples.

No Quality Control Sample Frequency Outliers exist.

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

QUALITY CONTROL REPORT

Work Order : **EM1114567** Page : 1 of 12

Amendment : 2

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : DR GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

 Telephone
 : +61 08 8424 3800
 Telephone
 : +61-3-8549 9608

 Facsimile
 : +61 08 8424 3810
 Facsimile
 : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

Site : ----

 C-O-C number
 : -- Date Samples Received
 : 22-DEC-2011

 Sampler
 : GB
 Issue Date
 : 30-MAR-2012

No. of samples received : 60

This report supercedes any prayious report(s) with this reference. Posults apply to the sample(s) as submitted. All pages of this report have been checked and appro-

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for release.

No. of samples analysed

This Quality Control Report contains the following information:

: ----

- Laboratory Duplicate (DUP) Report; Relative Percentage Difference (RPD) and Acceptance Limits
- Method Blank (MB) and Laboratory Control Spike (LCS) Report; Recovery and Acceptance Limits
- Matrix Spike (MS) Report; Recovery and Acceptance Limits



Order number

Quote number

NATA Accredited Laboratory 825

: ADBQ/037/11

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

: 60

| Signatories | Position | Accreditation Category |
|----------------|------------------------|------------------------|
| Eric Chau | Metals Team Leader | Melbourne Inorganics |
| Herman Lin | Laboratory Coordinator | Melbourne Inorganics |
| Varsha Ho Wing | Non-Metals Team Leader | Melbourne Inorganics |

Page : 2 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

Key: Anonymous = Refers to samples which are not specifically part of this work order but formed part of the QC process lot

CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

RPD = Relative Percentage Difference

= Indicates failed QC

Page : 3 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Laboratory Duplicate (DUP) Report

The quality control term Laboratory Duplicate refers to a randomly selected intralaboratory split. Laboratory duplicates provide information regarding method precision and sample heterogeneity. The permitted ranges for the Relative Percent Deviation (RPD) of Laboratory Duplicates are specified in ALS Method QWI-EN/38 and are dependent on the magnitude of results in comparison to the level of reporting: Result < 10 times LOR:-No Limit; Result between 10 and 20 times LOR:-0% - 50%; Result > 20 times LOR:-0% - 20%.

| ub-Matrix: WATER | | | | Laboratory Duplicate (DUP) Report | | | | | | | |
|----------------------|-----------------------|---------------------|------------|-----------------------------------|------|-----------------|------------------|---------|--------------------|--|--|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (% | | |
| G020T: Total Meta | Is by ICP-MS (QC Lot: | 2105258) | | | | | | | | | |
| EM1114567-001 | A1S | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.0 | No Limit | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.008 | 0.009 | 21.7 | No Limit | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.001 | 0.002 | 0.0 | No Limit | | |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.005 | 0.0 | No Limit | | |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.0 | No Limit | | |
| M1114567-010 | B3S | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.0 | No Limit | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.009 | 0.0 | No Limit | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.003 | 0.001 | 64.5 | No Limit | | |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.0 | No Limit | | |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.0 | No Limit | | |
| G020T: Total Meta | Is by ICP-MS (QC Lot: | 2105259) | | | | | | | | | |
| M1114567-021 | CcS | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit | | |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.0 | No Limit | | |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.0 | 0% - 50% | | |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit | | |

Page : 4 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| ub-Matrix: WATER | | | | | | t | | | |
|---------------------|-----------------------|----------------------|------------|--------|------|-----------------|------------------|---------|---------------------|
| aboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) |
| G020T: Total Meta | Is by ICP-MS (QC Lot: | 2105259) - continued | | | | | | | |
| M1114567-021 | CcS | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.0 | No Limit |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.0 | No Limit |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.0 | No Limit |
| M1114567-030 | D6S | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.0 | No Limit |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.0 | 0% - 50% |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.0 | No Limit |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.0 | No Limit |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.0 | No Limit |
| G020T: Total Meta | Is by ICP-MS (QC Lot: | | | | | | | | |
| M1114566-001 | Anonymous | EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | 0.0001 | 0.0001 | 0.0 | No Limit |
| | | EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | 0.013 | 0.014 | 11.3 | 0% - 50% |
| | | EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | 0.058 | 0.061 | 5.0 | 0% - 20% |
| | | EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | 0.002 | 0.002 | 0.0 | No Limit |
| | | EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | 0.001 | 0.001 | 0.0 | No Limit |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.017 | 0.017 | 0.0 | 0% - 50% |
| | | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | 0.015 | 0.016 | 7.4 | 0% - 50% |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | 0.082 | 0.087 | 5.1 | 0% - 20% |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.016 | 0.017 | 0.0 | 0% - 50% |
| | | EG020A-T: Nickel | 7440-66-6 | 0.005 | mg/L | 0.163 | 0.187 | 13.7 | 0% - 20% |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | 0.06 | 0.06 | 0.0 | No Limit |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | 1.67 | 1.71 | 2.4 | 0% - 20% |
| M1114567-050 | BbB | EG020A-T: IIOTI | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.0001 | 0.0 | No Limit |
| WITT 14307-030 | БОБ | EG020A-T: Cadmium | 7440-38-2 | 0.0001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Arsenic | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | | 7440-39-3 | 0.001 | mg/L | 0.005 | 0.006 | 0.0 | No Limit |
| | | EG020A-T: Chromium | 7440-39-3 | 0.001 | mg/L | <0.003 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Coholium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Cobalt | 7440-40-4 | | - | 0.010 | 0.001 | 0.0 | 0% - 50% |
| | | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.0 | 0% - 50% |

Page : 5 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | | | | Laboratory I | Duplicate (DUP) Report | t | |
|--------------------------------|------------------------|---|------------|--------|--------|-----------------|------------------------|---------|---------------------|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) |
| EG020T: Total Meta | Is by ICP-MS (QC Lot: | 2107796) - continued | | | | | | | |
| EM1114567-050 | BbB | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | 0.0 | No Limit |
| | | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | 0.005 | 0.005 | 0.0 | No Limit |
| | | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.009 | 32.8 | No Limit |
| | | EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| | | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | 0.06 | 0.10 | 42.2 | No Limit |
| EG035T: Total Reco | overable Mercury by F | IMS (QC Lot: 2115265) | | | | | | | |
| EM1114567-001 | A1S | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit |
| EM1114567-010 | B3S | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | 0.0007 | 151 | No Limit |
| EG035T: Total Reco | overable Mercury by F | IMS (QC Lot: 2115266) | | | | | | | |
| EM1114567-021 | CcS | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit |
| EM1114567-030 | D6S | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit |
| FG035T: Total Reco | overable Mercury by Fl | IMS (QC Lot: 2115267) | | | | | | | |
| EM1114567-041 | СЗВ | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit |
| EM1114567-050 | BbB | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0 | No Limit |
| EK055G: Ammonia | as N by Discrete Analy | /ser (QC Lot: 2115294) | | | 3 | | | | |
| EM1114567-001 | A1S | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.10 | 0.0 | 0% - 50% |
| EM1114567-010 | B3S | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.11 | 10.1 | 0% - 50% |
| | | /ser (QC Lot: 2115297) | 1001111 | 0.01 | 9.2 | 02 | U | | 070 0070 |
| EM1114567-021 | CcS | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.11 | 0.10 | 0.0 | 0% - 50% |
| EM1114567-030 | D6S | EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.09 | 0.0 | No Limit |
| | | | 7001111 | 0.01 | mg/L | 0.10 | 0.00 | 0.0 | 110 Emilio |
| EM1114567-041 | C3B | /ser (QC Lot: 2115299) | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.15 | 16.8 | 0% - 50% |
| EM1114567-050 | BbB | EK055G: Ammonia as N EK055G: Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.13 | 16.8 | 0% - 50% |
| | | | 7004-41-7 | 0.01 | IIIg/L | 0.11 | 0.13 | 10.0 | 0 /0 - 30 /0 |
| | N by Discrete Analyse | | | 0.04 | | 10.04 | 10.01 | 0.0 | NIa I insit |
| EM1114567-001 EM1114567-010 | A1S B3S | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 <0.01 | <0.01 <0.01 | 0.0 | No Limit |
| | | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| | N by Discrete Analyse | | | | | | | | |
| EM1114567-021 | CcS | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-030 | D6S | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| | N by Discrete Analyse | | | | | | | _ | |
| EM1114567-041 | C3B | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-050 | BbB | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EK059G: Nitrite plu | s Nitrate as N (NOx) b | y Discrete Analyser (QC Lot: 2115293) | | | | | | | |
| EM1114567-001 | A1S | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EM1114567-010 | B3S | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |
| EK059G: Nitrite plu | s Nitrate as N (NOx) b | y Discrete Analyser (QC Lot: 2115296) | | | | | | | |
| EM1114567-021 | CcS | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit |

Page : 6 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | | Laboratory Duplicate (DUP) Report | | | | | | | |
|----------------------|--------------------------|---|------------|-----------------------------------|------|-----------------|------------------|---------|---------------------|--|--|
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | LOR | Unit | Original Result | Duplicate Result | RPD (%) | Recovery Limits (%) | | |
| EK059G: Nitrite plu | s Nitrate as N (NOx) by | Discrete Analyser (QC Lot: 2115296) - continued | | | | | | | | | |
| EM1114567-030 | D6S | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| EK059G: Nitrite plu | s Nitrate as N (NOx) by | Discrete Analyser (QC Lot: 2115298) | | | | | | | | | |
| EM1114567-041 | C3B | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| EM1114567-050 | BbB | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| EK061G: Total Kjeld | ahl Nitrogen By Discrete | e Analyser (QC Lot: 2109542) | | | | | | | | | |
| EM1114567-001 | A1S | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.3 | 0.0 | No Limit | | |
| EM1114567-010 | B3S | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.2 | 0.0 | No Limit | | |
| EK061G: Total Kjeld | ahl Nitrogen By Discrete | e Analyser (QC Lot: 2109544) | | | | | | | | | |
| EM1114567-021 | CcS | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.1 | 0.0 | No Limit | | |
| EM1114567-030 | D6S | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.3 | 0.2 | 52.8 | No Limit | | |
| EK061G: Total Kjeld | ahl Nitrogen By Discrete | e Analyser (QC Lot: 2110810) | | | | | | | | | |
| EM1114567-041 | C3B | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | 0.2 | 0.0 | No Limit | | |
| EM1114567-050 | BbB | EK061G: Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | 0.0 | No Limit | | |
| EK067G: Total Phos | phorus as P by Discrete | Analyser (QC Lot: 2109543) | | | | | | | | | |
| EM1114567-001 | A1S | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| EM1114567-010 | B3S | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.01 | 0.0 | No Limit | | |
| EK067G: Total Phos | phorus as P by Discrete | Analyser (QC Lot: 2109545) | | | | | | | | | |
| EM1114567-021 | CcS | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |
| EM1114567-030 | D6S | EK067G: Total Phosphorus as P | | 0.01 | mg/L | 0.05 | 0.04 | 29.9 | No Limit | | |
| EK067G: Total Phos | phorus as P by Discrete | Analyser (QC Lot: 2110811) | | | | | | | | | |
| EM1114567-041 | C3B | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.01 | 0.0 | No Limit | | |
| EM1114567-050 | BbB | EK067G: Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | 0.0 | No Limit | | |

Page : 7 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Method Blank (MB) and Laboratory Control Spike (LCS) Report

The quality control term Method / Laboratory Blank refers to an analyte free matrix to which all reagents are added in the same volumes or proportions as used in standard sample preparation. The purpose of this QC parameter is to monitor potential laboratory contamination. The quality control term Laboratory Control Sample (LCS) refers to a certified reference material, or a known interference free matrix spiked with target analytes. The purpose of this QC parameter is to monitor method precision and accuracy independent of sample matrix. Dynamic Recovery Limits are based on statistical evaluation of processed LCS.

| Sub-Matrix: WATER | | | | Method Blank (MB) | | Laboratory Control Spike (LCS |) Report | |
|---|------------|--------|------|-------------------|---------------|-------------------------------|----------|------------|
| | | | | Report | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| Method: Compound | CAS Number | LOR | Unit | Result | Concentration | LCS | Low | High |
| EG020T: Total Metals by ICP-MS (QCLot: 2105258) | | | | | | | | |
| EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 106 | 86 | 110 |
| EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 68 | 128 |
| EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 87 | 115 |
| EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.1 mg/L | 102 | 87 | 111 |
| EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 103 | 87 | 115 |
| EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 101 | 88 | 114 |
| EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 88 | 110 |
| EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 102 | 90 | 114 |
| EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 87 | 113 |
| EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 107 | 89 | 113 |
| EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | 0.1 mg/L | 103 | 86 | 114 |
| EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | 0.1 mg/L | 104 | 82 | 116 |
| EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.5 mg/L | 101 | 81 | 125 |
| EG020T: Total Metals by ICP-MS (QCLot: 2105259) | | | | | | | | |
| EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 102 | 86 | 110 |
| EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 124 | 68 | 128 |
| EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 101 | 87 | 115 |
| EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.1 mg/L | 98.9 | 87 | 111 |
| EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 87 | 115 |
| EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 102 | 88 | 114 |
| EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 102 | 88 | 110 |
| EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 99.2 | 90 | 114 |
| EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 100 | 87 | 113 |
| EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 105 | 89 | 113 |
| EG020A-T: Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | 0.1 mg/L | 108 | 86 | 114 |
| EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | 0.1 mg/L | 101 | 82 | 116 |
| EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.5 mg/L | 112 | 81 | 125 |
| EG020T: Total Metals by ICP-MS (QCLot: 2107796) | | | | | | | | |
| EG020A-T: Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 110 | 86 | 110 |
| EG020A-T: Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 105 | 68 | 128 |
| EG020A-T: Barium | 7440-39-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 110 | 87 | 115 |
| EG020A-T: Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.1 mg/L | 103 | 87 | 111 |
| EG020A-T: Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 104 | 87 | 115 |
| EG020A-T: Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 107 | 88 | 114 |

Page : 8 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| | Sub-Matrix: WATER | | | | Method Blank (MB) | | Laboratory Control Spike (LCS |) Report | |
|--|---|----------------------|--------|--------|-------------------|---------------|-------------------------------|----------|------------|
| E00201** Total Metals by ICP-MS** (QCLot: 2107786) - continued C00201** Copper** | | | | | Report | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| EXCAPAT - Copper | Method: Compound | CAS Number | LOR | Unit | Result | Concentration | LCS | Low | High |
| FERDING Table Ta | EG020T: Total Metals by ICP-MS (QCLot: 2107796) | continued | | | | | | | |
| FERRORAN Managamese 7439-86-5 0.001 mgt. <0.001 0.1 mgt. 106 87 113 118 | EG020A-T: Copper | 7440-50-8 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 108 | 88 | 110 |
| FRODOR - Nuckel FA40-02-0 0.001 mg/L <0.001 0.1 mg/L 109 89 113 | EG020A-T: Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 107 | 90 | 114 |
| FRODUCAT: Variation | EG020A-T: Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 108 | 87 | 113 |
| FERDIZANT ZINC | EG020A-T: Nickel | 7440-02-0 | 0.001 | mg/L | <0.001 | 0.1 mg/L | 109 | 89 | 113 |
| Feedback Free Property Films Free Property Films Free Property Films Free Property Free Prop | EG020A-T: Vanadium | | 0.01 | mg/L | <0.01 | 0.1 mg/L | 103 | 86 | 114 |
| EG0351: Total Recoverable Mercury by FIMS (QCLot: 2115265) EG0351: Mercury 7439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 85.6 69 125 EG0351: Mercury 7439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 85.6 69 125 EG0351: Mercury 7439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 85.6 69 125 EG0351: Mercury 7439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 85.6 69 125 EK05351: Mercury 7439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 86.6 69 125 EK05351: Mercury 7439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 80.5 69 125 EK05531: Mercury 7439-97-6 0.0001 mg/L <0.001 0.5 mg/L 113 76 122 EK05561: Ammonia as N by Discrete Analyser (QCLot: 2115294) EK05563: Ammonia as N by Discrete Analyser (QCLot: 2115297) EK05563: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05563: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05563: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05563: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05563: Mirrite as N by Discrete Analyser (QCLot: 2115293) EK056563: Mirrite as N by Discrete Analyser (QCLot: 2115293) EK056763: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05763: Nitrite as N by Discrete Analyser (QCLot: 2105293) EK05763: Nitrite as N by Discrete Analyser (QCLot: 2105293) EK05763: Nitrite as N by Discrete Analyser (QCLot: 2105293) EK05763: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05763: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05763: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05663: Nitrite Pilitate as N (NOx) by Discrete Analyser (QCLot: 2115293) EK05663: Nitrite Pilitate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05663: Nitrite Pilitate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05663: Nitrite Pilitate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05663: Nitrite Pilitate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05663: Nitrite Pilitate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK06663: Total Kjeldahi Nitrogen By Discrete Analyser (QCLot: 210542) EK0616: Total Kjeldahi Nitrogen By Discrete Analyser (QCLot: 210542) | EG020A-T: Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | 0.1 mg/L | 110 | 82 | 116 |
| EG035T: Marcury T439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 89.2 69 125 | EG020A-T: Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.5 mg/L | 108 | 81 | 125 |
| EG935T: Total Recoverable Mercury by FIMS (QCLot: 2115266) EG935T: Mercury 7439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 85.6 69 125 EG935T: Mercury 7439-97-6 0.0001 mg/L <0.0001 0.0100 mg/L 80.5 69 125 EK095G: Ammonia as N by Discrete Analyser (QCLot: 2115294) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115297) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115297) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05GS: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK05GS: Mirite as N by Discrete Analyser (QCLot: 2115293) EK05GS: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05GS: Nitrite as N by Discrete Analyser (QCLot: 2115294) EK05GS: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05GS: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05GS: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05GS: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK05GS: Nitrite Pius Pius Pius Pius Pius Pius Pius Pius | EG035T: Total Recoverable Mercury by FIMS (QCLot | t: 2115265) | | | | | | | |
| EG035T: Mercury 7439-97-8 0.0001 mg/L <0.0001 0.0100 mg/L 85.6 69 125 | EG035T: Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | 0.0100 mg/L | 89.2 | 69 | 125 |
| EG035T: Mercury 7439-97-8 0.0001 mg/L <0.0001 0.0100 mg/L 85.6 69 125 | EG035T: Total Recoverable Mercury by FIMS (QCLot | t: 2115266) | | | | | | | |
| EG035T: Mercury | EG035T: Mercury | | 0.0001 | mg/L | <0.0001 | 0.0100 mg/L | 85.6 | 69 | 125 |
| EG035T: Mercury | EG035T: Total Recoverable Mercury by FIMS (QCLot | t: 2115267) | | | | | | | |
| EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115294) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115297) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115297) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115297) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115298) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2105293) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105296) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2115296) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2115296) EK057G: Nitrite as N lirate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK057G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analy | | | 0.0001 | mg/L | <0.0001 | 0.0100 mg/L | 80.5 | 69 | 125 |
| EK055G: Ammonia as N | | r· 2115294) | | | | | | | |
| EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115297) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2105293) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105293) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2105542) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2105544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2105544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2105544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2105544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2105544) | | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 113 | 76 | 122 |
| EK055G: Ammonia as N | | | | | | J. J. J. | | - | |
| EK055G: Ammonia as N by Discrete Analyser (QCLot: 2115299) EK055G: Ammonia as N by Discrete Analyser (QCLot: 2105233) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105234) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105234) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105234) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105234) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105235) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105235) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105235) EK057G: Nitrite plus Nitrate as N by Discrete Analyser (QCLot: 2105235) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115293) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahi Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahi Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahi Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahi Nitrogen By Discrete Analyser (QCLot: 2109544) | | | 0.01 | mg/l | <0.01 | 0.5 mg/l | 114 | 76 | 122 |
| EK055G: Ammonia as N 7684-41-7 0.01 mg/L <0.01 0.5 mg/L 110 76 122 EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105293) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2115293) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115293) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK061G: Total Kjeldahl Nitrogen as N | | | 0.0. | 9.2 | | 0.0 mg/2 | | | |
| EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105293) EK057G: Nitrite as N | | | 0.01 | ma/l | <0.01 | 0.5 mg/l | 110 | 76 | 122 |
| EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115293) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 210542) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) | | | 0.01 | IIIg/L | 40.01 | 0.5 mg/L | 110 | 70 | 122 |
| EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105294) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115293) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109542) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) | , | | 0.01 | ma/l | <0.01 | 0.5 mg/l | 100 | 0.4 | 110 |
| EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115293) EK059G: Nitrite Plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite + Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite + Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite + Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite + Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109542) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <-0.1 10 mg/L 10 mg/L 10 6 63 117 | | | 0.01 | mg/L | <0.01 | 0.5 Hig/L | 100 | 04 | 112 |
| EK057G: Nitrite as N by Discrete Analyser (QCLot: 2105295) EK057G: Nitrite as N | | | 2.24 | | 0.04 | 0.5 " | 00.4 | | 110 |
| EK057G: Nitrite as N 0.01 mg/L <0.01 0.5 mg/L 99.7 84 112 EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115293) EK059G: Nitrite + Nitrate as N 0.01 mg/L <0.01 0.5 mg/L 95.0 73 127 EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite + Nitrate as N 0.01 mg/L <0.01 0.5 mg/L 104 73 127 EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite + Nitrate as N 0.01 mg/L <0.01 0.5 mg/L 101 73 127 EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109542) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 99.1 | 84 | 112 |
| EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115293) EK059G: Nitrite + Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite + Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite + Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite + Nitrate as N (NOx) by Discrete Analyser (QCLot: 2109542) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) | | | | | | | | | |
| EK059G: Nitrite + Nitrate as N | EK057G: Nitrite as N | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 99.7 | 84 | 112 |
| EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115296) EK059G: Nitrite + Nitrate as N | EK059G: Nitrite plus Nitrate as N (NOx) by Discrete A | Analyser (QCLot: 211 | 5293) | | | | | | |
| EK059G: Nitrite + Nitrate as N | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 95.0 | 73 | 127 |
| EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser (QCLot: 2115298) EK059G: Nitrite + Nitrate as N | EK059G: Nitrite plus Nitrate as N (NOx) by Discrete A | Analyser (QCLot: 211 | 5296) | | | | | | |
| EK059G: Nitrite + Nitrate as N 0.01 mg/L <0.01 0.5 mg/L 101 73 127 EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109542) EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <0.1 10 mg/L 104 63 117 EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <0.1 10 mg/L 106 63 117 | EK059G: Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | 0.5 mg/L | 104 | 73 | 127 |
| EK059G: Nitrite + Nitrate as N 0.01 mg/L <0.01 0.5 mg/L 101 73 127 EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109542) EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <0.1 10 mg/L 104 63 117 EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <0.1 10 mg/L 106 63 117 | EK059G: Nitrite plus Nitrate as N (NOx) by Discrete A | Analyser (QCLot: 211 | 5298) | | | | | | |
| EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <0.1 | EK059G: Nitrite + Nitrate as N | | | mg/L | <0.01 | 0.5 mg/L | 101 | 73 | 127 |
| EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <0.1 | EK061G: Total Kieldahl Nitrogen By Discrete Analyse | r (QCLot: 2109542) | | | | | | | |
| EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 2109544) EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <0.1 10 mg/L 106 63 117 | | | 0.1 | mg/L | <0.1 | 10 mg/L | 104 | 63 | 117 |
| EK061G: Total Kjeldahl Nitrogen as N 0.1 mg/L <0.1 10 mg/L 106 63 117 | | r (OCL of: 2109544) | | | | | | | |
| The control of the co | | | 0.1 | ma/L | <0.1 | 10 ma/L | 106 | 63 | 117 |
| | | r (OCL et: 3440940) | | | | | | | |

Page : 9 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | Method Blank (MB) | | Laboratory Control Spike (LC | S) Report | |
|---|---------------|------|-------------------|---------------|------------------------------|-----------|------------|
| | | | Report | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| Method: Compound CAS Number | LOR | Unit | Result | Concentration | LCS | Low | High |
| EK061G: Total Kjeldahl Nitrogen By Discrete Analyser (QCLot: 211081 |) - continued | | | | | | |
| EK061G: Total Kjeldahl Nitrogen as N | 0.1 | mg/L | <0.1 | 10 mg/L | 87.1 | 63 | 117 |
| EK067G: Total Phosphorus as P by Discrete Analyser (QCLot: 2109543 |) | | | | | | |
| EK067G: Total Phosphorus as P | 0.01 | mg/L | <0.01 | 4.42 mg/L | 99.5 | 73 | 117 |
| EK067G: Total Phosphorus as P by Discrete Analyser (QCLot: 2109545 |) | | | | | | |
| EK067G: Total Phosphorus as P | 0.01 | mg/L | <0.01 | 4.42 mg/L | 89.7 | 73 | 117 |
| EK067G: Total Phosphorus as P by Discrete Analyser (QCLot: 2110811 |) | | | | | | |
| EK067G: Total Phosphorus as P | 0.01 | mg/L | <0.01 | 4.42 mg/L | 98.0 | 73 | 117 |

Page : 10 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



Matrix Spike (MS) Report

The quality control term Matrix Spike (MS) refers to an intralaboratory split sample spiked with a representative set of target analytes. The purpose of this QC parameter is to monitor potential matrix effects on analyte recoveries. Static Recovery Limits as per laboratory Data Quality Objectives (DQOs). Ideal recovery ranges stated may be waived in the event of sample matrix interference.

| ub-Matrix: WATER | | | Γ | | Matrix Spike (MS) Repo | rt | |
|----------------------|------------------------------|---------------------|------------|---------------|------------------------|----------|------------|
| | | | | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | Concentration | MS | Low | High |
| G020T: Total Metal | s by ICP-MS (QCLot: 2105258) | | | | | | |
| EM1114567-001 | A1S | EG020A-T: Arsenic | 7440-38-2 | 1 mg/L | 110 | 72 | 146 |
| | | EG020A-T: Beryllium | 7440-41-7 | 1 mg/L | 85.8 | 61 | 139 |
| | | EG020A-T: Barium | 7440-39-3 | 1 mg/L | 98.8 | 78 | 126 |
| | | EG020A-T: Cadmium | 7440-43-9 | 0.25 mg/L | 83.3 | 73 | 131 |
| | | EG020A-T: Chromium | 7440-47-3 | 1 mg/L | 85.1 | 65 | 131 |
| | | EG020A-T: Cobalt | 7440-48-4 | 1 mg/L | 101 | 68 | 132 |
| | | EG020A-T: Copper | 7440-50-8 | 1 mg/L | 84.6 | 71 | 125 |
| | | EG020A-T: Lead | 7439-92-1 | 1 mg/L | 95.5 | 68 | 130 |
| | | EG020A-T: Manganese | 7439-96-5 | 1 mg/L | 88.8 | 63 | 129 |
| | | EG020A-T: Nickel | 7440-02-0 | 1 mg/L | 89.5 | 72 | 128 |
| | | EG020A-T: Vanadium | 7440-62-2 | 1 mg/L | 95.6 | 66 | 130 |
| | | EG020A-T: Zinc | 7440-66-6 | 1 mg/L | 79.4 | 67 | 129 |
| G020T: Total Metal | s by ICP-MS (QCLot: 2105259) | | | | | | |
| M1114567-021 | CcS | EG020A-T: Arsenic | 7440-38-2 | 1 mg/L | 104 | 72 | 146 |
| | 114001 021 | EG020A-T: Beryllium | 7440-41-7 | 1 mg/L | 94.9 | 61 | 139 |
| | | EG020A-T: Barium | 7440-39-3 | 1 mg/L | 109 | 78 | 126 |
| | | EG020A-T: Cadmium | 7440-43-9 | 0.25 mg/L | 93.9 | 73 | 131 |
| | | EG020A-T: Chromium | 7440-47-3 | 1 mg/L | 105 | 65 | 131 |
| | | EG020A-T: Cobalt | 7440-48-4 | 1 mg/L | 111 | 68 | 132 |
| | | EG020A-T: Copper | 7440-50-8 | 1 mg/L | 93.9 | 71 | 125 |
| | | EG020A-T: Lead | 7439-92-1 | 1 mg/L | 105 | 68 | 130 |
| | | EG020A-T: Manganese | 7439-96-5 | 1 mg/L | 99.6 | 63 | 129 |
| | | EG020A-T: Nickel | 7440-02-0 | 1 mg/L | 101 | 72 | 128 |
| | | EG020A-T: Vanadium | 7440-62-2 | 1 mg/L | 110 | 66 | 130 |
| | | EG020A-T: Zinc | 7440-66-6 | 1 mg/L | 88.4 | 67 | 129 |
| G020T: Total Metal | s by ICP-MS (QCLot: 2107796) | | | | | | |
| M1114567-041 | C3B | EG020A-T: Arsenic | 7440-38-2 | 1 mg/L | 117 | 72 | 146 |
| | | EG020A-T: Beryllium | 7440-41-7 | 1 mg/L | 94.9 | 61 | 139 |
| | | EG020A-T: Barium | 7440-39-3 | 1 mg/L | 120 | 78 | 126 |
| | | EG020A-T: Cadmium | 7440-43-9 | 0.25 mg/L | 104 | 73 | 131 |
| | | EG020A-T: Chromium | 7440-47-3 | 1 mg/L | 112 | 65 | 131 |
| | | EG020A-T: Cobalt | 7440-48-4 | 1 mg/L | 125 | 68 | 132 |
| | | EG020A-T: Copper | 7440-50-8 | 1 mg/L | 101 | 71 | 125 |
| | | EG020A-T: Lead | 7439-92-1 | 1 mg/L | 112 | 68 | 130 |
| | | EG020A-T: Manganese | 7439-96-5 | 1 mg/L | 108 | 63 | 129 |

Page : 11 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | | | Matrix Spike (MS) Repo | rt | |
|---------------------------------------|--------------------------------------|--------------------------------------|------------|---------------|------------------------|----------|------------|
| | | | | Spike | Spike Recovery (%) | Recovery | Limits (%) |
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | Concentration | MS | Low | High |
| G020T: Total Metal | s by ICP-MS (QCLot: 2107796) | - continued | | | | | |
| EM1114567-041 | C3B | EG020A-T: Nickel | 7440-02-0 | 1 mg/L | 110 | 72 | 128 |
| | | EG020A-T: Vanadium | 7440-62-2 | 1 mg/L | 118 | 66 | 130 |
| | | EG020A-T: Zinc | 7440-66-6 | 1 mg/L | 99.9 | 67 | 129 |
| G035T: Total Reco | verable Mercury by FIMS (QC | Lot: 2115265) | | | | | |
| EM1114567-002 | B1S | EG035T: Mercury | 7439-97-6 | 0.0100 mg/L | 84.8 | 70 | 130 |
| G035T: Total Reco | verable Mercury by FIMS (QC | Lot: 2115266) | | | | | |
| EM1114567-022 | DdS | EG035T: Mercury | 7439-97-6 | 0.0100 mg/L | 84.9 | 70 | 130 |
| G035T: Total Reco | verable Mercury by FIMS (QC | Lot: 2115267) | | | | | |
| EM1114567-042 | D3B | EG035T: Mercury | 7439-97-6 | 0.0100 mg/L | 82.0 | 70 | 130 |
| K055G: Ammonia a | s N by Discrete Analyser (QC | | | | | | |
| EM1114567-002 | B1S | EK055G: Ammonia as N | 7664-41-7 | 0.5 mg/L | 94.4 | 70 | 130 |
| K055G: Ammonia | s N by Discrete Analyser (QC | | | | | | |
| EM1114567-022 | DdS | EK055G: Ammonia as N | 7664-41-7 | 0.5 mg/L | 98.3 | 70 | 130 |
| | is N by Discrete Analyser (QC | | | | 77.7 | | |
| EM1114567-042 | D3B | EK055G: Ammonia as N | 7664-41-7 | 0.5 mg/L | 90.3 | 70 | 130 |
| | N by Discrete Analyser (QCLo | | 7.00.7.17. | 0.0g/_ | 55.5 | | |
| EM1114567-002 | B1S | EK057G: Nitrite as N | | 0.5 mg/L | 118 | 70 | 130 |
| | N by Discrete Analyser (QCLo | | | 0.0 mg/L | 110 | 70 | 100 |
| EM1114567-022 | DdS | EK057G: Nitrite as N | | 0.5 mg/L | 118 | 70 | 130 |
| | | | | 0.5 Hig/L | 110 | 70 | 130 |
| :K057G: Nitrite as I EM1114567-042 | N by Discrete Analyser (QCLor D3B | | | 0.5/ | 115 | 70 | 400 |
| | | EK057G: Nitrite as N | | 0.5 mg/L | 115 | 70 | 130 |
| | Nitrate as N (NOx) by Discre | | | 0.5 " | 440 | | 100 |
| EM1114567-002 | B1S | EK059G: Nitrite + Nitrate as N | | 0.5 mg/L | 119 | 70 | 130 |
| | Nitrate as N (NOx) by Discre | te Analyser (QCLot: 2115296) | | | | | |
| EM1114567-022 | DdS | EK059G: Nitrite + Nitrate as N | | 0.6 mg/L | 110 | 70 | 130 |
| | s Nitrate as N (NOx) by Discre | te Analyser (QCLot: 2115298) | | | | | |
| EM1114567-042 | D3B | EK059G: Nitrite + Nitrate as N | | 0.5 mg/L | 117 | 70 | 130 |
| K061G: Total Kjeld | ahl Nitrogen By Discrete Analy | vser (QCLot: 2109542) | | | | | |
| EM1114567-002 | B1S | EK061G: Total Kjeldahl Nitrogen as N | | 25 mg/L | 107 | 70 | 130 |
| K061G: Total Kjeld | ahl Nitrogen By Discrete Analy | vser (QCLot: 2109544) | | | | | |
| M1114567-022 | DdS | EK061G: Total Kjeldahl Nitrogen as N | | 25 mg/L | 76.7 | 70 | 130 |
| K061G: Total Kjeld | ahl Nitrogen By Discrete Analy | vser (QCLot: 2110810) | | | | | |
| EM1114567-042 | D3B | EK061G: Total Kjeldahl Nitrogen as N | | 25 mg/L | 102 | 70 | 130 |
| K067G: Total Phos | phorus as P by Discrete Analy | | | | | | |
| EM1114567-002 | B1S | EK067G: Total Phosphorus as P | | 5 mg/L | 79.7 | 70 | 130 |

Page : 12 of 12

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ



| Sub-Matrix: WATER | | | | | Matrix Spike (MS) Repo | ort | |
|----------------------|---|-------------------------------|------------|---------------|------------------------|------------|------------|
| | | | | Spike | Spike Recovery (%) | Recovery I | Limits (%) |
| Laboratory sample ID | Client sample ID | Method: Compound | CAS Number | Concentration | MS | Low | High |
| EK067G: Total Phos | phorus as P by Discrete Analyser(QCLot: 2 | 2109545) | | | | | |
| EM1114567-022 | DdS | EK067G: Total Phosphorus as P | | 5 mg/L | 88.0 | 70 | 130 |
| EK067G: Total Phos | phorus as P by Discrete Analyser(QCLot: 2 | 2110811) | | | | | |
| EM1114567-042 | D3B | EK067G: Total Phosphorus as P | | 5 mg/L | 95.2 | 70 | 130 |

ALS Laboratory Group

ANALYTICAL CHEMISTRY & TESTING SERVICES



Environmental Division

CERTIFICATE OF ANALYSIS

Work Order : **EM1114567** Page : 1 of 14

Amendment : 2

Client : SINCLAIR KNIGHT MERZ Laboratory : Environmental Division Melbourne

Contact : DR GREG BARBARA Contact : Carol Walsh

Address : LEVEL 5, 33 KING WILLIAM ST Address : 4 Westall Rd Springvale VIC Australia 3171

ADELAIDE SA, AUSTRALIA 5000

 Telephone
 : +61 08 8424 3800
 Telephone
 : +61-3-8549 9608

 Facsimile
 : +61 08 8424 3810
 Facsimile
 : +61-3-8549 9601

Project : VE23610 609 QC Level : NEPM 1999 Schedule B(3) and ALS QCS3 requirement

Order number : ----

 C-O-C number
 : -- Date Samples Received
 : 22-DEC-2011

 Sampler
 : GB
 Issue Date
 : 30-MAR-2012

Site : ----

This report supersedes any previous report(s) with this reference. Results apply to the sample(s) as submitted. All pages of this report have been checked and approved for release.

This Certificate of Analysis contains the following information:

- General Comments
- Analytical Results



NATA Accredited Laboratory 825

Accredited for compliance with ISO/IEC 17025.

Signatories

This document has been electronically signed by the authorized signatories indicated below. Electronic signing has been carried out in compliance with procedures specified in 21 CFR Part 11.

| Signatories | Position | Accreditation Category |
|----------------|------------------------|------------------------|
| Eric Chau | Metals Team Leader | Melbourne Inorganics |
| Herman Lin | Laboratory Coordinator | Melbourne Inorganics |
| Varsha Ho Wing | Non-Metals Team Leader | Melbourne Inorganics |

Page : 2 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



General Comments

The analytical procedures used by the Environmental Division have been developed from established internationally recognized procedures such as those published by the USEPA, APHA, AS and NEPM. In house developed procedures are employed in the absence of documented standards or by client request.

Where moisture determination has been performed, results are reported on a dry weight basis.

Where a reported less than (<) result is higher than the LOR, this may be due to primary sample extract/digestate dilution and/or insufficient sample for analysis.

Where the LOR of a reported result differs from standard LOR, this may be due to high moisture content, insufficient sample (reduced weight employed) or matrix interference.

When sampling time information is not provided by the client, sampling dates are shown without a time component. In these instances, the time component has been assumed by the laboratory for processing purposes.

Key: CAS Number = CAS registry number from database maintained by Chemical Abstracts Services. The Chemical Abstracts Service is a division of the American Chemical Society.

LOR = Limit of reporting

^ = This result is computed from individual analyte detections at or above the level of reporting

- 27/2/12 This report has been amended and re-released to allow the reporting of additional analytical data.
- 30/3/12 This report has been amended following the request to issue the report without Suspended Solids results.
- EA025: Total Suspended Solids has been conducted outside of holding times due to laboratory constraints. Results should be scrutinised accordingly.
- It is recognised that Total Kjeldahl Nitrogen is less than Ammonia for sample EM1114567. However, the difference is within experimental variation of the methods.

Page : 3 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Cli | ent sample ID | A1S | B1S | C1S | D1S | A2S |
|-----------------------------------|------------------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Cl | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-001 | EM1114567-002 | EM1114567-003 | EM1114567-004 | EM1114567-005 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | 0.0001 | 0.0001 | <0.0001 | 0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.008 | 0.020 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.001 | 0.002 | 0.003 | 0.002 | 0.002 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.022 | <0.005 | <0.005 | 0.006 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercu | ury by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discre | te Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.11 | 0.10 | 0.09 | 0.10 |
| EK057G: Nitrite as N by Discrete | Analyser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete | Analyser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N | (NOx) by Discrete Ana | lyser | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen B | By Discrete Analy <u>ser</u> | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.3 | 0.3 | 0.4 | 0.2 |
| EK062G: Total Nitrogen as N (TKN | l + NOx) by Discrete Ar | nalyser_ | | | | | | |
| Total Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.3 | 0.3 | 0.4 | 0.2 |
| EK067G: Total Phosphorus as P b | y Discrete Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | 0.03 | <0.01 | <0.01 |
| | | | - | | <u> </u> | 1 | | |

Page : 4 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Canal | Sub-Matrix: WATER | | Cli | ent sample ID | B2S | C2S | D2S | A3S | B3S |
|--|--------------------------------|---------------------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| Control Metals by ICP-MS | | CI | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Assenic 7440-38-2 0.001 mg/L 0.001 0.001 0.0001 0. | Compound | CAS Number | LOR | Unit | EM1114567-006 | EM1114567-007 | EM1114567-008 | EM1114567-009 | EM1114567-010 |
| Sartium | EG020T: Total Metals by ICP-I | MS | | | | | | | |
| Sery lium | Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | Barium | 7440-39-3 | 0.001 | mg/L | 0.007 | 0.006 | 0.006 | 0.006 | 0.006 |
| Cobalt 7440-48-4 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 | Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | 0.0001 | <0.0001 |
| Copper | Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Manganese | Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel 7440-02-0 0.001 mg/L 0.002 0.002 0.002 0.002 0.003 | Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.010 |
| Lead 7439-92:1 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.0001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0. | Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | Nickel | 7440-02-0 | 0.001 | mg/L | | | | | |
| Zinc 7440-66-6 0.005 mg/L < 0.005 < 0.005 | Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Iron | Vanadium | 7440-62-2 | 0.01 | mg/L | | | <0.01 | <0.01 | |
| Mercury 7439-97-6 0.0001 mg/L 0.0001 | Zinc | 7440-66-6 | 0.005 | mg/L | | <0.005 | | 0.006 | |
| Mercury 7439-97-6 0.0001 mg/L <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 | Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EK055G: Ammonia as N by Discrete Analyser Ammonia as N | EG035T: Total Recoverable M | lercury by FIMS | | | | | | | |
| Ammonia as N 7664-41-7 0.01 mg/L 0.11 0.10 0.10 0.10 0.11 0.12 EK057G: Nitrite as N by Discrete Analyser Nitrite as N | Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Nitrite as N by Discrete Analyser Nitrite as N by Discrete Analyser Nitrite as N N N N N N N N N N | EK055G: Ammonia as N by Di | screte Analyser | | | | | | | |
| Nitrite as N | Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.11 | 0.10 | 0.10 | 0.11 | 0.12 |
| Nitrate as N by Discrete Analyser Nitrate as N by Discrete Analyser | EK057G: Nitrite as N by Discr | ete Analyser | | | | | | | |
| Nitrate as N 14797-55-8 0.01 mg/L 0.02 0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.0 | Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser Nitrite + Nitrate as N | EK058G: Nitrate as N by Disc | rete Analyser | | | | | | | |
| Nitrite + Nitrate as N 0.01 mg/L 0.02 0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 | Nitrate as N | 14797-55-8 | 0.01 | mg/L | 0.02 | 0.01 | <0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen By Discrete Analyser Total Kjeldahl Nitrogen as N 0.1 mg/L 0.4 0.3 0.2 0.2 0.2 0.1 EK062G: Total Nitrogen as N (TKN + NOx) by Discrete Analyser Total Nitrogen as N 0.1 mg/L 0.4 0.3 0.2 0.2 0.2 0.1 EK067G: Total Phosphorus as P by Discrete Analyser | EK059G: Nitrite plus Nitrate a | s N (NOx) by Discrete Ana | lyser | | | | | | |
| Total Kjeldahl Nitrogen as N 0.1 mg/L 0.4 0.3 0.2 0.2 0.1 EK062G: Total Nitrogen as N (TKN + NOx) by Discrete Analyser Total Nitrogen as N 0.1 mg/L 0.4 0.3 0.2 0.2 0.2 0.1 EK067G: Total Phosphorus as P by Discrete Analyser 0.4 0.3 0.2 0.2 0.1 | Nitrite + Nitrate as N | | 0.01 | mg/L | 0.02 | 0.01 | <0.01 | <0.01 | <0.01 |
| EK062G: Total Nitrogen as N (TKN + NOx) by Discrete Analyser Total Nitrogen as N 0.1 mg/L 0.4 0.3 0.2 0.2 0.2 0.1 EK067G: Total Phosphorus as P by Discrete Analyser | EK061G: Total Kjeldahl Nitrog | en By Discrete Analys <u>er</u> | | | | | | | |
| Total Nitrogen as N 0.1 mg/L 0.4 0.3 0.2 0.2 0.2 0.1 EK067G: Total Phosphorus as P by Discrete Analyser | Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.4 | 0.3 | 0.2 | 0.2 | 0.1 |
| Total Nitrogen as N 0.1 mg/L 0.4 0.3 0.2 0.2 0.2 0.1 EK067G: Total Phosphorus as P by Discrete Analyser | EK062G: Total Nitrogen as N (| TKN + NOx) by Discrete Ar | nalyser | | | | | | |
| | - | | | mg/L | 0.4 | 0.3 | 0.2 | 0.2 | 0.1 |
| Total Phosphorus as P 0.01 mg/L <0.01 <0.01 <0.01 0.01 0.01 <0.01 | EK067G: Total Phosphorus as | P by Discrete Anal <u>yser</u> | | | | | | | |
| ··· ··· ··· · · · · · · · · · · · · · | Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | 0.01 | <0.01 |

Page : 5 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | C3S | D3S | A4S | B4S | C4S |
|---------------------------------------|---------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | CI | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-011 | EM1114567-012 | EM1114567-013 | EM1114567-014 | EM1114567-015 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.007 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | 0.0006 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.014 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | 0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.007 | <0.005 | <0.005 | <0.005 | 0.028 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercury | by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete | Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.09 | 0.10 | 0.10 | 0.10 |
| EK057G: Nitrite as N by Discrete An | alyser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete Ar | alyser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | 0.01 | <0.01 | 0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (No | Ox) by Discrete Ana | lyser | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | 0.01 | <0.01 | 0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen By | Discrete Analyser | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.2 | 0.2 | 0.6 | 0.2 |
| EK062G: Total Nitrogen as N (TKN + | NOx) by Discrete Ar | alyser | | | | | | |
| ^ Total Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.2 | 0.2 | 0.6 | 0.2 |
| EK067G: Total Phosphorus as P by I | Discrete Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | 0.01 | 0.05 | 0.03 | 0.03 | 0.04 |

Page : 6 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Arsenic 7440-36-2 0.001 mg/L 0.006 | Sub-Matrix: WATER | | Cli | ent sample ID | D4S | X4S | Y4S | AaS | BbS |
|--|--------------------------------------|-----------------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| Control Cont | | CI | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Arsenic 7440-98.2 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 < | Compound | CAS Number | LOR | Unit | EM1114567-016 | EM1114567-017 | EM1114567-018 | EM1114567-019 | EM1114567-020 |
| Arsenic 7440-98.2 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 < | EG020T: Total Metals by ICP-MS | | | | | | | | |
| Beryllium | Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmitum | Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Cobalt | Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium 7440-47-3 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 | Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Copper | Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Manganese 7439-96-5 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 | Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel 7440-02-0 0.001 mg/L 0.002 0.002 0.002 0.002 0.002 0.002 0.000 0.000 0.000 0.0001 0.00 | Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.011 | 0.010 | 0.010 | 0.010 |
| Lead 7439-92-1 0.001 mg/L <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.005 <0.0 | Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium 7440-62-2 0.01 mg/L < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.005 Value | Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| Zinc | Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| From 7439-89-6 0.05 mg/L < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.05 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.001 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.01 < 0.0 | Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Construction Cons | Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.131 | <0.005 | 0.010 | <0.005 |
| Mercury 7439-97-6 0.0001 mg/L <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.0001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.001 <0.0 | Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| Control Cont | EG035T: Total Recoverable Mercui | ry by FIMS | | | | | | | |
| Ammonia as N 7664-41-7 0.01 mg/L 0.10 0.09 0.08 0.10 0.10 EK057G: Nitrite as N by Discrete Analyser Nitrite as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 EK058G: Nitrate as N by Discrete Analyser Nitrate as N 14797-55-8 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 EK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser Nitrite + Nitrate as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 EK061G: Total Kjeldahl Nitrogen By Discrete Analyser Total Kjeldahl Nitrogen as N (TKN + NOx) by Discrete Analyser Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.2 0.2 EK067G: Total Nitrogen as N (TKN + NOx) by Discrete Analyser Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.2 0.2 0.2 EK067G: Total Phosphorus as P by Discrete Analyser | Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Counting | EK055G: Ammonia as N by Discret | e Analyser | | | | | | | |
| Nitrite as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0 | Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.09 | 0.08 | 0.10 | 0.10 |
| Nitrate as N by Discrete Analyser Sk058G: Nitrate as N by Discrete Analyser Sk059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser Nitrite + Nitrate as N (NOx) by Discrete Analyser Nitrite + Nitrate as N | EK057G: Nitrite as N by Discrete A | nalyser | | | | | | | |
| Nitrate as N | Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| CK059G: Nitrite plus Nitrate as N (NOx) by Discrete Analyser Nitrite + Nitrate as N | EK058G: Nitrate as N by Discrete A | Analyser | | | | | | | |
| Nitrite + Nitrate as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0 | Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Nitrite + Nitrate as N 0.01 mg/L <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0.01 <0 | EK059G: Nitrite plus Nitrate as N (l | NOx) by Discrete Ana | lyser | | | | | | |
| Total Kjeldahl Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.2 0.2 0.2 0.2 KK062G: Total Nitrogen as N (TKN + NOx) by Discrete Analyser Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.2 0.2 0.2 KK067G: Total Phosphorus as P by Discrete Analyser | Nitrite + Nitrate as N | | | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Total Kjeldahl Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.2 0.2 0.2 0.2 KK062G: Total Nitrogen as N (TKN + NOx) by Discrete Analyser Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.2 0.2 0.2 KK067G: Total Phosphorus as P by Discrete Analyser | EK061G: Total Kjeldahl Nitrogen By | y Discrete Analy <u>ser</u> | | | | | | | |
| Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.2 0.2 0.2 K067G: Total Phosphorus as P by Discrete Analyser | Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| Total Nitrogen as N 0.1 mg/L 0.2 0.2 0.2 0.2 0.2 0.2 0.2 K067G: Total Phosphorus as P by Discrete Analyser | EK062G: Total Nitrogen as N (TKN | + NOx) by Discrete Ar | nalyser | | | | | | |
| | ` Total Nitrogen as N | | | mg/L | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| Total Phosphorus as P 0.01 mg/l <0.01 0.01 0.04 0.02 0.02 | EK067G: Total Phosphorus as P by | / Discrete Analyser | | | | | | | |
| Total i nospinorus as i | Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.01 | 0.04 | 0.02 | 0.02 |

Page : 7 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | CcS | DdS | A5S | B5S | C5S |
|--------------------------------------|-----------------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|
| | Cl | ient sampli | ng date / time | 19-DEC-2011 15:00 |
| Compound | CAS Number | LOR | Unit | EM1114567-021 | EM1114567-022 | EM1114567-023 | EM1114567-024 | EM1114567-025 |
| EG020T: Total Metals by ICP-MS | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.010 |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.003 | 0.002 | 0.004 | 0.003 |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | <0.005 | <0.005 | <0.005 |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 |
| EG035T: Total Recoverable Mercui | ry by FIMS | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 |
| EK055G: Ammonia as N by Discrete | e Analyser | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.11 | 0.09 | 0.11 | 0.08 | 0.10 |
| EK057G: Nitrite as N by Discrete A | nalyser | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK058G: Nitrate as N by Discrete A | Analyser | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK059G: Nitrite plus Nitrate as N (I | NOx) by Discrete Ana | lyser | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| EK061G: Total Kjeldahl Nitrogen By | y Discrete Analy <u>ser</u> | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| EK062G: Total Nitrogen as N (TKN | + NOx) by Discrete Ar | alyser_ | | | | | | |
| ` Total Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| EK067G: Total Phosphorus as P by | Discrete Analyser | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.04 | 0.05 | 0.09 | 0.03 |
| | | | _ | | 1 | 1 | I | |

Page : 8 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | D5S | A6S | B6S | C6S | D6S | | | |
|---------------------------------------|---------------------------|------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|--|--|--|
| | Cli | ent sampli | ng date / time | 19-DEC-2011 15:00 | | | |
| Compound | CAS Number | LOR | Unit | EM1114567-026 | EM1114567-027 | EM1114567-028 | EM1114567-029 | EM1114567-030 | | | |
| EG020T: Total Metals by ICP-MS | | | | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | | | |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | | |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.010 | | | |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.003 | 0.002 | 0.002 | 0.002 | | | |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | 0.006 | 0.006 | <0.005 | <0.005 | | | |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 | | | |
| EG035T: Total Recoverable Mercury | by FIMS | | | | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | | |
| EK055G: Ammonia as N by Discrete | Analyser | | | | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.10 | 0.11 | 0.11 | 0.10 | | | |
| EK057G: Nitrite as N by Discrete Ana | alyser | | | | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK058G: Nitrate as N by Discrete An | alyser | | | | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK059G: Nitrite plus Nitrate as N (NC | Ox) by Discrete Anal | lyser | | | | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK061G: Total Kjeldahl Nitrogen By I | Discrete Analys <u>er</u> | | | | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.2 | 0.2 | 0.1 | 0.3 | | | |
| EK062G: Total Nitrogen as N (TKN + | NOx) by Discrete An | alyser | | | | | | | | | |
| Total Nitrogen as N | | 0.1 | mg/L | 0.2 | 0.2 | 0.2 | 0.1 | 0.3 | | | |
| EK067G: Total Phosphorus as P by D | Discrete Analyser | | | | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | 0.02 | 0.02 | 0.02 | <0.01 | 0.05 | | | |

Page : 9 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | A1B | B1B | C1B | D1B | A2B | | |
|---------------------------------------|---------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|--|--|
| | CI | ient sampli | ng date / time | 19-DEC-2011 15:00 | | |
| Compound | CAS Number | LOR | Unit | EM1114567-031 | EM1114567-032 | EM1114567-033 | EM1114567-034 | EM1114567-035 | | |
| EG020T: Total Metals by ICP-MS | | | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | | |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.010 | 0.010 | 0.010 | | |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.003 | 0.002 | 0.002 | 0.002 | | |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.013 | 0.009 | <0.005 | <0.005 | 0.006 | | |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 | | |
| EG035T: Total Recoverable Mercury | by FIMS | | | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | |
| EK055G: Ammonia as N by Discrete | Analyser | | | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.12 | 0.11 | 0.09 | 0.12 | | |
| EK057G: Nitrite as N by Discrete Ana | alyser | | | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK058G: Nitrate as N by Discrete An | alyser | | | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK059G: Nitrite plus Nitrate as N (NO | Ox) by Discrete Ana | lyser | | | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK061G: Total Kjeldahl Nitrogen By I | Discrete Analyser | | | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.1 | 0.2 | 0.2 | 0.2 | | |
| EK062G: Total Nitrogen as N (TKN + | NOx) by Discrete Ar | nalyser | | | | | | | | |
| ^ Total Nitrogen as N | | 0.1 | mg/L | 0.1 | 0.1 | 0.2 | 0.2 | 0.2 | | |
| EK067G: Total Phosphorus as P by D | Discrete Analyser | | | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.01 | 0.02 | <0.01 | 0.04 | | |

Page : 10 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Cli | ent sample ID | B2B | C2B | D2B | A3B | B3B | | | |
|-------------------------------------|-----------------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|--|--|--|
| | Cl | ient sampli | ng date / time | 19-DEC-2011 15:00 | | | |
| Compound | CAS Number | LOR | Unit | EM1114567-036 | EM1114567-037 | EM1114567-038 | EM1114567-039 | EM1114567-040 | | | |
| EG020T: Total Metals by ICP-MS | | | | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.006 | 0.006 | 0.006 | 0.006 | | | |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | 0.0001 | <0.0001 | <0.0001 | 0.0001 | <0.0001 | | | |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.010 | 0.011 | 0.010 | 0.010 | | | |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.002 | 0.003 | 0.003 | 0.004 | 0.002 | | | |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | <0.005 | 0.006 | <0.005 | <0.005 | | | |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | <0.05 | <0.05 | <0.05 | | | |
| EG035T: Total Recoverable Mercu | ry by FIMS | | | | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | | |
| EK055G: Ammonia as N by Discret | e Analyser | | | | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.13 | 0.10 | 0.09 | 0.12 | 0.09 | | | |
| EK057G: Nitrite as N by Discrete A | nalyser | | | | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK058G: Nitrate as N by Discrete | Analyser | | | | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK059G: Nitrite plus Nitrate as N (| NOx) by Discrete Ana | lyser | | | | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK061G: Total Kjeldahl Nitrogen B | y Discrete Analy <u>ser</u> | | | | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | 0.2 | 0.1 | 0.1 | 0.2 | | | |
| EK062G: Total Nitrogen as N (TKN | + NOx) by Discrete Ar | nalyser | | | | | | | | | |
| ^ Total Nitrogen as N | | 0.1 | mg/L | <0.1 | 0.2 | 0.1 | 0.1 | 0.2 | | | |
| EK067G: Total Phosphorus as P by | / Discrete Analyser | | | | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | 0.02 | 0.01 | 0.06 | 0.02 | 0.04 | | | |
| | | | - | | | 1 | | 1 | | | |

Page : 11 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Cli | ent sample ID | C3B | D3B | A4B | B4B | C4B | | |
|-----------------------------------|-------------------------|-------------|----------------|-------------------|-------------------|-------------------|-------------------|-------------------|--|--|
| | Cl | ient sampli | ng date / time | 19-DEC-2011 15:00 | | |
| Compound | CAS Number | LOR | Unit | EM1114567-041 | EM1114567-042 | EM1114567-043 | EM1114567-044 | EM1114567-045 | | |
| EG020T: Total Metals by ICP-MS | | | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.005 | 0.006 | 0.006 | 0.006 | 0.006 | | |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.009 | 0.009 | 0.010 | 0.010 | 0.010 | | |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.004 | 0.004 | 0.005 | 0.005 | 0.005 | | |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| Zinc | 7440-66-6 | 0.005 | mg/L | <0.005 | <0.005 | 0.007 | 800.0 | 0.007 | | |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | <0.05 | 0.08 | <0.05 | 0.06 | | |
| EG035T: Total Recoverable Mercu | ury by FIMS | | | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | |
| EK055G: Ammonia as N by Discre | te Analyser | | | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.12 | 0.13 | 0.12 | 0.10 | 0.09 | | |
| EK057G: Nitrite as N by Discrete | Analyser | | | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK058G: Nitrate as N by Discrete | Analyser | | | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK059G: Nitrite plus Nitrate as N | (NOx) by Discrete Ana | lyser | | | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK061G: Total Kjeldahl Nitrogen E | By Discrete Analyser | | | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | <0.1 | 0.1 | <0.1 | | |
| EK062G: Total Nitrogen as N (TKN | I + NOx) by Discrete Ar | nalyser | | | | | | | | |
| ` Total Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | <0.1 | 0.1 | <0.1 | | |
| EK067G: Total Phosphorus as P b | y Discrete Analyser | | | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| | | | | | | | | | | |

Page : 12 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Clie | ent sample ID | D4B | X4B | Y4B | AaB | BbB 19-DEC-2011 15:00 EM1114567-050 <0.001 0.005 <0.001 <0.0001 <0.001 0.010 <0.001 0.005 <0.001 0.006 0.006 <0.001 <0.001 <0.001 <0.001 <0.001 <0.010 <0.001 <0.010 <0.001 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 <0.011 | | |
|---------------------------------------|-----------------------------|------------|----------------|-------------------|-------------------|-------------------|-------------------|---|--|--|
| | Cli | ent sampli | ng date / time | 19-DEC-2011 15:00 | | |
| Compound | CAS Number | LOR | Unit | EM1114567-046 | EM1114567-047 | EM1114567-048 | EM1114567-049 | EM1114567-050 | | |
| EG020T: Total Metals by ICP-MS | | | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.005 | 0.007 | 0.006 | 0.005 | | |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | 0.0001 | <0.0001 | <0.0001 | 0.0001 | <0.0001 | | |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.011 | 0.010 | 0.010 | 0.010 | | |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.005 | 0.005 | 0.004 | 0.004 | | | |
| Lead | 7439-92-1 | 0.001 | mg/L | 0.001 | <0.001 | <0.001 | <0.001 | | | |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.010 | 0.006 | 0.008 | 0.007 | | | |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.06 | 0.06 | 0.10 | 0.06 | | |
| EG035T: Total Recoverable Mercury | by FIMS | | | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | |
| EK055G: Ammonia as N by Discrete | Analyser | | | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.08 | 0.16 | 0.11 | 0.12 | 0.11 | | |
| EK057G: Nitrite as N by Discrete Ana | alyser | | | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK058G: Nitrate as N by Discrete An | alyser | | | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | 0.04 | <0.01 | <0.01 | 0.01 | <0.01 | | |
| EK059G: Nitrite plus Nitrate as N (NC | Ox) by Discrete Ana | lyser | | | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | 0.04 | <0.01 | <0.01 | 0.01 | <0.01 | | |
| EK061G: Total Kjeldahl Nitrogen By [| Discrete Analys <u>er</u> | | | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | <0.1 | 0.1 | <0.1 | | |
| EK062G: Total Nitrogen as N (TKN + I | NOx) by Discre <u>te An</u> | alyser_ | | | | | | | | |
| Total Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | <0.1 | 0.1 | <0.1 | | |
| EK067G: Total Phosphorus as P by D | Discrete Analys <u>er</u> | | | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | <0.01 | 0.02 | <0.01 | 0.03 | <0.01 | | |

Page : 13 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Cli | ent sample ID | СсВ | DdB | A5B | B5B | C5B | | |
|-------------------------------------|-----------------------------|-------------|-----------------|-------------------|-------------------|-------------------|-------------------|-------------------|--|--|
| | Cl | ient sampli | ing date / time | 19-DEC-2011 15:00 | | |
| Compound | CAS Number | LOR | Unit | EM1114567-051 | EM1114567-052 | EM1114567-053 | EM1114567-054 | EM1114567-055 | | |
| EG020T: Total Metals by ICP-MS | | | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.006 | 0.007 | 0.005 | 0.005 | 0.006 | | |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0001 | <0.0001 | <0.0001 | | |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.011 | 0.010 | 0.011 | 0.010 | | |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | 0.001 | <0.001 | <0.001 | <0.001 | | |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.005 | 0.005 | 0.005 | 0.005 | 0.004 | | |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | |
| Vanadium | 7440-62-2 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | 0.01 | <0.01 | | |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | <0.005 | 0.011 | 0.006 | 0.006 | | |
| Iron | 7439-89-6 | 0.05 | mg/L | <0.05 | 0.09 | 0.17 | 0.14 | 0.12 | | |
| EG035T: Total Recoverable Mercu | iry by FIMS | | | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | |
| EK055G: Ammonia as N by Discre | te Analyser | | | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.10 | 0.13 | 0.08 | 0.12 | 0.10 | | |
| EK057G: Nitrite as N by Discrete | Analyser | | | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK058G: Nitrate as N by Discrete | Analyser | | | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK059G: Nitrite plus Nitrate as N (| (NOx) by Discrete Ana | lyser | | | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | |
| EK061G: Total Kjeldahl Nitrogen B | By Discrete Analyser | | | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | 0.1 | <0.1 | 0.2 | 1.0 | | |
| EK062G: Total Nitrogen as N (TKN | l + NOx) by Discrete Ar | nalyser | | | | | | | | |
| ↑ Total Nitrogen as N | | 0.1 | mg/L | <0.1 | 0.1 | <0.1 | 0.2 | 1.0 | | |
| EK067G: Total Phosphorus as P b | y Discrete Analy <u>ser</u> | | | | | | | | | |
| Total Phosphorus as P | | 0.01 | mg/L | 0.02 | 0.14 | 0.02 | 0.05 | 0.05 | | |
| | | | | | | | | | | |

Page : 14 of 14

Work Order : EM1114567 Amendment 2
Client : SINCLAIR KNIGHT MERZ

Project : VE23610 609



| Sub-Matrix: WATER | | Cli | ent sample ID | D5B | A6B | B6B | C6B | D6B | | | |
|-----------------------------------|-------------------------|-------------|-----------------|-------------------|-------------------|-------------------|-------------------|-------------------|--|--|--|
| | CI | ient sampli | ing date / time | 19-DEC-2011 15:00 | | | |
| Compound | CAS Number | LOR | Unit | EM1114567-056 | EM1114567-057 | EM1114567-058 | EM1114567-059 | EM1114567-060 | | | |
| EG020T: Total Metals by ICP-MS | | | | | | | | | | | |
| Arsenic | 7440-38-2 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Barium | 7440-39-3 | 0.001 | mg/L | 0.007 | 0.005 | 0.006 | 0.005 | 0.006 | | | |
| Beryllium | 7440-41-7 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Cadmium | 7440-43-9 | 0.0001 | mg/L | <0.0001 | <0.0001 | 0.0001 | 0.0001 | <0.0001 | | | |
| Cobalt | 7440-48-4 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Chromium | 7440-47-3 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Copper | 7440-50-8 | 0.001 | mg/L | 0.010 | 0.012 | 0.011 | 0.010 | 0.012 | | | |
| Manganese | 7439-96-5 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | | | |
| Nickel | 7440-02-0 | 0.001 | mg/L | 0.006 | 0.005 | 0.005 | 0.005 | 0.006 | | | |
| Lead | 7439-92-1 | 0.001 | mg/L | <0.001 | <0.001 | <0.001 | <0.001 | 0.001 | | | |
| Vanadium | 7440-62-2 | 0.01 | mg/L | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | | | |
| Zinc | 7440-66-6 | 0.005 | mg/L | 0.006 | 0.025 | 800.0 | 0.009 | 0.014 | | | |
| Iron | 7439-89-6 | 0.05 | mg/L | 0.10 | 0.16 | 0.17 | 0.16 | 0.14 | | | |
| EG035T: Total Recoverable Mercu | ury by FIMS | | | | | | | | | | |
| Mercury | 7439-97-6 | 0.0001 | mg/L | <0.0001 | <0.0001 | <0.0001 | <0.0001 | <0.0001 | | | |
| EK055G: Ammonia as N by Discre | te Analyser | | | | | | | | | | |
| Ammonia as N | 7664-41-7 | 0.01 | mg/L | 0.09 | 0.08 | 0.08 | 0.11 | 0.10 | | | |
| EK057G: Nitrite as N by Discrete | Analyser | | | | | | | | | | |
| Nitrite as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK058G: Nitrate as N by Discrete | Analyser | | | | | | | | | | |
| Nitrate as N | 14797-55-8 | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK059G: Nitrite plus Nitrate as N | (NOx) by Discrete Ana | lyser | | | | | | | | | |
| Nitrite + Nitrate as N | | 0.01 | mg/L | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | | | |
| EK061G: Total Kjeldahl Nitrogen B | By Discrete Analyser | | | | | | | | | | |
| Total Kjeldahl Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | 0.1 | 0.1 | <0.1 | | | |
| EK062G: Total Nitrogen as N (TKN | I + NOx) by Discrete Ar | nalyser | | | | | | | | | |
| ^ Total Nitrogen as N | | 0.1 | mg/L | <0.1 | <0.1 | 0.1 | 0.1 | <0.1 | | | |
| EK067G: Total Phosphorus as P b | v Discrete Analyser | | | | | | | 1 | | | |
| Total Phosphorus as P | | 0.01 | mg/L | 0.04 | 0.02 | 0.04 | 0.04 | 0.02 | | | |
| . 3.aoopiioi ao ao i | | | 5 | | 1 1 1 | 1 1 1 | 1.1 | 1 1 1 | | | |



Appendix E Infauna Sampling Results



Client
Contact
Location
Collection Date

SKM - Adelaide Greg Barbara Unknown 19/12/2011

Taonomist
Id Date

Shona Hocknull 05/02/2012

| Phylum | Class/order | Family | Species | A1 | A4 | A5 | Α6 | B1 | B1 | В4 | В5 | В6 | C1 | C4 | C5 | C6 | D5 | D6 | DI |
|-----------------------|-------------|-----------------|-----------------|-----------|----|----|----|----|----|----|----|----|-----------|----|-----------|----|----|----|----|
| Annelida | Polychaeta | Nephtyidae | | 1 | | 1 | | 1 | | | | | | | | | | | |
| Annelida | Polychaeta | Opheliidae | Armandia sp.1 | | | | | | | | | 1 | | | | | | | |
| Annelida | Polychaeta | Syllidae | | | 1 | | 1 | | | | | | | | | | | | |
| Annelida | Polychaeta | Terebellidae | | | | | | | | | | | | | | | | | 1 |
| Arthropoda/ Crustacea | Amphipoda | Cheirocratidae | | | 1 | 1 | 1 | | | | | 3 | | | | | | | |
| Arthropoda/ Crustacea | Amphipoda | Damaged | | | | 1 | | | | | | | | | | | | | |
| Arthropoda/ Crustacea | Amphipoda | Eusiridae | | | | | 2 | | | | | | | | | | | | |
| Arthropoda/ Crustacea | Amphipoda | Lysianassidae | | | | | | | | | | 1 | | | | | | | |
| Arthropoda/ Crustacea | Amphipoda | Stegocephalidae | | | | | | 1 | | | | | | | | | | | 1 |
| Arthropoda/ Crustacea | Anaspidacea | | | | 3 | | | | | | | | | | | | | | |
| Arthropoda/ Crustacea | Asellota | | | | | | 2 | | | | | | | | | | | | |
| Arthropoda/ Crustacea | Isopoda | | | | | 1 | 3 | | | | | | | | | | | | |
| Arthropoda/ Crustacea | Leptostraca | | | | | | | | | | | 2 | | | | | | | |
| Arthropoda/ Crustacea | Tanaidacea | Kalliapseudidae | | | 1 | | | | | | | | | | | | | | |
| Cnideria | Actiniaria | | | | | | | | | | | | | 1 | | | | | |
| Echinodermata | Asteroidea | Asteroidea | | | | | | | | | | 1 | | | | | | | |
| Echinodermata | Echinoidea | Echinoidea | | | | | | | | | | | | | 1 | | | | |
| Mollusca | Bivalvia | Chamidae | | 1 | | | | | | | | | | | | | 1 | | |
| Mollusca | Bivalvia | Limidae | | 1 | | | | | | | | | | | | | | | |
| Mollusca | Bivalvia | Littorinidae | | | | | 2 | | | | | | | | | | | | |
| Mollusca | Bivalvia | Nuculanidae | Nuculana sp.1 | | | | | | | 2 | 1 | | | | | | | | 1 |
| Mollusca | Bivalvia | Tellinidae | Tellina sp.1 | | | | | 2 | | | | | 1 | | 1 | | | | |
| Mollusca | Bivalvia | Thyasiridae | Tauraxinus sp.1 | | | | 1 | 1 | | | | | | | | | | | |
| Mollusca | Bivalvia | Veneridae | | | | | 1 | | | | | | | | | | | | |
| Mollusca | Gastropoda | Patellidae | | | | | 1 | | | | | 1 | | | | | | | |
| Mollusca | Gastropoda | Turbinidae | Astralium sp.1 | | | | | | | | | 1 | | | | | | | |
| Mollusca | Gastropoda | Volutidae | | | | | | 1 | | | | | | | | | | | |
| Nemertea | Nemertea | Nemertea | | | | | | | | | | 1 | | | | | | | |



Appendix F EPBC Search Result



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about <u>Environment Assessments</u> and the EPBC Act including significance guidelines, forms and application process details.

Report created: 13/11/13 18:26:01


Summary

Details

Matters of NES
Other Matters Protected by the EPBC Act
Extra Information


Caveat

<u>Acknowledgements</u>



This map may contain data which are ©Commonwealth of Australia (Geoscience Australia), ©PSMA 2010

Coordinates
Buffer: 10.0Km



Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the <u>Administrative Guidelines on Significance</u>.

| World Heritage Properties: | None |
|---|------|
| National Heritage Places: | None |
| Wetlands of International Importance: | None |
| Great Barrier Reef Marine Park: | None |
| Commonwealth Marine Areas: | None |
| Listed Threatened Ecological Communities: | None |
| Listed Threatened Species: | 34 |
| Listed Migratory Species: | 33 |

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage-values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place and the heritage values of a place on the Register of the National Estate.

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

A <u>permit</u> may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

| Commonwealth Land: | None |
|------------------------------------|------|
| Commonwealth Heritage Places: | None |
| Listed Marine Species: | 62 |
| Whales and Other Cetaceans: | 12 |
| Critical Habitats: | None |
| Commonwealth Reserves Terrestrial: | None |
| Commonwealth Reserves Marine | None |

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

| Place on the RNE: | 1 |
|----------------------------------|------|
| State and Territory Reserves: | 1 |
| Regional Forest Agreements: | None |
| Invasive Species: | 17 |
| Nationally Important Wetlands: | None |
| Key Ecological Features (Marine) | None |

Details

Matters of National Environmental Significance

| Listed Threatened Species | | [Resource Information |
|--|--------------|--|
| Name | Status | Type of Presence |
| Birds | | |
| Acanthiza iredalei iredalei | | |
| Slender-billed Thornbill (western) [25967] | Vulnerable | Species or species habitat likely to occur within area |
| Diomedea epomophora epomophora | | |
| Southern Royal Albatross [25996] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea epomophora sanfordi | | |
| Northern Royal Albatross [82331] | Endangered | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea exulans antipodensis | | |
| Antipodean Albatross [82269] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea exulans exulans | En den maned | 0 |
| Tristan Albatross [82337] | Endangered | Species or species habitat may occur within area |
| <u>Diomedea exulans (sensu lato)</u> | | |
| Wandering Albatross [1073] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Halobaena caerulea | | |
| Blue Petrel [1059] Leipoa ocellata | Vulnerable | Species or species habitat may occur within area |
| · · · · · · · · · · · · · · · · · · · | Vulnerable | Species or species |
| Malleefowl [934] | vuirierable | Species or species habitat likely to occur within area |
| Macronectes giganteus | | |
| Southern Giant-Petrel [1060] | Endangered | Species or species habitat may occur within area |

| Name | Status | Type of Presence |
|--|-----------------|--|
| Macronectes halli | | |
| Northern Giant-Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Psophodes nigrogularis leucogaster | | |
| Western Whipbird (eastern) [64448] | Vulnerable | Species or species habitat likely to occur within area |
| Pterodroma mollis | | |
| Soft-plumaged Petrel [1036] | Vulnerable | Species or species habitat may occur within area |
| Rostratula australis | | |
| Australian Painted Snipe [77037] | Endangered | Species or species habitat may occur within area |
| Sternula nereis nereis | | |
| Australian Fairy Tern [82950] | Vulnerable | Species or species habitat likely to occur within area |
| Thalassarche cauta cauta | | |
| Shy Albatross, Tasmanian Shy Albatross [82345] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche cauta steadi | | _ |
| White-capped Albatross [82344] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche melanophris | | |
| Black-browed Albatross [66472] | Vulnerable | Species or species habitat may occur within area |
| Thalassarche melanophris impavida Campbell Albatross [82449] | Vulnerable | Species or species habitat may occur within |
| Mammals | | area |
| | | |
| Balaenoptera musculus Blue Whale [36] | Endangered | Species or species habitat may occur within area |
| Eubalaena australis | | |
| Southern Right Whale [40] | Endangered | Breeding known to occur within area |
| Megaptera novaeangliae | V. de a rala la | Canaina ar annaina |
| Humpback Whale [38] Neophoca cinerea | Vulnerable | Species or species habitat likely to occur within area |
| Australian Sea-lion [22] | Vulnerable | Foraging, feeding or |
| Sminthopsis psammophila | 7 4.11014010 | related behaviour likely to occur within area |
| Sandhill Dunnart [291] | Endangered | Species or species habitat likely to occur within area |
| Plants | | within area |
| Acacia enterocarpa | | |
| Jumping-jack Wattle [17615] | Endangered | Species or species habitat likely to occur within area |
| Acacia pinguifolia Fat-leaved Wattle [5319] | Endangered | Species or species habitat may occur within area |
| Caladenia tensa Greencomb Spider-orchid, Rigid Spider-orchid [24390] | Endangered | Species or species habitat likely to occur within area |
| Frankenia plicata | | |
| [4225] | Endangered | Species or species habitat likely to occur within area |

| Name | Status | Type of Presence |
|--|---------------------------|--|
| Prostanthera calycina West Coast Mintbush, Limestone Mintbush, Red Mintbush [9470] | Vulnerable | Species or species habitat likely to occur within area |
| Ptilotus beckerianus Ironstone Mulla Mulla [3787] | Vulnerable | Species or species habitat may occur within area |
| Pultenaea trichophylla Tufted Bush-pea [12715] | Endangered | Species or species habitat may occur within area |
| Reptiles | | 4.04 |
| Caretta caretta Loggerhead Turtle [1763] | Endangered | Breeding likely to occur within area |
| Chelonia mydas Green Turtle [1765] | Vulnerable | Foraging, feeding or related behaviour known to occur within area |
| Dermochelys coriacea Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Foraging, feeding or related behaviour known to occur within area |
| Sharks | | |
| Carcharodon carcharias Great White Shark [64470] | Vulnerable | Species or species habitat may occur within area |
| Listed Migratory Species * Species is listed under a different scientific name on | the EPBC Act - Threatened | [Resource Information Species list. |
| Name | Threatened | Type of Presence |
| Migratory Marine Birds | | |
| Apus pacificus Fork-tailed Swift [678] | | Species or species habitat likely to occur within area |
| Diomedea antipodensis Antipodean Albatross [64458] | Vulnerable* | Foraging, feeding or related behaviour likely to occur within area |
| <u>Diomedea dabbenena</u> Tristan Albatross [66471] | Endangered* | Species or species habitat may occur within area |
| Diomedea epomophora (sensu stricto) Southern Royal Albatross [1072] | Vulnerable* | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea exulans (sensu lato) Wandering Albatross [1073] | Vulnerable | Foraging, feeding or related behaviour likely to occur within area |
| Diomedea sanfordi Northern Royal Albatross [64456] | Endangered* | Foraging, feeding or related behaviour likely to occur within area |
| Macronectes giganteus Southern Giant-Petrel [1060] | Endangered | Species or species habitat may occur within area |
| Macronectes halli Northern Giant-Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Puffinus carneipes Flesh-footed Shearwater, Fleshy-footed Shearwater [1043] | | Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche cauta (sensu stricto) Shy Albatross, Tasmanian Shy Albatross [64697] | Vulnerable* | Foraging, feeding or |

| Name | Threatened | Type of Presence |
|---|--------------|--|
| | | related behaviour likely to occur within area |
| Thalassarche impavida | | to occur within area |
| Campbell Albatross [64459] | Vulnerable* | Species or species habitat may occur within |
| Thologographo molonophric | | area |
| Thalassarche melanophris Black-browed Albatross [66472] | Vulnerable | Species or species |
| | | habitat may occur within area |
| Thalassarche steadi White capped Alberross [64462] | Vulnerable* | Foraging fooding or |
| White-capped Albatross [64462] | Vullierable | Foraging, feeding or related behaviour likely to occur within area |
| Migratory Marine Species | | |
| Balaenoptera edeni Bryde's Whale [35] | | Species or species |
| | | habitat may occur within |
| Balaenoptera musculus | | area |
| Blue Whale [36] | Endangered | Species or species |
| | | habitat may occur within area |
| Caperea marginata Pygmy Pight Whale [20] | | Species or species |
| Pygmy Right Whale [39] | | Species or species habitat may occur within |
| Carcharodon carcharias | | area |
| Great White Shark [64470] | Vulnerable | Species or species |
| | | habitat may occur within area |
| Caretta caretta | | |
| Loggerhead Turtle [1763] | Endangered | Breeding likely to occur within area |
| Chelonia mydas | Mala analala | |
| Green Turtle [1765] | Vulnerable | Foraging, feeding or related behaviour known |
| <u>Dermochelys coriacea</u> | | to occur within area |
| Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Foraging, feeding or |
| | | related behaviour known to occur within area |
| Eubalaena australis | | to occur within area |
| Southern Right Whale [40] | Endangered | Breeding known to occur within area |
| <u>Lagenorhynchus obscurus</u> | | |
| Dusky Dolphin [43] | | Species or species habitat may occur within |
| Lampa pagus | | area |
| <u>Lamna nasus</u> Porbeagle, Mackerel Shark [83288] | | Species or species |
| | | habitat likely to occur within area |
| Megaptera novaeangliae | | within area |
| Humpback Whale [38] | Vulnerable | Species or species habitat likely to occur |
| Oreinus area | | within area |
| Orcinus orca Killer Whale, Orca [46] | | Species or species |
| | | habitat may occur within area |
| Migratory Terrestrial Species | | area |
| Haliaeetus leucogaster White-bellied Sea-Eagle [943] | | Species or species |
| vvinte semes ees Esgie [e ie] | | habitat known to occur |
| <u>Leipoa ocellata</u> | | within area |
| Malleefowl [934] | Vulnerable | Species or species |
| | | habitat likely to occur within area |
| Merops ornatus Painhow Receptor [670] | | Charles ar anacias |
| Rainbow Bee-eater [670] | | Species or species habitat may occur within |
| | | |

| name | rnreatened | Type of Presence |
|---|--------------|--|
| | | area |
| Migratory Wetlands Species | | |
| <u>Ardea alba</u> | | |
| Great Egret, White Egret [59541] | | Species or species |
| | | habitat likely to occur |
| Ardon ibis | | within area |
| Ardea ibis Cottle Faret [50542] | | Charles or angeles |
| Cattle Egret [59542] | | Species or species |
| | | habitat likely to occur within area |
| Charadrius veredus | | within area |
| Oriental Plover, Oriental Dotterel [882] | | Species or species |
| Chemai i lover, Chemai Botterer [002] | | habitat may occur within |
| | | area |
| Gallinago hardwickii | | 55 5. |
| Latham's Snipe, Japanese Snipe [863] | | Species or species |
| | | habitat may occur within |
| | | area |
| Rostratula benghalensis (sensu lato) | | |
| Painted Snipe [889] | Endangered* | Species or species |
| | | habitat may occur within |
| | | area |
| | | |
| Other Matters Protected by the EPBC A | ct | |
| Listed Marine Chasins | | |
| Listed Marine Species | | [Resource Information |
| * Species is listed under a different scientific name | | · · · · · · |
| Name | Threatened | Type of Presence |
| Birds | | |
| Apus pacificus | | |
| Fork-tailed Swift [678] | | Species or species |
| | | habitat likely to occur |
| | | within area |
| Ardea alba | | |
| Great Egret, White Egret [59541] | | Species or species |
| | | habitat likely to occur |
| Ardea ibis | | within area |
| | | Species or species |
| Cattle Egret [59542] | | Species or species habitat likely to occur |
| | | within area |
| <u>Catharacta skua</u> | | within area |
| Great Skua [59472] | | Species or species |
| | | habitat may occur within |
| | | area |
| <u>Charadrius veredus</u> | | |
| Oriental Plover, Oriental Dotterel [882] | | Species or species |
| | | habitat may occur within |
| | | area |
| <u>Diomedea antipodensis</u> | | |
| Antipodean Albatross [64458] | Vulnerable* | Foraging, feeding or |
| | | related behaviour likely |
| Diamodos deblesis sus | | to occur within area |
| <u>Diomedea dabbenena</u> | | |
| Tristan Albatross [66471] | Endangered* | Species or species |
| | | habitat may occur within |
| Diomedea epomophora (sensu stricto) | | area |
| | Vulnerable* | Foreging fooding or |
| Southern Royal Albatross [1072] | vumerable | Foraging, feeding or |
| | | related behaviour likely to occur within area |
| Diomedea exulans (sensu lato) | | to occur within area |
| Wandering Albatross [1073] | Vulnerable | Foraging, feeding or |
| | v diriorabio | related behaviour likely |
| | | to occur within area |
| Diomedea sanfordi | | |
| Northern Royal Albatross [64456] | Endangered* | Foraging, feeding or |
| | - | related behaviour likely |
| | | |
| | | |

Threatened

Name

Type of Presence

| Name | Threatened | Type of Presence |
|---|-------------|---|
| | | to occur within area |
| Eudyptula minor | | |
| Little Penguin [1085] | | Breeding known to occur within area |
| Gallinago hardwickii Latham's Snipe, Japanese Snipe [863] | | Species or species habitat may occur within area |
| Haliaeetus leucogaster White-bellied Sea-Eagle [943] | | Species or species habitat known to occur within area |
| Halobaena caerulea Blue Petrel [1059] | Vulnerable | Species or species habitat may occur within area |
| Larus novaehollandiae Silver Gull [810] | | Breeding known to occur within area |
| Macronectes giganteus Southern Giant-Petrel [1060] | Endangered | Species or species habitat may occur within area |
| Macronectes halli | | |
| Northern Giant-Petrel [1061] | Vulnerable | Species or species habitat may occur within area |
| Merops ornatus Rainbow Bee-eater [670] | | Species or species habitat may occur within area |
| Pandion haliaetus Osprey [952] | | Species or species habitat known to occur within area |
| Phalacrocorax fuscescens Black-faced Cormorant [59660] | | Breeding known to occur within area |
| Pterodroma mollis Soft-plumaged Petrel [1036] | Vulnerable | Species or species habitat may occur within area |
| Puffinus carneipes Flesh-footed Shearwater, Fleshy-footed Shearwater [1043] | | Foraging, feeding or related behaviour likely to occur within area |
| Rostratula benghalensis (sensu lato) Painted Snipe [889] | Endangered* | Species or species habitat may occur within area |
| Sterna bergii Crested Tern [816] | | Breeding known to occur within area |
| Sterna fuscata Sooty Tern [794] | | Breeding known to occur within area |
| Sterna nereis Fairy Tern [796] | | Breeding known to occur |
| Thalassarche cauta (sensu stricto) Shy Albatross, Tasmanian Shy Albatross [64697] | Vulnerable* | within area Foraging, feeding or related behaviour likely to occur within area |
| Thalassarche impavida Campbell Albatross [64459] | Vulnerable* | Species or species habitat may occur within area |
| Thalassarche melanophris Black-browed Albatross [66472] | Vulnerable | Species or species habitat may occur within area |
| Thalassarche steadi White-capped Albatross [64462] | Vulnerable* | Foraging, feeding or |

| Name | Threatened | Type of Presence |
|---|------------|--|
| | | related behaviour likely |
| This are is an abricallia, an abricallia | | to occur within area |
| Thinornis rubricollis rubricollis Hooded Plover (eastern) [66726] | | Species or species habitat known to occur within area |
| Fish | | |
| Acentronura australe | | |
| Southern Pygmy Pipehorse [66185] | | Species or species habitat may occur within area |
| Campichthys tryoni Tryon's Pipefish [66193] | | Species or species |
| | | habitat may occur within area |
| Filicampus tigris Tiger Pipefish [66217] | | Species or species |
| | | habitat may occur within area |
| Heraldia nocturna Upside-down Pipefish, Eastern Upside-down | | Species or species |
| Pipefish, Eastern Upside-down Pipefish [66227] | | habitat may occur within area |
| Hippocampus breviceps Chart has all Casharras Chart assessed Casharras | | On a sing an an asing |
| Short-head Seahorse, Short-snouted Seahorse [66235] | | Species or species habitat may occur within area |
| Histiogamphelus cristatus | | |
| Rhino Pipefish, Macleay's Crested Pipefish, Ringback Pipefish [66243] | | Species or species habitat may occur within area |
| Hypselognathus rostratus | | 0 |
| Knifesnout Pipefish, Knife-snouted Pipefish [66245] | | Species or species habitat may occur within area |
| Kaupus costatus | | |
| Deepbody Pipefish, Deep-bodied Pipefish [66246] | | Species or species habitat may occur within area |
| <u>Leptoichthys fistularius</u> Brushtail Pipefish [66248] | | Species or species |
| | | habitat may occur within area |
| Lissocampus caudalis Australian Connette Dinefich | | Charles ar anasias |
| Australian Smooth Pipefish, Smooth Pipefish [66249] | | Species or species habitat may occur within area |
| Lissocampus runa | | Consiss or opening |
| Javelin Pipefish [66251] | | Species or species habitat may occur within area |
| Maroubra perserrata | | 0 |
| Sawtooth Pipefish [66252] | | Species or species habitat may occur within area |
| Notiocampus ruber | | 0 |
| Red Pipefish [66265] | | Species or species habitat may occur within area |
| Phycodurus eques Leafy Seadragon [66267] | | Species or species |
| | | habitat may occur within area |
| Phyllopteryx taeniolatus Common Seadragon, Weedy Seadragon [66268] | | Species or species |

Species or species

Species or species

Species or species

area

area

habitat may occur within

habitat may occur within

Common Seadragon, Weedy Seadragon [66268]

Pugnose Pipefish, Pug-nosed Pipefish [66269]

Robust Pipehorse, Robust Spiny Pipehorse

Pugnaso curtirostris

Solegnathus robustus

| Name | Threatened | Type of Presence |
|--|------------------|--|
| [66274] | | habitat may occur within |
| r | | area |
| Stigmatopora argus | | |
| Spotted Pipefish, Gulf Pipefish [66276] | | Species or species |
| | | habitat may occur within |
| | | area |
| Stigmatopora nigra | | |
| Widebody Pipefish, Wide-bodied Pipefish, Black | | Species or species |
| Pipefish [66277] | | habitat may occur within |
| Stipecampus cristatus | | area |
| Ringback Pipefish, Ring-backed Pipefish [66278] | | Species or species |
| rangback ripensii, rang backed ripensii [66276] | | habitat may occur within |
| | | area |
| <u>Urocampus carinirostris</u> | | |
| Hairy Pipefish [66282] | | Species or species |
| | | habitat may occur within |
| | | area |
| Vanacampus margaritifer | | |
| Mother-of-pearl Pipefish [66283] | | Species or species |
| | | habitat may occur within |
| Vanacampus phillipi | | area |
| Port Phillip Pipefish [66284] | | Species or species |
| Fort Frillip Fiperish [00204] | | habitat may occur within |
| | | area |
| Vanacampus poecilolaemus | | <u></u> |
| Longsnout Pipefish, Australian Long-snout | | Species or species |
| Pipefish, Long-snouted Pipefish [66285] | | habitat may occur within |
| | | area |
| <u>Vanacampus vercoi</u> | | |
| Verco's Pipefish [66286] | | Species or species |
| | | habitat may occur within |
| Mammals | | area |
| Arctocephalus forsteri | | |
| New Zealand Fur-seal [20] | | Species or species |
| | | habitat may occur within |
| | | area |
| Arctocephalus pusillus | | |
| Australian Fur-seal, Australo-African Fur-seal | | Species or species |
| [21] | | habitat may occur within |
| | | area |
| Neophoca cinerea | | - · · · · |
| Australian Sea-lion [22] | Vulnerable | Foraging, feeding or |
| | | related behaviour likely to occur within area |
| Reptiles | | to occur within area |
| Caretta caretta | | |
| Loggerhead Turtle [1763] | Endangered | Breeding likely to occur |
| | J | within area |
| <u>Chelonia mydas</u> | | |
| Green Turtle [1765] | Vulnerable | Foraging, feeding or |
| | | related behaviour known |
| Dawn a also live as also se | | to occur within area |
| Dermochelys coriacea | Final and second | Fanonina Garati |
| Leatherback Turtle, Leathery Turtle, Luth [1768] | Endangered | Foraging, feeding or |
| | | related behaviour known to occur within area |
| | | to occur within area |
| Whales and other Cetaceans | | [Resource Information] |
| Name | Status | Type of Presence |
| Mammals | | |
| Balaenoptera acutorostrata | | |
| Minke Whale [33] | | Species or species |
| | | habitat may occur within |
| | | area |
| Delegandes eden: | | |
| Balaenoptera edeni | | |
| Bryde's Whale [35] | | Species or species |
| • | | habitat may occur within |
| • | | |

| | 01.1 | T (D |
|---|------------|--|
| Name | Status | Type of Presence |
| Balaenoptera musculus Blue Whale [36] | Endangered | Species or species habitat may occur within area |
| Caperea marginata | | |
| Pygmy Right Whale [39] | | Species or species habitat may occur within area |
| <u>Delphinus delphis</u> | | |
| Common Dophin, Short-beaked Common Dolphin [60] | | Species or species habitat may occur within area |
| Eubalaena australis | | |
| Southern Right Whale [40] | Endangered | Breeding known to occur within area |
| Grampus griseus | | |
| Risso's Dolphin, Grampus [64] | | Species or species habitat may occur within area |
| <u>Lagenorhynchus obscurus</u> | | |
| Dusky Dolphin [43] | | Species or species habitat may occur within area |
| Megaptera novaeangliae | | |
| Humpback Whale [38] | Vulnerable | Species or species habitat likely to occur within area |
| Orcinus orca | | |
| Killer Whale, Orca [46] | | Species or species habitat may occur within area |
| <u>Tursiops aduncus</u> | | |
| Indian Ocean Bottlenose Dolphin, Spotted Bottlenose Dolphin [68418] | | Species or species habitat likely to occur within area |
| Tursiops truncatus s. str. | | |
| Bottlenose Dolphin [68417] | | Species or species habitat may occur within area |

Extra Information

| Places on the RNE | | <u>[Resource miormation]</u> |
|---|-------|--------------------------------|
| Note that not all Indigenous sites may be listed. | | |
| Name | State | Status |
| Natural | | |
| Lipson Island Conservation Park | SA | Registered |
| State and Territory Reserves | | [Resource Information] |
| Name | | State |
| Lipson Island | | SA |

[Resource Information]

Invasive Species [Resource Information]

Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resouces Audit, 2001.

| Name | Status | Type of Presence |
|-----------------|--------|--------------------|
| Birds | | |
| Alauda arvensis | | |
| Skylark [656] | | Species or species |

| Name | Status | Type of Presence |
|--|--------|--|
| | | habitat likely to occur within area |
| Carduelis carduelis | | within area |
| European Goldfinch [403] | | Species or species habitat likely to occur within area |
| Columba livia Rock Pigeon, Rock Dove, Domestic Pigeon [803] | | Species or species |
| | | habitat likely to occur within area |
| Passer domesticus House Sparrow [405] | | Species or species |
| | | habitat likely to occur within area |
| Streptopelia chinensis Spotted Turtle-Dove [780] | | Species or species |
| | | habitat likely to occur within area |
| Sturnus vulgaris Common Starling [389] | | Species or species |
| Turdus merula | | habitat likely to occur within area |
| Common Blackbird, Eurasian Blackbird [596] | | Species or species |
| | | habitat likely to occur within area |
| Mammals <u>Felis catus</u> | | |
| Cat, House Cat, Domestic Cat [19] | | Species or species |
| Muo muoquluo | | habitat likely to occur within area |
| Mus musculus House Mouse [120] | | Species or species |
| | | habitat likely to occur within area |
| Oryctolagus cuniculus Rabbit, European Rabbit [128] | | Species or species |
| Vulpes vulpes | | habitat likely to occur within area |
| Red Fox, Fox [18] | | Species or species |
| Plants | | habitat likely to occur within area |
| Asparagus asparagoides | | |
| Bridal Creeper, Bridal Veil Creeper, Smilax, Florist's Smilax, Smilax Asparagus [22473] | | Species or species habitat likely to occur within area |
| Chrysanthemoides monilifera Bitou Bush, Boneseed [18983] | | Species or species |
| | | habitat may occur within area |
| Lycium ferocissimum African Boxthorn, Boxthorn [19235] | | Species or species |
| | | habitat likely to occur within area |
| Olea europaea Olive, Common Olive [9160] | | Species or species |
| | | habitat may occur within area |
| Rubus fruticosus aggregate Blackberry, European Blackberry [68406] | | Species or species |
| | | habitat likely to occur within area |
| <u>Ulex europaeus</u> Gorse, Furze [7693] | | Species or species |
| | | habitat likely to occur within area |
| | | |

Coordinates

-34.18179 136.323,-34.19738 136.32383

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World Heritage and Register of National Estate properties, Wetlands of International Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the qualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

For species where the distributions are well known, maps are digitised from sources such as recovery plans and detailed habitat studies. Where appropriate, core breeding, foraging and roosting areas are indicated under 'type of presence'. For species whose distributions are less well known, point locations are collated from government wildlife authorities, museums, and non-government organisations; bioclimatic distribution models are generated and these validated by experts. In some cases, the distribution maps are based solely on expert knowledge.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- -Department of Environment, Climate Change and Water, New South Wales
- -Department of Sustainability and Environment, Victoria
- -Department of Primary Industries, Parks, Water and Environment, Tasmania
- -Department of Environment and Natural Resources, South Australia
- -Parks and Wildlife Service NT, NT Dept of Natural Resources, Environment and the Arts
- -Environmental and Resource Management, Queensland
- -Department of Environment and Conservation, Western Australia
- -Department of the Environment, Climate Change, Energy and Water
- -Birds Australia
- -Australian Bird and Bat Banding Scheme
- -Australian National Wildlife Collection
- -Natural history museums of Australia
- -Museum Victoria
- -Australian Museum
- -SA Museum
- -Queensland Museum
- -Online Zoological Collections of Australian Museums
- -Queensland Herbarium
- -National Herbarium of NSW
- -Royal Botanic Gardens and National Herbarium of Victoria
- -Tasmanian Herbarium
- -State Herbarium of South Australia
- -Northern Territory Herbarium
- -Western Australian Herbarium
- -Australian National Herbarium, Atherton and Canberra
- -University of New England
- -Ocean Biogeographic Information System
- -Australian Government, Department of Defence
- -State Forests of NSW
- -Geoscience Australia
- -CSIRO
- -Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the Contact Us page.

© Commonwealth of Australia

Department of Sustainability, Environment, Water, Population and Communities

GPO Box 787

Canberra ACT 2601 Australia

+61 2 6274 1111